

Justicia . Cal. simpliciter vel duplex . Cor. irregularis vel
Subaequalis . Caps: 2-valvis, 2-loocularis: Dissepimen
~~valvis~~ ^{valvis} Contraria in ~~analis~~ retinaculi semi numeri nuncii
natis Gestr.

paniflora L. t. p. 2.

Justicia

Paps: bilobularis, bivalvis: valvulis elastice desilientibus.
Dissegi mentis contrariae, Petaliae ad eis seminum uniuersale

Justicia

Pal. simplex vel duplex. Cor. irregulans. Paps: 2-valvis,
2-locularis: valvulis elastice desilientibus: Dissegi
mentis contrariae ad eam.

Cupulina ♀

Gastera Cupulina specie novati Verminalibus pustulibus, brachia pubes
datis imbreatis ciliato vellopi calyce majoribus La Mart.

T. Enc: pag 40

Sloan hist. I. tab. 109. fig. 1.
Habitat in Madagasca.

Muallia Blechum.

non vid.

~~Galappopolea Gustavii pauciflora umbellae pedunculatis dichotomis, pedicelli
lateralibus in floris, intermedioque floris, caule superne dichotomo~~

Habitat in Java herb. Horren

glabra. Caulis herbaeus, superne dichotomus - folia petiolata, bipinnata, orata,
attenuata, integrifolia, extropum obsoletae crassata, subvenosa. Pedunculi disti-
tans, bipinnatos; apice pedicelli tres, rist unguiculatae, laterales triflori
~~intermediis que floris~~; folia duae orata parva. Bractea lanceolata, acutæ, à
lyra pium longior. Calyx foliata subulata. Corolla ~~racemosa~~ purpurea.
~~integerrimum~~: infusus emay natum. Anthene annata. Stamina longior
dine labii purpurea. Corolla purpurea. Seta nigra ex axilla

Dearthera

nitida Braetca

Habitat in montibus ad Ryan.

Pedunculi ^{partiales} inferiores utriusque gemino, polluians; alter ^{quam} brevior; pedicelle 5-8, unguiculatae; subumbellati: pedunculi superiores simplices. Braetca utriusque pubulata, unguiculans: ad basin pedunculum partialeum

Cvinda 118 *Glossia pedunculata* axillaris solitariae sub quadrifloris, brac.
foliorum oblongis ciliatis. Schub. fl. p. p. 21. t. 4 (Dianthus) aut. Stev. Bor.
^{et al.} n. 938.

Habitat in Taponia & Herb. Royeri C

Paulis erectus, hexagonus, pedalis, venis & cune villosum. Solia petiolata
1-2 pollicaria, oblonga, acuto, basi attenuata. Pedunculi alterni,
pedicelli, 2-3-4 florae, flore-pedicellata. Bractea duo subulata ad basim
cellorum, duo oblonga ad basim calycis concava extansores; 1-2-3 florae 4
ungueulares interiores & subulata. Corolla purpurea, bracteis longior

~~Giffaria~~
membrana cca herba cca erecta, foliis subfalcato-lanceolatis piceo
axillaris oblongi, bracteis bellum do membranacei quadrangulari
subovatis hinc angustioribus ciliatis aet. soc. h. nat Par. 1. pag 105.
Habitat in aere mendicali.

~~non vnde~~ polygala L. mea

3 Elytria 8 quamis floriferis ovatis, 8 lapi lanicollatis margine villoso, 8 lapi Lanicollatis
9 integrum mis, foliis lanceolatis integris 9 10 2
tubi flora Fusca aculeo, foliis lanceolatis integris glabri, 8 quamis floriferis 4 5 6 7 8 9 10 11 12
integerrimis ovatis caeruleisque lanceolatis integrum mis
Anonymus carolinensis; foliis radicibus, ovato-oblongis; scapo un
d' imbricato, 8 quamis cuneiformibus; spica confusa, bracteata,
terminali Walt. fl. Carol. p. 60.

Tubiflora carolinensis Gral. ed. Syst. Lin. 2. p. 1. p. 27.

Habitat on Carolina Rose.

Solia radicaria, subresilia, pluosa, bi-Tripollaria, acuta,
asenaria, vix oblique nervosa. Scapus pedalis, ^{stultus,} simplex, Testis pycno
mis alternis, puberugineis, unguiculatis, ovato-lanicollatis,
attenuatis, glabri, nitidis, post apice patulis, supraemis floriferis
4. interdum poli-^{ris} latioribus. Specie vix pollens imbricatus 8 quamis ovatis
tono, interdum acuminatis, marginis membranaceis et villoso. Flos unicus sub
sque, peduncula ^{spatula} angulo 8 quamis. Bractea linearis ad basin. Calyx profunde
5 que partitus, lacinia ^{spatula} sed opposita villosa, duo opposito oblongo
lacionis, altera apice bidentata, altera integra, trii anteriores
lineares angustiorer. Corolla limbis bilobata, lobis superiore bifido
inferiore trifido: lacinia crenata. Ita terminaliter

Lineanfolia G.

Lineanfolia ~~tutcheri~~ ~~color~~ *Lineanbus*, species at all *lambus* alternis
longe pedunculatis La Marsh T. Enc: pag: 41.
Habitat in Virginia Florida 3

3440 Daffusa

Genistifolia Fusca sp. ex axillaris fusca Verminalibusque, calycibus Tetraphyllis, foliis Lanceolatis.

Habitat in India orientali

Pauli herba eius, procumbens diffusus, sexangulatus, glaber, ceterum
inferne geniculatae intactis. Folia brevissime petiolata, desquipollita
via, acuta, integerrima, glabra, avenia vix nervosa. Spica sit longi-
tudine foliis angusto. Flores oppositi, solitarii. Bractea longitudine caly-
cis, Calyxis laciniis linearibus, lanceolatis, pallido dorso linea-
vindi. Corolla purpurea.

Planta distincta a varietate Engystipholia Fusca repentis, & multi minus
anno prostrato ab amiss. Rottlero communematum ^{sub non} nec antea nihil resu-
cula Bractea in hoc huius teste, longitudine calycis sunt nec brevioribus
ut in plantis herb. Herm. Etiam potius Wild. sed procumbens sic ut ex quo
procumbens potius diffusa di morninanda.

² 4 alpina ~~Pinguicula~~ neotanacetiæ o petalo breviore. S.V. p. 65:3. Flor. Lapp. 12. f. 12. fig. 3.
Fl. fin. 26. Spec. pl. 25:3. Flor. D. 453.

~~Pinguicula flore albo minore, calcaris brevissimo~~ Rijks Nat. 752.

~~Pinguicula calcare brevissimo~~ Hall fl. 293.

B. flore albo

~~Pinguicula flore albo~~ Bluf. Pan. 360

~~Scop. Flon. 22.~~

Habitat in alpibus Laponicus, Helvetiorum, Pyrenæis 4

Solia angustiora lere lanceolate. Scapi palmarer. foliæ plures
glabri. calcar corollæ brevis.

Synon. gesneri huc transfer

40 50
picta *Austicia racemis axillaris terminalibusque, floribus vestitissimis,*
foliis ^{ellipticis} ovoideis pictis. Symb. bot. 2. p. 14.

§. 129. *Tigridia - maram.* Rheed. Mat. 6. p. 60.

Solidum bracteatum Thunb. Amb. 4. p. 73. t. 30

Austicia picta; fructuosa, foliis lanceolato-ovatis pictis, corollis fuscis in
filiis luteis. Spec. pl. # 21. Synt.

Habitat in India orientali +

Ramorum articuli inferiores angulati. Solia brevissime petiolata, pro-
tentissima, acuta, glabra; disco macula conjugata irregulariter
lucida, integerrima. Pedicelli solitari vel 2. 3 oppositi. Racemi soli-
tanii, tripli pollicares: Pedicelli pollicares vel ^{dubius} 2. 3 oppositi, superne crassi-
res unguiculares, basi bracteis binis, rotundis, ovatis. Polys quinqueparti-
ti: laciniis lanceolato-porro. Corolla pollicaris! Tubus basi angustatus,
superne ampliatus quod nuptio latior: Labium superius lanceolatum, bilo-
bium; lobis brevibus obtusis: inferius trifidum; laciniis lanceolatis obtusis.
Antherae et stylus ^{longior} superiore longiores.

erisurgens Justicia foliis ovatis integerrimis, bracteis pubescentibus, ramis hexagonis L.V. p. 63.

25. Amoni Ad. 5. p. 391. * Spec. pl. p. 23. 14

Justicia herbacea erisurgens, ad alas alternas nodos et hummitates florida Brown
Jan. 110. f. 2. f. 1.

Habitat in Jamaica

Rami superne fortangulati glabri angulis glabris. Folia ovata integerrima acuta-
glabra pollinaria. Petiolis foliis ramorum floriferorum lanceolata mucronata
breviter petiolata. Petoli foliis duplo breviores. Pedunculi terminales sub-
dichotomi, tetragonoi articulati superne incrassati pedicellis patulis bracte-
filiaceis utrinque ad fragilans diffusam flores sessiles remoti b. bracteati bracteis
calycis longioribus altero ligulato majoro altero laci linearis brevore. Calyxplex
glauco linearis linearis ligulatus acuminatus tenuior, laevis membranaceis tenuioribus

Carthaginensis Linnae R. Let. 4.

affinis bracteolata

49

nubra Fuscia racemis terminalibus simplicibus solitariis, bracteis petal =
ceis, foliis lanceolatis. El. Amer. 2. p. 1.

Habitat ad St. Matham in America meridionali Von Rohr 5
Caulis ^{fruticosus} ramosus, Vener. Rami oppositi, patentib; bini, glabri, cortex canescens,
atticulati; articulis apice utrinque dilatatis. Solia petiolata, tripli-
cata, attenuata, acuta, integerrima, venosa, utrinque glaberrima, subtus
pallidiora. Racemi longitudinali foliorum, erecti. Bracteae oppositi,
versanti, inferiores subtriflori, superiores uniflori. Bractea minuta.
Calyx lacinia parva, lobulae rotundatae. Corolla pollicaris, gla-
bra. Filamenta corolla breviora. Anthers parallelae.

~~ster nitidum et flave~~

~~270~~ secunda Folii foliis ovato-lanceolatis acuminate, rameis terminalibus
compositis racemulis secundis, foliis ovato-lanceolatis acuminatis
Habita in Insula Trinitatis Dr von Rohr

Paulis herbaceis, cerasus, globosus, uti tota planta, Tetragona
~~fulci exaratus~~ ~~se~~ ~~le~~ ~~fulci exaratus~~ ~~angulatus~~ subexangulatus, lateribus
colea petiolata, bipinnata; attenuata integerrima.

Racemos terminalis, ~~racemis~~ ad apicem compotus est rameu
lis erecti, pinnatis oppotis, superne ^{verso} simplices summi
est simplices floribus oppotis, ~~laci~~ in racemulis alterni
4, breve pedicellata. Base Bracted duo, petiolo, longitu
dine pedicelli, ad base pedicellorum

Calyx spaditus: lacinia linearis lanceolatis, semivirgineis
ribus.

Corolla pinnata, purpurea, bifida: labio superiore linearis,
integri; inferiore sub trifido

Stamena capillaria, longitudine corollo. Antho binis
pedicellata, parallelo.
Style longitudine corollo

Determinate

171

21

174

50

Gnida Gusticia racemos axillaris bipinnatis, floribus secundis, foliis lanceolatis
ellipticis, filamentis glabris. Symb. bot. 2. p. 6.
Euphrasia minor Latione folio Madagasca ^{Dura}, conicum pubescentibus. Pluth
Alm. p. 142 t. 279 fig 4.

Gusticia pubescens; folio ovali-acuminatis, pedunculis lateralibus ramos ^{sc}
ramis floris, bracteis subulatis; calyx brevioribus L. Sormark March illust. p. 40
fida herb. Gustici

Habitat in Malabarum.

Paulus herba ev., aculeo Vitex agonis, glaber, lateribus utrinque sulco exarata ^{ad}
internodio bipollis arbus: Ramuli axillares, pediculis parum longioribus
floriferi. Solia longe petiolata, pollia avia et ulna, patentia, inde-
germina, basi per petiolum parum decurrentia, supra glabra, subtus
pollidiora, venosa. Pediculus foliis dimidiobrevior. Pedunculi et axillios
cicum ramulorum, oppositi, filiformes, patentissime, foliis triplo breviore
res: pedicelli oppositi, erecti. ⁴ Raceme lineariter vocen pedicellorum ^{conum}
que longitudinaliter duas ad basin divisura pedunculi. Flores pauciglilio
septe

histi

lum teneare, apice et inferne parum dilatatum, bilineare, basi calycem ad dimidiam
partem tegens. Calyx profunde quinque fidae, bractea parum brevior. Laciniis 2. Sta 200,
tredo, sc. pedicelli longitudinaliter tubi. Corolla tubum inferius rotundatus.
Affinis uidea differet; cæcile longitudinaliter sulcato; foliis lationibus longe petiolatis; pe-
dunculis foliis triplo brevioribus. Semel bifidis non ramiis et ramulis et floribus oppositis;
calyculis cæcioribus, basi bracteatis et filamentis hyalini.

~~U. L. P. O. N. O. T. A. R. I. S. U. L. C. T. R. I. M. I. N. D.~~

51
elongata ~~est~~. Fusca racemi axillaribus elongatis, floribus remotissimis subternis, foliis ovaliis.

Habitat in Indio orientali Röttler

Paulis herbaeaeus, strictus, tetragonus, angustiæ ~~et~~ ^{marginatus,} ~~angustibus~~ ^{atticulatis;}

Rami oppositi articulis & pitham et ultra, glabri # solo pedicello, remotissima,
brachiatæ pollicaria, attenuata, integerrima, glabra, nervosa. Pedunculi axil-
lares, oppositi, sœpe pedales, simplices, raro basi ramo cuncti, tetrago-
niæ. Flores pedicellati lini vel torni. Bractea gemina, minuta. Calyx
calycinis ^{pubescens} betaceus ^{pubescens} Corolla parva, Capsula angusta, semipollucare, ova-
+ pubescens lo armato villoso.

Pannaria

12

Habitat in Africa herb Ricch.

Rami teretis, rufi, superne glabri, superne flavo hispido Tomentosi. Sola herba petiolati subcordato-ovata, palmata, rigida, apice subrotunda, supra glabra, nitida, obscurè nevosa, subbasivirgata, venosa, & dilatata, Tomentosa, invenia, inde primaria. Pedunculus ~~latus~~ Verminalis, florae cinctus Tomentosus, uti calyx. Flores racemosi. Pedicelli breves, approximati. Petalum
laevis oblongo, obtuso

52

paniculata ~~Fuscia~~ panicula axillaris terminalis usque dicto horizonte; foliis basi crenatis,
dis, filamentis hispidis, capsulis compressis, foliis lanceolatis. Symb. bot. 2. p. 5.

Cara Canisom Rheed. Mol. g. p. 109. t. 56.

Fuscia paniculata; diffusa, panicula axillaris, foliis persistib; lanceo-

lati; Burm. Ind. g.

Habitat in India Orientali.

Caulis herbaceus; squame pedalis, strictus, tetragonus; anguli acuti; glaber,
lateralibus bistratiis. Solia fere 4; pectoralis, squame pollicaria, utrinque
attenuata, in petiolam brevissimum desinens, integrerrima, obtusa; stylus,
glabra. Pedunculi tetragoni, glabri, patentissimi; axillares solitarii, oppositi;
plenumque bifidi, subramosi. Verminalis paniculata ramosa; spiculae
culis partiglobis oppositis. Pedicelli alterni, remoti, capillares, solita-
rii, sive cum sedimento secundi in axillis bracteatum. Bractea oppo-
sitea, subramificatione primo linearis lanceolata; pedicellorum

Signos et huius ad omnibus miti- notis capsulis planis compressis undique eiusdem latitudini, nec basi angustata;

subtus late, brevissime, opposito, ad basin: nullas ~~basas~~ calycis. Calyx
quinq; partibus: lacinia pubulata, subobtusa, pubescens. Corolla
extus pubescens: labium superius bifidum; lineare, obtusum; in series
infidum: lacinia lanceolata. Filamenta longitudinaliter hispidissima,
longitudo fere corolla. Capsula six pollicaria, marginata, exoda utrinque
acuta, mucronata; glaucescens nitida, mediotinea purpurea; ante con-
giatio in aliis compressa undique

Gnaphalium aculeatum, foliis cuneatis integerrimi, spicis imbricatis, bracteis ovatis inter
gris mucronatis ~~foliis cuneatis integerrimi pedatais~~ Scopi marginae membranacei & levi
floriferae.
Habitat in africo. foliis cuneatis integerrimi

Saxos &c aculeato. Radix longa. Sola radicalia, 2-3 pollicaria, in petiolam attenue-
tur, extrorsum dilatata, obtusifolius, integriformis, ovata, remote pubescens, utrius-
que glabra, uti caput radicis. Scapi plures, digitales et ultra, glabri, a basi imbrica-
ti squamis ovatis, mucronatis, lanatis, margine membranaceo ando ciliatis. Caly-
cis foliola linearis-lanceolata, quadrangularis, dorso vnde glabra, excepto apice, mu-
cronata, aquatica, papuam versus ciliata.

53^{ff}
Scandens Luzula pedunculis axillaribus Trichotomis Divaricatis; foliis crassis
acuminatis subrepandis, ramis villosis. Lymb. bot 2. p. 7.

Habitat in Malabarica.

Rami scandentes, teretes, articulati. Folia petiolatae, bipinnatae, late ova-
ta, base acuta, venosa; venis utrinque ^{sub}vilosis. Pedicoli semiungui-
culares, villosi. Pubescens. Pedunculi oppositi, foliis brevioribus; primum
tripartiti, deinceps bimaculati, pedicelli bifidi. Flores pedicellati. Brac-
tea duae, opposita, minutiobimaculatae, ad basim pedicellarum calycis que-
que. Calyx laevis et staceo.

Venit ramis senisque foliorum minor vel magis villosis. Pedunculi parti-
les interdum racemosi, oppositi, duum planum,

^{Primum tripliciter} deinde
^{Biqualiter} bisecti hornis

polygonoides Ver. hispida, floribus solitariis subspatulatis, foliis alternis oblongis
caule simplici ab uno ad apicem floriferum laeviuscula
Habitat in pastuis Italicis et Gallis.

~~non vidi~~

cadem cum bellardii

54 ~~55~~
Gangetica Fuscia racemis bracteatis terminalibus que simplicibus, calycibus alter
ni his pedis, bracteis obsoletis +

Fuscia gangetica; foliis ovatis, racemis simplicibus longis, floribus alter
nis secundis, bracteis obsoletis Lin. Lv. Cimon. ac. 4. p. 299. Baron fl.

India q. t. 4. fig. 2.

Habitat in India orientali.

Caulis herbaeus, Tetragonus, ad genicula julurus. Solia brevis. Bime petiolata.
integerrima, glabra ~~aut~~, margine basaeos ciliata. Racemi ex aliis foliorum su
periorum solidani, elongati, erecti. Bractea pubulata, gemina. Calyx oblongo
porollo magistrali. Lin. L. C. ~~+~~

nemorosa & caule herbaeо ветвягono erecti usque, foliis ovato-lanceolato,
fructu ovatis Sw. prod v. p. 14.

Habitat in Jamacia, Hispaniola & Swanne inspt

Caulis pedalis sum pleris angulis vellopsi. Sola petiolata, 3 vel quadri
polluviaя nemorosa avenca oculo armata, lenei minutis aliis puliformis
bus oculo armato redendi, petroli pubescentes. Specie terminali
pollucans, umbonata bracteis ovatis acutis vellopsi

Lanceolato ovatis

non vidi

~~54~~ 55
nudans Fusussia racemis terminalibus apice nudantibus; floribus resupinatis, foliis ovato-lanceolatis dentatis Symb. bot. 2. p. 6

Fusussia nudans; herbacea, glabris lanceolatis denticulatis, pedunculis terminalibus cernuis, bracteis petiolatis Burm. Ind. p. 10. t. 5. f. 1

Habitat in Java Commerson

Paulo herbaceis; Veretivis tenuis, subtiliter hirsutis; glaber: Rami alterni. Solia brevissime petiolata, pollicaria vel ultra, attenuata, rara pilosa, patentissima, remote denticulata, venosa; venis alterni, simplicibus: humma lanceolata, integerrima. Racemi pubescentes: pedicelli breves. Bractea ad basin pedicellorum, et duo sub singulo flore. Calyx profundus qui que fidus: lacinia linear-lanceolatis, corolla triplo brevioribus. Corolla divaricata; operus (inferius) angustius, bidentatum; operus (superior) trifidum: lacinia lanceolata.
Stamina in labio inferiore.

labio

56 61

tintoria. Fuscia pedunculis axillantibus congestis unifloro; foliis lanceolatis - oblongis pubescentibus, caule pubescens, caule procumbente +
Fuscia Tinctoria; foliis lanceolatis, pubescens, pubescentibus; floribus congestis axillatis. Lourero fl. Corrense. p. 25.
Habitat in incultis Coquimboano.

Caulis herbaceus; ramosus, longus; teres; siliqua. Dolia acuminata.
Bractea conica, obtusata. Corolla rosae; labio reflexo; superius oratum integrinum; inferius oblongum levigatum. Lourero C.C.

~~floribus solitariis luteis lato-lanceolatis oblongis quibus resimilis
et foliis membranaceis lato-lanceolatis suboblongis, longioribus, lepto-~~

~~oymodes~~ 4

~~Gaffira orymodes~~ caule angulato, foliis ovatis petiolatis, pedunculis
astillibus multifloris brevissimis, bracteis parvis apice pinnatip.
La Mat. T. En. pag 39.
adhatoda jalapa folio Burch. n. color.
Habitat in ameria meridionali

mehi ignota

57
Palicinum F. Justicia floribus terminalibus solitariis sessilibus; foliis lanceolatis.

Habitat in Peru F

Paulis fructuosis, ramosis. Folia versus apices ramorum, confertis, breviter petiolata, bipinnata, obtusiuscula. Flores terminali ramorum.

Palicis foliola quinque, lanceolata. Corolla sesqui-pinnata, elegans nubens; labium superius erectum, emarginatum inferius. Vitellum lobis oblongis. Stamina labio superiore breviora. Anthene parallela. Stigma indivisum.

Descripta secundum figura 900. Justicii quo in museo Ant. L. Justicii abservatatur.

16. 17

colcinea

³¹ Luffuria Specie terminalibus, foliis bracteisque ellipticis
margine nudis, corollatum galea lanceolata apice reflexa, Styrnate
Fusussia colcinea floribus ampli spicatis; folio ovato-oblongo
acute ang. st. gl. Ray. 10. tab. 5.
Pasticia colcinea intricosa, corollis bilabiatis; labio superiore indi-
viso, foliis bracteisque spicarum ellipticis acuminatis, anthers
parallelis Smith. K. piet. pag. 8.

⁴⁸ Papenna Brief Dr von Rode

Primi glabri. Sola elliptica lanceolata ferruginea, glabra,
integerrima, venosa. Specie terminalis, palmato-bracteo-
ta in rami ramos attenuata. Caliginis foliola linearis lanceolata.
Styra bilamellatum

1964
Secundo Justicia racemis terminalibus compositi, racemulus secundus mul
ti florit, bracteis petiolaris, foliis orato oblongis attenuatis.

Justicia secunda; racemis terminalibus compositi: racemulus secundus,
foliis orato-lanceolatis aluminatis. Symb. bot. 2. p. 7.

Habitat in insula Trinitatis

Paulio herbarius, erectus, glaber nimis, e bulbis tenuis subexangulatus,
angulis levibus. Solia petiolata, bipinnata, integrifolia, glabra.

Racemus terminalis, palmatus, erectus, usque ad apicem floriferus
compositus. Pedunculi pediculares oppositi, pinnatis vel panini ultra,
in speciminiis majoribus brevissimi inferiorne brevissimi apice peduncul
culos plures subumbellatos inaequales gerentes, superne subverticillati;
doloris ope lenti pubescentes. Flores alterni, quinque novem trorse pedicellati.

Bractea duo petiolaris ad basim pedunculorum pediculatum et pedicel
lum: Calyxis laciniis quinque partitus: laciniis linearis-lanceolatis, semiungui
culares. Corolla pollicaris purpurea: labium superius lineare, int
erius; inferius bibrigidum. Filamenta capillaria, longiora corolle.
Anthera binata, pedicellata. Stylus longitudine corolle.

~~americanae~~ *Dianthera specis foliariae alternis* L. V. p 63. 1. Sp. pl. p. 24. Grana 4.
~~gratiola affinis floridana, foliis floribus et capsulis in spica brevi, pediculis e foliorum~~
~~alis in hirsutis~~ Plu. Kn. Amalth. 1/4. t. 423. f. 5.

Dianthera grana. Virg. 4

Habitat in Virginia, Florida 4 brevis

Caulis ~~se~~ angulatus sileatus glaber folia petiolata canaliculata
ut unque attenuata vel quadriplicaria integriforme
glabra simpliciter pubescens pedunculi axillares folio longiore
angulati. Spica terminalis densa oblonga semplicans
bracteas et lacrimas calyces canescentia canaliculata

ciliaria L. *Festuca* ^{spicata} ~~lanceolata~~ specie *terminalibus foliis hispidis*
Festuca ciliaria Gaertn. Hort. Vind. 2. p. 47. t. 104.

Festuca ciliaria; foliis lanceolatis, floribus oppositis sessilibus; bracteis
 calycibusque petiolaris hispidis flore longioribus Lin. Suppl. 84.

Habitat - O

Caulis erectus, pedalis, pilosus, scaber: ramuli axillares, bi-quinque
 pollicares. folia breviter petiolata, remota, integrifolia, obtusiuscula,
 subtilis ne rosa, nervis hispidis. Pelioli, uti bracteis et calycibus hispidi
 floribus oppositi, spha unicus in axilla foliorum. Bractea utrinque ad
basin singuli flori, calyx longior linearis, acuto, erecto patens. Ca-
 lycis laciniis linearibus, inferne marginie membranaceo, superne lateri-
 bus hispido. Corolla alba: labium superius brevius, ovatum emar-
 ginatum, viride-flavescens, triatum: inferioris trilobum; lobis rotun-
 datis equalibus, prope faciem ferrugineo bipunctata: Tubus brevis;
 subtilis ad laevem gibbosus, supra intus; intus forniciatus, stans ex cla-
 vescendibus, basi purpureo. Filamento basi barbata. Anthera superiore glo-
 bosa; inferior longior aristata. Capsula obovata, glabra. *

61
ganda nuda Justicia spicis terminalibus foliorum floribus verticillatis, folios lanceolatis elongatis.

Vada hodi Rheed. Mat. q. p. 79. t. 42

Adhatoda maderaspatensis hydrocypensis folio Petiv. ⁱⁿ Act. angl.

n. 244. p. 319.

Ranuncular galeatus Balicis glabro folio Bisnaganicus floribus aurantiis Blattw. Alm. 316. t. 55. f. 3.

Gendana nuda Knorph. Emb. 4. p. 70 t. 20 et p. 72. t. 29

Justicia gendarussa; glandulosa, folios lanceolatis, integerrimis, spicis terminalibus simplicibus Burm. Ind. # 10

Justicia gendarussa glandulosa, glabra, folios lanceolatis elongatis, spicis verticillatis Lin. Suppl. # 85

Habitat in India orientali. 5

Anthem
inferior aristata

Rami glabri, verticiles. folia brevisima petiolata, bipinnata et ultimis rami glabri, verticiles. folia brevisima petiolata, venis simplicibus. Specie palmata taguroma, glabra, obtriangularis, venis simplicibus. Specie palmata flore utrinque duovel tres quartos; folium tereare - lanceolatum per utrinque pubescens vel verticello utrigulare, superioribus membrinis brachio petiolatum longitudine florum. Bractea petiolata calyx lacinia petiacea. folia lobata inferius pedatis, amplius verte

Porollo labium inferius patens, amplum, conversum, apice tridentatum;
crena media rugosa, calix, marginata, subrevoluta, concava: laterales
promiscuas; convexus circulo, gibbo thicko.

~~Ludovicus Linneus. Exsiccatum. 1753. Ed. 1. Vol. 1. No. 1. 1753. Th~~

62 ~~by Quotia~~
debilitate spicis axillaris terminalibusque, bracteis ovatis imbrevis et secundis
ciliatis Symb. bot. 2 p. 15

Dianthera debilis; foliis oblongis, spicis axillaris imbratis; bracteis
late ovatis ciliatis Dorsh. Descr. p. 9.

Habitat in Arabia felice

Rami obsoletae quod ruranguli, quadrivalvis, retrosum pilosi, articulati;
articulati per que pollicis ari bus. Sola basi in petolum brevi simum congu-

dimidiata ~~spicis~~ ~~axillaris~~ rotata, obtusincola, arenaria, battus nevrignatior
rurum, parallatis villosis. Spica oblonga, tetragona, foliis brevioribus, axillari
alternis, sessiles, terminali pedunculata. Verticilos verticillata, qua
ternis, articulata, secundum nervos villosa, duo majores. Palyn
quinq[ue] partiti: lacinia linearis-lanceolata, ciliata, bracteis bre
vioribus. Corolla ex his pubescens, bracteis paniculata longior
Diffusa. rami dulcioris retrosum pilosis, foliis desigibus, spi
cis tetragonis axillaris, bract

Dianthera debilis; suffruticosa, foliis elliptico-lanceolatis, spicis
terminalibus, anthenis inferioribus anistatis. Symb. bot. 1 p. 5

~~pedinata~~ Deacte Justicia diffusa, spicis axillariis sessilibus terminalibus, secundis
dorsi imbricatis: bracteis semi lanceolatis s. v. p. 63. 19. Amon. Acad. 4. p. 299.*
Spec. pl. p. 22. q.

Habitat in Apia Diffusa

Caulis fructiferus feliformis retrogoni: angulus puluis reversus. foliis
petiolata oblonga obtusifolia glabra ope lenti puluis roris adspersa
angustiulana: spicis axillariis et terminalibus foliis confertis
pedunculatis inferioribus pedunculis foliis rami formibus longitudine
foliorum: spicis imbricatis adie florum dorsi bracteis imbricatis oblon-
gatis, foliis offerto: spicis imbricatis adie florum dorsi bracteis imbricatis oblon-
gatis, membranaceo: bracteis interioris altera lata subrotunda marginae uno que mem-
branaceo: bracteis interioris sub rotundata calycis vellosa
coctenium redit: bracteis interioris sub rotundata calycis vellosa
Guttiera parviflora Diffusa, foliis ovatis integris spicis secundis
bracteis lanceolatis: anterioribus alatis Retz obj. fof. 5 p. q

636
Rohrii Festuca spicis terminalibus compositis imbricatis pubescentibus, bracteis
ovatis, foliis ellipticis integerrimis. Symb. bot. 3. p. 6

Habitat in Rajenna

Paulus herbaceus, erectus, ramosus, Petrazonus, superne Venerabilime pubescens,
pubescens. Solia petiolata ^{ben octo} pollicaria, petioles ^{duo - tres} lata, attenuata,
autem acuta, nervosa, integerrima, glabra. Petiolis bipartitum
Spicula partialis breviter pedunculata, opposita. Pedunculi pubescentes.
Flores oppositi. Bractea binas, unguicularer, longitudine calectis. Ca-
lyxis laciniis linearis-lanceolata, pubescentes. Corolla labium
superius linearis-lanceolatum, in duplo; inferius trigonum.
anthera una super alteram.

castthaginensis *Gustavia* spicis axillaris terminalibusque, bracteis oblongis im
bractis ciliatis obtusis

Gustavia castthaginensis, foliis ovalibus, utrinque acuminatis, floribus
spicatis, bracteis oblongo-cuneiformibus Farg. am. ^{v.} 5. t. 5.

Gustavia mirabiloides, foliis ovalis acutis margine undatis, spicis gemi-
nis terminalibus, bracteis ovali-calyce longioribus ^{Lamark} Le Maist. illustr.

I. p. 39.

Habitat in Panboi v. Rohr

Racemos herbaceus. Rami oppositi, inferiore verticis glabi, superne obscure
setigere illosiusculi. Sola petiolata, lvs. Tripollaria, ovata, base
acuminata vel attenuata, acuta, venosa, nervosa, pubescens, pilis
vans ad sparsa, frequentioribus in costa et nervis. Spicis terminali-
bus ramorum, pleniusque solitario interdum gemino, bipinnatores. Brac-
tes ^{exterioris} et ^{interioris} unguiculatae, tace ad basin calycis cuneatae, obtusae, breviorae
stigma obtusum.

Spathe foliis ovato-oblongis et ellipticis ovatis. Bracteis exterioribus ma-
gebrotundis oblongis vel magis oblongis. Anthera inferior apud basi membranacea
D ^{anista} sec. Farg. an- bet.

~~clarata Fustera~~

~~Dianthus clarata~~ 80ff.

~~Habitat~~ hec Mus.

~~Ram. Tetragoni, glabri, ad articulos parum latentes. Solia petiolata, fere pollicacea
elliptica, utrinque acuta, glabra, subitus annectim nervosa, marginè subundulata, inde
genima. Pedunculi articulares, oppositi; folios fere longiores. Inclivitudo. Bracteas
pedicelli subulato minimo, ad flore linearis lanceolata, obtusa, flore parvi~~

retusa Astuccia spicis terminalibus, bracteis obvatis subtus is imbricatis glabris, foliis
ovatis adumbratis. Symb. bot. 2. p. 8

Habitat in insula St. Eustacii.

Paulis herbarius, Veres, glaber, ut: Voda plantae. Solia petiolata, bipinnicaria
et ultra, ne propter inique elevatio, integerrima, obtusiuscula, subtruncata,
lodicula. Pedunculus Terminalis, Simplex, Trivel quadruplicaria;
apice floriferus. Spica pollicaris. Flores solitarii, oppositi. Pructus
terno ad singulum florem: extenor valvis multo latior, apice angustata,
lato, oblonga, basi sensu angustata; duos laterales, oblongi, basin ver-
tus attenuato, exterior breviores, pilis minuti subtilitate. Calyx quin-
que partitus: Cacinis lanceolatus. Corolla magna, purpurascens, ultra
medium fissa; labium superius lanceolatum, bidentatum, in inferior bre-
vius; inferior triplo latius, Trifidum; Cacinis Oblongi. Stamina
longitudine labii superioris.

~~oblonga
basin sensu
angustata~~

rad. ~~66-42-68~~
radicans foliis lanceolatis oblongis glabris, pectinis axillaris terminalibus
oblongis, bracteis oblongis ciliatis anastomatis, foliis lanceolatis oblongis glabris +
Dianthus ciliata: foliis lanceolatis, floribus axillaris pectatis confertis,
bracteis calycibusque ciliatis fl. penne 1. p. 12. t. 14 f. 7 a.

Habitat in Penuria nemoribus.

Caulis herbaceus, bipinnatus, erectus, simpliciter hexagonus, radicans. Sola
petiolata integriflora. Flores in forma capitate collecti compacti, per opposi-
tum sessiles. Bractea acuminata in forma longa ciliata, ad apicem pubescens
ad basim calyculis. Calyx laevis ^{quinq} pubescens. Corolla purpurea
violacea, fons extus pubescens. Labium superius emarginatum
in formam trifidum. Semina scabra. Fl. p. 12. t. 14 f. 7 a. 187. Per. l. c.

Secundum figuram

+ seorsim

strobile

6869

nemorosa *Justicia* spicis terminalibus; bracteis imbricatis ovatis adutis - ^{villosas}
foliis ovato-lanceolatis

Dianthera nemorosa foliis oblongo-ovatis cum acuminis, spicis
florum geminatis Brown Ham: 110.

Gustavia nemorosa, caule herbaceo strigoso eretiusculo, foliis
ovato-lanceolatis, spicis oblongis Swartz prodr. # 14. St. Ind. Occi-
dent. I. p. 37.

It habitat in umbrosis paludosis & silvis Americas calidioris i-
Paulis herbariis pedalioid ultra, ang. simplex, angulis villosi: 80-
lio ~~per~~ ^{trunc} ~~quadr~~ pollicaria, nervosa, avenia, ~~de~~ ad cuncta fineolis
albidis foliis immersis ad spesso, acuminata, obtuse serrata.

Petioli longi pubescentes. Bractea nervosa, viole-purpurea et
lato

interno lanceo Corolla purpurea-purpurea. Posa labium superius emarginatum:
inferius trigidum. Stigma obtusum.
valde affinis catharinensis.

~~bedd~~ 8 annos

~~fastigata~~ ~~Turritis~~

~~Habitat~~ herba ~~Mantii~~

Paulus herbarius obtuse tetragonus, glaber. Sola petiolata, oblonga, ovata
attenuata, lepida et ultra, glabra, intermedium subrenata, supra
nitida. Racemi axillares et terminales subcymopodi. Pedunculi villo
so-viscovi. Pedicelli uniflori. Bractea 4-5. villosa, uncinularia; no
qualsis; villosa, calyx triplo brevioris. Palyn 5que partitus: lacinia
pubulata, pilosa glandulosa. Corolla pollicans. tubus lineans, calyx
longior, glaber. limbus regulans extus oculo armato villosus. Anthere
parallela villosa
an Eranthem. herb. ~~Herrmanni~~

~~206~~ *polystachya* *Gustavia* spicis axillaribus oppositis imbricatis secundis; bracteis ovatis hispido- foliis lanceolatis oratis integrinatis Symbola bot. 2. p. 7. t. 26.

Gustavia membranacea; herbaea, erecta; foliis subfoliatis lanceolatis. Specie axillaribus oblongis. bracteis pallidis membranaceis, subserratis, hispidis angustioribus, ciliatis. Act. Soc. h. n. Paris. 1. p. 105.

Gustavia polystachya; foliis oblongo-lanceolatis, spicis alternis axillaribus, bracteis ovalibus villosis non rosopallidis ^{La marck} ~~Cathart~~ illust. 1.

p. 40.

Habitat in Cognac von Rohr. 4

Paulis erectus, striatus, obscure Vetraceous, angulis retrorsum pilosis. foliis petiolata patens ^{bracteis} ^{lanceolatis} ^{oppositis} ^{angustioribus} ^{ciliatis} ^{oblongis} ^{lanceolatis} ^{oppositis} ^{angustioribus} nitida, subtilis pilosa.

Specie solitaria, ^{axillaris} ^{ciliata}, folio dimidiis breviores; imbricatis; terminali foliis via. Bracteae anteriores oratae, membranaceae, semiunguisulares, nervo 10, ciliatae, dorsalem semioratae, ciliatae, cuspidatae, unica ad basim singuli florae. Flor solitarius, perennis, in aliis bracteana. Calyx quinque parted: laciniis petiolatis. filamenta apice bifidae laciniis altera erecta, altera deflexa. Anthera unita in stylola latissima filamento.

peruviana Guadalupe

~~Labium superius bifidum, angustum, inferioris trifidum. Anthera altera
supra alteram, inferior anisata: anta subbifida~~

~~villoso~~ lobos oscillantes ^{sub} lacinia terminalibus vernis felis cani edato
ovatis

et S. peruv. Corion. minima la Marthe foliosa Jacq.

69 70 violacea Faschia spicis terminalibus; bracteis lanceolatis imbricatis; foliis lanceolatis. Symb. bot. 2. p. 15.

Habitat in Arabia felice s.
Rami pulvinulenti villosi, obsolete quadranguli, articulati; articulis
sesqui pollicinibus. Dolia petiolata, longitudo per foliorum, in inferiora
oblonga, superiora lanceolata, basi dilatata, integrerrima, glabra, late viridia,
avenia. Petiolis brevis. Spica brevissime pedunculata, semi-gollucaria;
cylindrica. Bractea alumina & subpungente terminalis; in fine longitudo
per spicam, alia 2 ad basim calycis. Calyx coriaceus ^{quinq}, pubulatus,
glaber, bracteis brevioris. Corolla extus pubescens.

Dianthera violacea; caule fruticoso, spicis terminalibus imbricatis; bra-
cteis lanceolatis, floribus bicarpellatis. Symb. bot. 1. p. 6.

~~picra~~ Justicia ~~picra~~ foliis lanceolato-ovatis petiolis, corollis facie inflatis. S. V.
p. 62. 6. Spec. pl. 21. 5. *

Thunb - Manam Rheed. mal. 6. t. 60

Tolium bracteatum Rumph. Amb. 4. p. 73. Tab. 30

Habitat in Asia

Ramo ^{ramo} angulato articuli angulati, superiores ut in congeneribus. Folia petiolis
brevisimo, patentissima, elliptica acuta, glabra integriforma diffuso maculata
longitudinali irregularis lucida, bipinnata. Flores in Racemis laterali bus
terminalibusque, solitariis, 3-4 pollicaribus. Pedicelli solitarii vel 2-3 oppositi
^{superne incaustati} (superne incaustati) bracteis brevibus, minutis, ovatis. Calyx sparsus lacinias lanceolatas
paras. Corolla magna pollicaris tubo basi angustato superne ampliata quadriplo
latis. Labium superius lanceolatum in earlobium lobis brevibus ^{oblongis} rotundatis; inferius
in fidum lacinias lanceolatis obtusis. Anthera et Stylus labio superiore longiores

~~#~~ 70
flava Austuria spicis terminalibus, floribus geminis, bracteis lanceolatis ob-
tusis, foliis lanceolato-oblongo

Dianthera amencana & flava, spicis terminalibus umbellatis bracteis
lanceolatis; foliis & calyculis; floribus flavis labio inferiore latiore,
anthera inferiore condita Dorsk. desert. p. 9.

H Habitat in Arabia

Paulis ~~fructuosa~~ ^{opaca} fructuosa, heretius culus, crassitie penno aveneria.
Rami obsoleti angulati, subpubescentes; lateribus linea extant,
angulati; articulis pollicaribus. Folia petiolata, articulis duplo lon-
giora, integerrima, rare pilosa, venosa. Petiolis brevis. Spica pedun-
culata, bipinnatis, subpilosa. Flores contigui. Bracteae tri subpingue-
lo flore, obtusa, subtruncata villosa, unica majora longitudine floris.

Polyx quinquefida, villosa: lacinia subulata. Corolla extus villosa,
longitudine bractearum: labium inferius trifidum; lacinia lateralis
lanceolata, intermediana oblonga, obtusa. Anthera inferior aristata

Dianthera flava; suffruticosa, foliis elliptico-lanceolatis, spicis terminali-
bus, antheris interioribus aristatis. Symb. bot. I. p. 5

~~72~~⁷¹
alata Fusca racemis axillaribus, foliis linearē lanceolatis, caule subd-
ato.

Habitat in India Orientali König.

Paulis herbaceo, elongato, membranis quatuor angustissimis
decurrentibus quadrangulari-alatus, tenuis. Dolia subsessilia
bitexta triplacaria, remota binia, suprema angustissima, glabra
pubes pallidiora, integerrima. Flores remoti; pedicelli breves,
ope lenti villosuli. Bracteas triangularē ad basin.

Dicotyledon

~~euflachiana~~ *Dianthera* ~~lolicis~~ lanceolato-oblongis, pedunculis multifloris, bracteis linearibus apice latiusculis acuminatis s. v. p. 64. 5 Gaeq. Am

Habita in Apula & Eufachii t

Folium erectum Tripedale lato superne herbace caeruleo-laceo
et decurrentibus aliis angulatis glabris. Tolo petiolata obata
sequi pollicis oria inferiora obtusa cum mucrone superiore acutiora
nervosa pubescens puberula. Specie ~~an~~ ^{vera} flavescentia terrore elonga-
to inferiore basi rameo. Flores ^{vera} ~~vera~~ ^{vera} flosiles inferne gemini
remoti altero pedicellato superiore foliatis distantem ^{rum} bractea numeris
ap cuneiformes altera parum longior. Calix calyris pubulata

72 ~~#~~
Americana Fuscia spicis axillariis; floribus confertis; pedunculis elongatis ac
ternis; foliis lanceolatis

Gratiola affinis floridana; foliis floribus et capsulis in spica brevi;
pedunculis a foliorum alio enixa. Pleben. abneth. 114. t. 423. f. 5.

Dianthera monos. v. ro. 4.

Dianthera americana; spicis solitariae alterno Lin. Syst. Vag. 64

~~#~~ Fuscia linearifolia; foliis linearibus; spicis axillariis alterno longe
pedunculatis ^{Lamaf.} ~~laevi~~. illustr. sp. 41.

Habitat in Virginia, Florida, in littoribus hieme inundatis fluminum
Ohoi st. Langenti, mississippi Michaux
Paulus ex angustis, Buleatus, glaber. folia breviter petiolata, et
utrinque attenuata, trinervia quadruplicaria, integerrima, glabra,
simpliciter subvenosa. Pedunculi foliis ^{elongatis} angulati. Spica
oblonga, densa, semi-pedicellata. Bractea et Calyx lacinia linea
ri lanceolata.

~~#~~ Fuscia ensiformis Michaux
Dianthera ensiformis; spicis subcapitatis, pedunculo longo solita
rio, floribus rubris, foliis ensiformibus. Walt. fl. car. 63.

~~moreiana~~ G. flonbus ~~stellariis~~ solitariae, foliis lanceolatis integerrimis Rumpf

herb. 61723

Thunb. herb. - amb. 6-723

Habitat in India orientali

Rumpf

nom. vnde

procumbens 7³ Fuscia ^{Lateralibus} Spicis axillaris terminalibusque, calycibus & oophyllo-
lincaribus pilosis, glabriuscule
Setae pilosa, labia in seriore subhispidis, foliis oblongis.
Euphrasia Binaria parvula foliis rubie modo spicatis. Plana. aln.
83. t. 392 f. 4.

Hulu-ray-kola. Herm. regl. 2. 4. 15. 42. 32.

Fuscia foliis sublanceolatis integerrimis, spicis terminalibus
lateralibusque alternis, bracteis setaceis Lin. fl. regl. 7.

Fuscia procumbens; foliis lanceolatis integerrimis, spicis terminali-
bus lateralibusque alternis, bracteis setaceis, caule procumber-
te Lin. 8 yst. Veg. 63.

angustifolia ^{pro} foliis lanceolatis.

rotundifolia foliis subrotundo-ovatis, caule subhispido.

Thymifolia foliis ovatis obtusiusculis, caule ramosissimo

Habitat in India orientali et China, / ad vias Tranquebariorum
Paulis ^{petrae aquilonis} ~~procumbens~~, diffusus; geniculis inferne & cernue que radice
infractis, infimis radicibus. Rami dichotomi; articulis bipinnatis
lincaribus, petiolaris, one lenticulis villosiusculis. Solia pediolata, remota
verticillata.

Demi pollicaria, alata, integrerima, scabriuscula a lineis albido elevatis orulo armata
so videndis. Spicæ terminalis vel ad dichotomia, ramulorum basi folia duæ, pollua-
rei, erectæ foliæ conferti, oppositi, parvi. Bractæ binæ ad basin singuli calycis.
Calyx membrana angustissima magnitate utique marginata. Corolla ante explica-
tum pilosiuscula, dein glabrior; Labium inferius purpureum, trilobum.

Anthera inferior membranæ albida anisata.

Quod fusto procumbens Lin. a mea eadem ac mea est, ex collatio Specimi-
mum quia quisque hinc habet hunc spicæ ipsius.
ut quisque hinc habet hunc spicæ ipsius.

Varietas qd' gaudet foliis subrotundis vel ovato rotundatis & caulibus elongatis.
foliis vix magnitudine Thymi & capilli majoribus, plenumque ovatis vel oblongo-subrotun-
dis, spicæ gracilibus, bractæ calycis binæ angustissimis
foliis vix magnitudine Thymi & capilli majoribus, plenumque ovatis vel oblongo-subrotun-
dis, spicæ gracilibus, bractæ calycis binæ angustissimis

~~at allatam mif. satis multorum ruforum~~

Diffusa 14 Fusicia Specie axillari bus terminalibus que, calycibus & traphyl-
lio lanceolatis glabis, labio inferiore barbato, foliis lanceolatis.

Fusicia diffusa; Specie axillaribus terminalibus que, calycibus &
traphyllis glabis, bracteis calycis brevioribus; foliis ellipticis Wid.
Spec. pl. I. p. 87.

Habitat in India orientali. Röthler

Diversa a varia late angustifolia fust. procumbens, mecum anno
prosternito communicata a Röthlero, nec antea mihi visa. Differ-
re videtur ab antecedente: foliis duplo vel triplo longioribus: bracteis
vix villosis: margine membranaceis foliorum calycis latiore: Labio
inferiore subtiliter versus marginem pilis frequentibus barbato. anthera
superiore oculo armato pilis raro adspersa. Versor ramen quod a mi-
nus exterritate difficile dignoscitur, prostatina cum prior invenie-
tur calyculis magis rara glabris, foliolisque latioribus vel an-
quistioribus. In quodquod vidi bracteas longitudo calycis inveni-
variat foliis latioribus & angustioribus. Postenam modus consciendi,
superficies foliorum, color florum & antherae ~~per~~ precedentis procedenti.

Corolla

75 # punctata *Gaudia* Specie terminalibus, floribus remoto verticillatis, bracteis lanceolatis acuminatis, folio lanceolato-ovati. Symb. bot. 2. p. 15

Dianthera americana B) alba, flore I anthero albo; labio inferiori in fovea violaceo punctata, verticillis remotis, spicatis; bracteis lanceolatis calyci duplo brevioribus. Dorb. Descr. E. 9.

Habitat in Arabia

Dianthera Raulis herbarius, crederem, Berolinus Culicis, locis, articulatus; articuli Tripoli punctata; caule latis arcu: ramulis axillares, filiformes, longitudine articulorum. Dolia herbarico, specie terminalibus, caulina petiolata, lanceolato-ovata articulatis Diantherae breviorum foliis lanceolato-acuminata, vix crenata, glabra, & implexa rasa; ramulorum ovatis acuminatis, anthers metitio. Symb. bot. 1. p. 4

peculiaris, unguiculata. Specie in apice rauulorum, pedunculata, pollicares, glabra. Pedunculus filiformis. Flores verticillati, uncinque ^{duo-tri}, brevisime pedicellati, parvi, inferiores remotiores. Rhizome lanceolata, acuminata & lineam longo.

Calyx bracteis longior, quinque fiduc: calycinis puberulatis. Corolla calyci duplo longior, et non glabra; fovea violacea punctata - anthers metitio.

76 ~~#~~ *Gustavia* spiciformis terminalibus ^{axillaris} ^{tacca} filiformibus, spiculis verticillatis

#

Amborenum minus angustifolium, flore dilute coeruleo Sloan.

Gen Catal. 49. Hist. 1. p. 160. t. 103. f. 2.

Dianthera folio lanceolato-ovatis, racemo spathoso aspergente,
spiculis verticillatis Brown. Jam. 110.

Dianthera comata; spicis filiformibus verticillatis, inferioribus
umbellatis Lin. Syst. ~~Sp~~ 64

#

Habitat in Jamaica 4

Paulis terer, ramosus, globos. folia bipinnata, breviter petiolata,
lanceolata, attenuata, glabra, subtus pallidiora, simpliciter ve-
nosa. Pedunculi axillares pariales utriusque ^{due} ~~tri~~ bipinnata
ver, inaequales, patent. Flores alterni, frequentes, raro. Brac-
tentias ad basim calycis et unica oppositi flora, subulata, longi-
tudine calycis.

Gustavia comata folio lanceolatis acutis. subserratis, spicis filiformi-
bus verticillatis umbellatisque Swartz. obs. p. 14

~~aciculans fructi ova diffusa, spinos at foliis oblongis mucronatis, spinis
at lacinibus petiolis, pedunculis at lacinibus solitariis
Austriaca aciculans fructi ova diffusa, spinosa, spinis at petiolis
foliis pedunculatis at lacinibus solitariis su-prodo. pag. 13~~

Habitat in Tamaqua Swartz

Rami Verteret cotyledone cinereo glabi non dosi, ramuli oblique
tetragoni subpubescentes. Spinae venae patentia foliis parum
breviser folia plenariaque solitaria, breviter petiolata oblonga
unguiculana obtusa mucronata, basi acuta integerrima pubes-
cens pallidiora supra, rursum linearibus alba v. orato armato redendo.

17-~~26~~-44

reptans Justicia spicis terminalibus solitariis, floribus secundis, foliis ovaliis,
tacete repente

Quotida reptans; coula herbacea repente, foliis obtusis, spica Var =
minali indivisa Swartha podo. p. 13. 88¹ Ind. Sci. D. p. 33.

Habitat in fastosi umbrosis madidi ad ripas fluviorum in Hispania
niola & inque Domingo

Paulis filiformis, 23 pollicans, angulatus. Tolia petiolata, ~~sep vix~~
unguis plana, interdum acuta, ~~sep~~ obtusa, nervosa, glabra. Spica
~~tri-quadrata~~ ~~&~~ flora, multiflora rufa, flores minuti. Bractea tricuspidata, val-
ce brevior. Petioles latinius ~~quin~~, linearis, acuto ~~#~~ lobium superius emar-
ginatum; inferioris tripartitum. Stigma obtusum

Corolla:

Dicanta. bts.

polyachya ~~Gustria folii lanceolati integriformis, specie at illaribus oppositis
imbrevis, foliis ex parte herpeti, foliis lanceolato-oratis integriformis
Habitat in Cognac In von Roht Herd. di Lunc
Caulis herbaceus, erectus, strictus, oblique tetragonus; angulis retrospun
pelosis.~~

Folia petiolata, lanceolata & in pollicaria, lanceolata, attenuata,
acuta, supra glabra, notida, subtilis pelosa, patentissima

Petiolas brevissimas.

Secundo

Specie axillare, foliariae opposita, erecta, folio dimidio breviora, imbi
lata, terminali solitaria

~~apicis anterius~~ ~~ovato, membranaceo, nervoso, peloso,~~ ~~foliata~~
~~apicis posterius~~ ~~triangulo-~~ ~~membranaceo, nervoso, peloso,~~ ~~foliata~~
~~Dorsum~~ ~~unica~~ ~~foliata~~
~~oblique~~ ~~lanceolata, cuspitata, ad base paguli floris.~~

Flos solitarius, sessilis, in aliis bracteorum.

Calyx 5-partitus, simplex; Corolla setacea:

Anthera gemina

Filamenta apice bifida, latitia alterna ereta Obtusa deflexa: Antera
unica in pagula latitia

Gustria polyachya foliis oblongo lanceolatis specie
alternis axillaretis, bracteis ovalibus vellosis nervosis
pellucido la mark. T. En pag. 46

4 membranaceae
a d. for h. nat
Pani fed Lest
Rich

1829-24

~~humilis~~ 77. 70

~~humilis~~ 72. Justicia spicis solitariis vero in aliibus que panicifloris, floribus disti-
tibus, ~~pedunculis~~ ^{bracteis} foliis longioribus; foliis lanceolatis oblongo-
Dianthus ovata; spicis solitariis alternis feliformibus; foliis duplo
longioribus; foliis puberulis; floribus rubro-purpureis Walt. fl. Carol.

63

Habitat in Carolina. Bosc.

Paulio herbaceus angustatus, x - Volo plantae glaber, ramum ramosus.
Solida petiolata, remota, ~~sesquipedalis~~ foliaria, inferiora minora interdum
vix semipeltaria, integriformia, acutangula, interdum subrotata,
plenumque lanceolato-oblonga, ^{angustiora tenuiora. folio longiori} foliis non
seruis capiunt floriferous. Flores cinq. 5 apie, oppositi. Calyx
profunde pentagoni; lacinias linearibus acutis.

Branches
lineares, longi
stems in calyx
cav. protex.
non nullo
minutis
et basi

Justicia humilis; glabra: caulis herbaceus, & virpe repente asper-
gentibus: foliis ovali-lanceolatis: spicarum bracteis minutissi-
mis; floribus subbimaculatis sessilibus: capsula longe stipitata,
compreensa. Michaux. fl. br. Amer. t. p. 8.

~~ovata~~

~~Scandentia~~ ~~ovata~~ foliis subovatis, fructu solitario alterno pediforme
foliis duplo longioribus Walth. fl. car. p 63.

~~Habitat in Carolina~~

Scand. Walth.

non vid.

Eustachiano Justicia spicis axillaris terminalibusque, floribus inferne geminis superne solitaria, bracteis cuneatis. Symb. bot. 2. p. 15

Justicia eustachiana. Dianthera; foliis lanceolato-oblongis, pedunculis multifloris; bracteis linearibus apice latiisculis aluminatis. For gr.
Gomes p. 4. t. 4.

Habitat in insula St. Eustachii 5

Truncus erectus, Tripedalis. Rami ex lineis decurrentibus angulatis, glabri. Dolio petiolata, usque polliciana, ovata, inferioria obtusa cum mucrone, superiora acutiora, nervosa, pubescentia subvenosa. Spica axillaris, alterna, elongata, in inferiori base ramos. Flores 5 cavi ad: El in inferiori gemini, alter pedicellatus, alter perspicillatus, distantes. Bractae duae cuneiformes, mucronatae, altera parum longior. Calyx & lacinia subcato.

Tranquebariensis ~~Gutierrezia~~ suffruticosa, caule Vereti, foliis orbiculatis, fructu terminalibus,
floribus solitariis, bracteis obtusatis s. v. p. 62. 4. Suppl. pag. 35.

Habitat in Tranquebar t. oppositis diffusi

Suffruticosa ramis teretibus, decaricatis, in eam ^{tempore florit aste} ~~tempore florit aste~~ ^{tem-}
~~meano~~ incanus. Dolia orbiculata petiolata, bafi a cuto, integerrima
~~uniquaternaria~~, ~~quaternaria~~, venis incanis, unquaternaria; petiolis longis
dura fere foliis. Flores (axillares, solitarii, sessiles) in Spica terminali,
bracteis oppositis, obtusatis, adpresis, in quinque axillis flores solitarii
soliti. in Suppl. sed flores potius ~~est~~ in parentesi duci.

Spica axillaris foliosa primus. quae axilli
ut in ^{tempore florit aste} ~~tempore florit aste~~ Ecliptice

~~24~~ 80

nasuta Guttaria pedunculis axillaris dichotomis, foliis ellipticis integrimis

Symb. bot. 2. p. 15

Pul. coll. Rhed. Mal. q. p. 135. t. 69.

Guttaria nasuta; foliis lanceolato-ovatis integrimis, pedunculis
dichotomis Lin. Spec. pl. f. 23. Quat. mag. vol. 10 t. 325

Habitat in India Orientali

Rami obsoleti angulati, rix pubescentes. Sola petiolata, bipinnata
nigra, obtusa, integrimma, glabra. Pedunculi solitarii, alterni, foliis

pubescentes, opacantes, bis vel ter divisi, par foliorum ellipticorum sub prima divi-
sura: pedicelli biflori; flores sessiles. Bractea lineare, minuta.
Calyx minutus: laciniis rotatis. Corolla tubus linearis, calyx quasi
triplo longior: labium superius lineare, brevius, angustum, reflexum;
inferius opacum, trifidum.

~~Betonica Guettiera graminifolia, foliis lanceolato-ovatis, bracteis ovatis acuminate venofo
reticulatis coloratus. L. V p. 62. 4. Flor. Ley 18*. Sp. pl. 21. 6. Tabr. Helm. 2. p. 217.~~

~~Betonica Gratiensis Boott. Jav. 146~~

~~Bemurini Rhod. Mal. 2. p. 126. t. 21.~~

~~Habitat in India~~

~~rami obscuri angusti, articulati, paperne lat. planum latitores. Folia
petiolata ova' lanceolata - ovata, margini inaequaliter
acuminata, pollicaria. Petioles breves. Specia terminalis bi. f. triplite
cane. bracteate & bracteis ovatis, acuminate, pallidis, venis vindibus
pictis reticulatis. Flores solitarii, oppositi; lacinia calycinis lanceola
ta villosa~~

Antonio