

Launis

Jara Rohri est Launi frucei & quidem foba Pecunim. Mich.
Cnicu Aublotii Launi fruceo. Mich.
Juani. est Launis foliis lancolatis subtus reticulatis cortice flavo spon.
Launis globosa L. Mich. Plum. 4^o. 60

Launis foliis magnis glaucis a Nyanois est
Arbor glandifera foliis scolorandii herb. Vaillant
Vernx a Perdus à grande feuille Tomaco comalea Gadonem herb. Sur

melisifolia L. foliis cordato laccolatis venosi membranacei subtus
pubescentibus, gemmis simplicibus trifloris Walt. fl.

Car. pag. 10134.

Habitat in Carolina

Cyathiferus Launus foliis lanceolato-ellipticis supra nitidis, floribus paniculae ramis
dichotomis, calycibus persistentibus cyathiformibus lobatis.

Habitat in Montserrat de Reyan.

Rami glabri, cortice fusco, apice cinerei.

Folia petiolata, albida, 4 vel quinque pollicaria, attenuata vel acumina-
ta, basi subacuta, membranacea, glaberrima; venosa, supra nitida,
subtus pallidiora, costa, ubi ⁱⁿ venis majoribus ^{per}persea: junctura subter in
costa leviter villosa.

Petioles unguiculares, triquetri, supra plani.

Panicula infra foliacea et ex axillis superioribus, erecto patente, foliis
brevior. Pedunculi partiales albescens, brevis: dichotomi plerumque in-
terdum triflori, fructiferi fusci. ~~flori tomentoso lanuginosi~~ ⁱⁿ ~~erecti~~ ^{erecti}.

Calycer unicani: fructiferi glabri crassi lobis, cyathiformes, diametro un-
guiculari, margine integerrimo.

1500. ~~no 205~~ Montserrat

an exactata Swartzii,

Bicolor Laurus foliis elliptico lanceolatis subtus glaucis purpureo fereantibus, paniculis
erectis fructibus oblongis, calycibus deciduis.

Habitat in Montserrat in Hyar?

Rami glabri, fusci, apice teniter canescentes, tuberculati & capsulis foliorum

Folia ad apicem ramorum, approximata, petiolata, quadripolliciana,
obtusiuscula, basi acuta, vix undulata, membranacea, glabra, nervosa,
obscure venosa, subtus glaucis-purpureo fereantibus, venis majoribus fasciis.

Petiolum

racemi laterales et axillares superioribus, foliis brevioribus, erecto-patentibus.

Longitudine foliorum, pedunculi partiales alterni, triflori vel
dichotomi: ramulis trifloris.

Calyx deciduus, glaber; laciniis obtusis.

Fructus pollicaris, glaber, oblongus.

(An Laurus indica her.?)

Coryifolia Laurus foliis lato lanceolatis sesquipedalibus glabris, racemis terminalibus
buis decompositis, calycibus persisistentibus, fructibus oblongis.

Habitat in montserrat de Ryan.

Arbor. Rami teretes, glabri, cortice cinereo, infera crassitie pennis anfractis,
inferne cicatricati.

Folia in ramis florentibus versus apicem ramorum, ternato-verticillata
in fructiferis alterna, interdum opposita, distantia; omnia brevis
sime petiolata, sesquipedalia vel minora, elliptico-lanceolata
integerrima, inferne versus sensim angustata, basi plerumque
obtusissima, apice attenuato, acuto, membranacea, utrinque
glabra, supra laevia, glauco viridia, subavenia, secundum nervos
obscura lineata; subtus pallidiora viridi flavescens, venosa.

Racemi floriferi in apice ramorum subumbellati, decompositi
6-8 pollicares, pat diffusi. Pedunculi angulati: partem
basim sparsi filiformes; supremi simpliciores uti pedunculus
communis et pedicelli glaucescentes. Fructiferi axillares
vel laterales: pedicelli unguiculares versus calycem
sensim incrassati, pedunculo communi duplo crassiores.
Flores parvi, ordioii?

Calyx 6 fidus, canescens: fructiferus cyathiformis, magnitudine nuclei Coryli, in
regemini rugosi, validi.

Fructus oblongus, pollicaris, lavis, obtusis

elongata P. 17 caulis foliis ovato-lanceolatis, attenuatis membranaceis, ^{simplex} ~~ramiculis~~ ^{ramiculis} axillaribus; caly-

ribus persistentibus; 6fidis

Habitat in montserrat & Pyrenis, in G. polo St. Crucis West

Rami vix glabri, cortice fusco.

Folia petiolata, altera nuda, quinque pollicaria, extrorsum attenuata, ^{obtusius venata,} tuberculata, membranacea, nitida, subtus pallidiora, 3ripalinervis, nervis exstantibus alternis, superiora vix persistentia, utrinque gla-

Petiole

Pedunculi axillares, ramosi, perparvascentes, longitudine foliorum, parthalibus, alteris, 3stantibus, quadriferi in rascati.

Calycis laciniis oblongis, ovatis, glabris, persistentibus.

Fructus oblongus

Leucoxylon laevius foliis oblongo-lanceolatis planis perennantibus, vixenni foliis
brevibus, calycibus unisulcatis verrucosis. Sw. prodr. pag. 65.

In Montserrat legit Dr. Ryan 5

Rami teretes, superne subangulati, cortice fusco cinerascente, sparse
punctati, glabri.

Folia petiolata, alterna, 4. 5. pollicaria, elliptica vel lanceolata-ellip-
tica, acumine brevi terminata, subcoriacea, glabra, obscure venosa,
supra nitidiuscula, subtus opaca.

Petioles

Paniculae, axillares, solitariae, folio breviores. Pedunculis parvis ac
ternis, distantibus, trifloris vel trifidis ramis trifloris. Flores glabri, pedi-
cellati.

Cinnamomum L. foliis trinerviis ovato-oblongis: nervis apicem evane-
scentibus fl. zeyl. 1454

Cinnamomum f. Canella Zeylanica Baud. pin. 400

Kiatou-Karva Rhed. mal. 5. p. 105. t. 53.

Cassia Cinnamomea Herm. huydb. 129. t. 655.

Cinnamomum foliis latis ovatis, frugiferum Burm. zeyl. 62. t. 27.

Habitat in Zeylona, in Mathura Tacquia 3

Rami cortice fusco superne compresso subangulati juniores saepe
uti gemmo villosi. folia petiolata opposita superiora inter-
dum alterna ovata vel ovato lanceolata. velle elliptico
lanceolata saepe 5 ad 6 pollicaria supra nitida subtus
glaucescentia ^{saepem} avenia. nervi tres plerumq^a basi orti
duo interdum ^{maximum} supra basin hinc interdum triplernervia
Peduncululi ex axillis supremis et apice ^{longi} flores raiemq^o
paniculati pedicelli et calyci roseo glanduloso adpressi

myrtha launi foliis ovatis trinerviis acumine longo, floribus axillaribus sessilibus congestis Lour. fl. Cochinch. pag. 251.

Caralla zeylanica sylvestris odore et sapore *Myrtha* floribus verticillatis Herm. Zeyl. pag. 26.

Cassa cinnamomea, myrtha odore, foliis trinerviis, subtus copio Plukh. Alm pag. 89.

Habitat in Cochinchina f. 7

Arbor 5 pedali, ramis supra. Folia crassa integerrima alterne glabra, supra obscure viridia, subtus pallida, petiolata. Flores albi, gemerati. Cal. communis 4 phyllis foliolis subrotundis concavis, basi exterioribus minoribus pilosis quinquefloris pedicellatis. Corolla 6 petala ovata, concava. filamenta 9, interiora glandulosa. Stylus brevis stigma obtusum. Basia ovata glabra perva. Perianth. proprium. 0

accedit ad involucentam

Chlorostylon foliis trinerviis ovatis coriaceis; nervis apicem attingentibus
Linn. S. V.

Chlorostylon foliis ovatis glabris rigidis trinerviis; floribus singulatis
Brown-jam 1817 Tab. 7. fig. 1.

Habitat in Jamaica &

Rami glabri teretes fusci folia petiolata alterna glabra
tenuissime venosa, nervis purpurascens. palmaria ve
ultra

_____ membranaceis
folia mihi videntur esse membranacea

curvifolia Laurus foliis oblongis obfete trinerviis intus
micis pelt comenalebus parvis Lour. fl. Cochinch. pag

252.

Habitat in fylvis Cochinchina.

Arbor magna. Folia acuminata, integerima, venosa, palli-

da, dura, nitida, alterna. flos albus. Calyx carnosus

lobatus infensis persistens. Stamina ^{de data} ~~superne~~ ^{foraminibus 4.}

chique quodlibet filamentum emittens filamenta 4. in quibus
anthera. Massa oblongo-ovata nigra

non vidi

Caryophyllus laurus foliis trinerviis alternis oppositisque, corymbis terminalibus ramosis, ramis bracteatis Lour. fl. Cochinch. pag. 250.

Habitat in sylvis Cochinchinae.

arbor mediocri. Folia ovato-lanceolata integerrima glabra

lesquipollentia, inferne glauca, petiolata, duruscula.

Cal. 0. Corolla 6 petala ovato-lanceolata. Stamina 4. 3 in
senora biglandulosa. Nectarium glandulosa. Baccina
nava ovata.

non vidi

polyadelphia Laurus folius obsolete Viniverris Carocolatis, floribus
axillaribus polyadelphis Less. fl. Cochinch. pag. 251

Habitat in montibus Cochinchinae 5

Arbor magna. Folia integerrima, plana, coriacea, venosopetio
lata, alterna, obscure viridia. Corymbi racemosi ~~ex~~ flore
albo rubri. Cal. lyathi formis coloratis, bracteis; laciniis oblongis,
erectis. Corolla nulla nisi calycem per diem. Nectarium glandula
6. pedunculata. Stamina 9. tria appendiculata in tres fascicu
los connata. anthera apice 4 perforata. Stigma obtusum.
baccia coronata minima fere sperma

castro L. foliis triplinerviis lanceolatis L'fl. zeyl. 146 + L. V. p. 303y

Cinamonum f. canella malabarica f. javanensis Baub. pin. 409

Pania Rheed. mal. 1. p. 107. t. 59.

Cinnamomum perpetuo florens, folio tenuiore acuto Burm zeyl.

63. t. 20

Habitat in malabarica, Sumatra, Java &

varietas forte Cinnamomi

Rami fissi. Folia opposita, firma glaberrima coria avenae
nervi parum supra basin subtus glauca petioli unguiculares ^{obtus.}
subtus con. vidi. Pedunculi ~~stellati~~ folio longiores

varietas modo Cauri cinnamom

folia opposita ~~ovatis~~ elliptico lanceolatis ve. triplinerviis subtus
glauces

Culclaban d. foliis Triplineris oppositio Lin. Mant.

Coelidlawan. Eph. N. B. Dei. 2. ann. 3. p. 53.

Cortex canophylloides Rumph. amb. 2. pag. 63. 7. 14

Habitat in India orientali

Camphora L. foliis triplinerviis lanceolato-ovatis

Camphora officinarum Baub. pin 500

Launus camphorifera Kompp. jap.

Arbor camphorifera japonica Boeyn. prod. p. 7. L. p. 94. u.

16. t. 2.

Commet. hort. 1. p. 185. t. 95.

Habitat in Japonia 5

Rami glabri cortice fusco purpurascente. Folia alterna 3-4
pollicari acuminata coria supra tuta obfusa venosa
nervis alter nervum supra alterum. petioli pollicari

— alternis

Cotoneflora Launi foliis elliptico lanceolatis triplinerviis axillis nervosa
subtus glandulosis villosis, floribus lateralibus axillaribus
Habitus dedit Schumacker
glabra tota. Folio petiolata 2-3 polliciana acuminata
utrinque acuta glauca vix venosa. flores pedicellati

glandulosa Laur. foliis ovatis triplinerviis, axillis nervorum subtus
glandulosis villosis

Habitat in insula Bourbonica Thourin 7

Simi glabri cortice cinereo. Folia petiolata, alterna ovata
attenuata 3 pollicaria Venius ^{obtusiuscula} sume venosa glaberrima
triplinervis (nervi alter panem supra alterum
axilla nervorum supra panem supra substantiam folii ele-
vata subtus fere villosa. Petioli purpurascens
pollicares. Flores axillares pluri

Sub

Tuberculatus L. foliis elliptico ovatis triplinerviis venosi racemis compositis
foliis longitudine folii ramisque glabris axillis nervorum villosis

Habitat in Canariis

Rami tuberculati folia membranacea utraque acuta interdum
parum acuminata 2-3 polliciana alterna versus apicem
confertiora subtus pallidiora. Pedunculi et flores ^{exteri} glabri

nervi apicem
non attingentes pedunculi parvales inferiores triflori superiores uniflori
subtus ~~marginibus~~ ^{marginibus} venis flores intus villosi

montana l. foliis triplinerviis ovato-acuminatis persistentibus, flombus
racemosis paniculatis Sw. prodr. pag. 65.

Habitat in Tamaulaco

Rami teretes superne angulati, foveis tomento tenui foveis grisei folia
attenuata superiora sequi pollicaria subtus oculo armato tomento
tenuisimo tecti supra arenaria

ramis subtomentosis fimbriisque
ramis subtomentosis canescentibus

~~Thunbergia~~
Thunbergii L. foliis nervosis lanceolatis perennibus, ramulis tuberculatis
floribus solitariis Thunb. fl. jap. pag. 173. glaucis
Habitata Japonia 5

non vidi

pedunculata L. foliis nervosis oblongis integris, floribus solitariis pedun-
culatis. Thunb. fl. jap. P. V. pag

Habitat in Japonia. 5

non vidi

Persea d. foliis ovatis coriaceis transverse venosis perennantibus, floribus
corymbosis L. V. Jacq. off. 37. Amer. 60. t. 264. fig. 92.

Persea Pluf. hist. 1. p. 2.

Persea americana Bauh. pin. 441.

Pyrus similis fructu in nova hispania nucleo magno Bauh. pin.
439

Arbor americana, amplissimo pergamenis foliis; superficiei nitidissima.
fructu pyriformi crustaceo coriaceo Conato Pluk. alm. 39. t. 267.

fig. 1.

Prunifera arbor fructu maximis pyriformi viridi, pericarpio esculento
butyraceo nucleum unicum maximum nullo ospiulo tertium Cengen
de Sloan hist. 2. p. 132. t. 222. fig. 2.

Habitatio americana calida

Rami teretes glabri, superne tomento tenuissimo grisei angulati. Folia
petiolata alterna 6-8 pollicaria venosa profert. in subtus
glabra juniora pubescentia. Petiolus sesqui pollicaris
Flores racemati ^{lymphi} ~~corymbi~~ versus apicem corymbi ramosi pedunculus compressus
si ~~est~~ in tomento tenuissimo incano in venter vndis

canaliculata Launus; foliis rigidis ovatis canaliculatis subtus quasi
pinnatifidis raris terminalibus sessilibus; e ramellis simplicibus
apice subtus floris Act. Soc. h. nat. par. 1. pag. 108.
Habitat in Amer. merid.

Benzoin. *L. folius enervis ovatis utraque acutis integris annuis*

Arbor virginiana, Citrea vel lemoni folio, Benzoinum pendens Com
Kort. 1. p. 109. t. 97.

Arbor virginiana, p. *shaminis folio, baccata Benzoinum sedolens Plum*
alm. 42. t. 139. fig. 3. 4.

Habitat in Virginia

92
pilosa ^{ovato-}Launus foliis oblongis enerviis pilosis, racemosis ^{Yermunellibus}
amplius Lour. fl. Cochinchinens. 257.

Habitat. in Cochinchina

arbor magna. Folia integerrima, acuminata, supra petio-
lata, quinquipolliciana. Flores polygami & in distincta
planta ^{eg. 6.} foliolis ovatis lanuginosis. Anthera & loculares. vid
hermaphro in hermaphrod: filamenta pilosissima. Stigma
2 fidum. Bacca globosa coriacea

non vidi

puberula laevis ramulis nervi rachibus minute puberulis. foliis
obovatis subabrupte ~~perpetua~~ acuminatis, rachibus axillaribus
et terminalibus subsessilibus dict. for. h. n. Par. 1. pag. 100
Habitat in Amer. merid.

non vidi

Conacea L. foliis ovatis acuminatis planis venis nitidis conatis, racemis
erectis foliis brevioribus Sw. prodr. pag. 65.

Habitat in Yamarea ramuli in axillis foliorum floriferi
Rami glaberrimi, supra striati. Folia petiolata alterna
distanta, tenuissime venis reticulata venis constantibus ab utro
3-4 pollicaria. Racemi in axillis foliorum ramulorum pedunculati
tri-floris, calyces rosei et alii oculo armato videndi subia
rescentes

an Doria
plantam a Pyreneo habui huius quam maxime simili
quoad folia diversa floribus minutis petiolis sub folio duplo
longioribus

— rami glabri

patens. foliis ovato-lanceolatis planis membranaceis, racemis erectis diffusis
foliis longioribus Sw. prodr. pag. 65.

Habitat in Jamaica ^{et} purpurascens

Rami glabri ~~tan~~ ^{sub} ~~laevigati~~ ~~et~~ ~~non~~ ~~non~~ ~~all~~ ~~tuberculati~~
folia supra nitida ~~et~~ venosa, glabra, basi acuta ^{et} interdum
rarius angustata saepe acumina brevi obtuso ~~et~~ terminata
calyces incani ~~et~~ pedunculi pariter patentissimi ramis
divaricatis

elliptico ovatis planis membranaceis supra nitidis, racemis
erectis ^{et} ~~longioribus~~ ramis ~~sub~~ ~~erectis~~ pedunculis pariter ~~erectis~~ dichotomis
racemis erectis ~~et~~ ~~compositis~~

Difformis *Launus* *Glaberrima*, foliis ovato-lanceolatis membranaceis laevibus acuminatis, promissis obtusis pedunculis floribus subcapellatis; defloratis incrassatis - *Difformibus*
Act. f. h. nat. par. 1. pag. 100
Habitat in Amer. merid.

exaltata L. foliis ovato lanceolatis coriaceis perennantibus planis; racemis
erectis compositis, calyce cyathiformi persistente Sw. prodr. p 65.
Habitat in Tamarica }

Coefornt. inter amer. Ryani sub nomine Cyathiform