

Lela

Erpina l. caule angulato-fimbriato Lin Mant. 124. * P. V. p. 045.
Habitat in Cap. bono p. 24

See

agrostata L. caule tereti pubescente Lin Mant. 25124. * S. V. pag. 255.
Habitat in India orientali.

Lin

crispa L. petiolis crispato alatis tetragonis
Leca crispa Caulis angulato fimbriato her. Mant. 124

Habitat in Cap. b. Spei t

Folia petiolata pennate bipinna - foliola petiolata opposita
oblonga grosse fissurata oblique transverse nervis, inter
nervos venis striis subtilis albidis vix tamen villosa circumscissa.

Folia ter nata vel quinquies pennata her. Thonn.

Lema Celtis foliis ovato-lanceolatis acuminatis oblique cordatis serratis superioribus

Rapernimis Sw. prodr. p. 53

Lati arboris foliis anguste arbor baccifera Sloan. hist. 2. p. 80

Habitat in Ind. occidentali

Lambucina A. Ken. Mant. 211. * P. V. p. 243

Sapindus indica foliis bipinnatis: infemus ternatis: seminis quinis
Burm. Ind. 73. t. 24. fig. 2.

~~Habitat in Java~~ ^{in oppido adle Tereti pubescente} Ken. Mant. May 124

~~equata~~ ^{equata} ~~seca~~ ^{seca} petiolis femiberebibus supra canaliculatis

Habitat in India orientali

Rami teretes folia petiolata alterna pedalia, subbipin-
nata ^{bi p. 4 p. 4} foliola inferiora saepe ternata: petiolata

opposita ^{elliptica} ~~ovata~~ ^{transverse} lanceolata nervosa, ^{opposita} ~~ternata~~ acuminata

5-6. pollicaria, simpliciter venosa juniora subtus pilis alipis
Cyma terminalis trichotoma