

Leonurus[?]

1. Cardica L. foliis ^{floriferis} ~~linearibus lanceatis~~ ^{fidis} ~~trilobis~~ calycibus pediculis
2. crispus L. foliis omnibus acutis serratis nigrosermis margine inaequaliter reflexis
caulis quinquelobis
3. Sibiricus L. foliis ^{trifloris} ~~tripartitis lanceatis~~, calycibus vellosis, floribus ~~calycibus duplo longioribus~~
4. Sibiricus L. foliis ^{tripartitis} ~~tripartitis~~ multifidis lanceolatis obtusis, floribus
^{calycibus duplo majoribus.}
5. mammosum L. foliis ovatis lanceolatisque serratis, calycibus sesquialteris
spinosis

crispus L. foliis omnibus acute serratis, rugosis semis margine inaequaliter
reflexis, caulibus quinquelobis. Mur in Com. Gott. 1777. p. 44. t. 4.
S. V. pag. 530.

~~Carduus Turphutensis~~

~~maritimus curvatus. Dietrich. Bot. t. 1. p. 230.~~

~~Carduus crispus Raj. hist. 572.~~

Habitat

non vidi

carduac L. foliis caulibus lanceolatis trilobis L. V. fl. Dan. 7. 727

Carduac Duchf. hist. 395

Marrubium carduac dictum Bauh. pin. 230

Habitat in Europa meridionali &

Caulis erectus interdum biumbellatus, simplex vel ramif. Folia petiolata
patentissima infera

superiora floralia trifida laciniis lanceolatis serratis summis
integerrimis attenuata, omnia utraque leviter pubescentia
subtus albida, venosa. Caliculi 14 flori cincti involu-
cri subulatis calycibusque rigidis pungentibus pilosiusculis

varietas hirsutissima an crispus Merr.

Tataricus L. foliis tripartitis laciniatis, calycibus villosis her. S. V. p. 5304

Hort. Ups. 1714

Cardua foliis tenuis et profundius incisae glabra s. villosa

amm. nith. 49. 50 mill. ix. 00

Habitat in Tataria ♂

Species her. Cardui. Colia Paulis Erectus leviter pubescens
Colia petiolata inferiora quinquefida laciniis lanceolatis
incisae superiora trifida ultima indivisa lanceolata
attenuata omnia subtus villosa, Calycis pungentes

Siboncus L. foliis tripartitis multifidis linearibus obtusiusculis Lin. S.V.

1704

Ballote inodora, foliis coronopi Am. r. th. 40. Y. S.

Habitat in Siberia, China &

Caulis pubescens. laciniis foliorum ~~in~~ extusum in ~~is~~ a
supra glabra subtus ~~laeviter~~ ^{parva} ~~tame~~ villosa mollia ultima
femina linearia ~~integris~~ ^{parva} flores triplo majores quam in
cordata

var setos prostrat. corollis majantibus Gmel. hb. 3 p 239.

manubraftum L. foliis ovatis lanceolatisque serratis, calycibus sessilibus pino
si Lin. T. V. Mant. 412. Faerg. fl. austr. 5. T. 405.

Manubraftum foliis cordato Bosc. mus. 2. p. 120. T. 90.

Habitat in Bohemia, Ucraina et Java

Caulis pubescent. folia petiolata supra glabra trinervia
subtus molliter villosa carinata, trinervia attenuata bi
pollucina serraturis utraque 2-3 remotis magnis. verticilli
multi flori. involucri petalae calycis dentes petalae sub
perungentes.

Leon. membranaceum a Walden. distinguitur genere sub nomine Chaetoni
ob calycis laciniis erectis et corollae labio superiore non plano ut in heo
nunc sed Hornuato differentia sane haud magni momenti