

Lindsea

Lindsoa

Ryan.

of the
met.

pellucida Melastoma foliis lanceolatis integerrimis glabris, panicula terminali
muc. in pediculis in videri
racemosa dichotoma floribus ^{petalis} ~~subdichotoma~~ ^{subdichotoma} subdecimpositis

refescente Habitat in Jamaica. Ramuli ^{petalis} ~~subdichotoma~~ ^{subdichotoma} subdecimpositi compressi
in pulvere Ramuli obscure tetragon. folia petiolata 3-4 pollicaria basi acuta, utrinque
lenti ^{attenuat} ^{probas nervi margini}

tres vel quatuor divariata. Flori minuti lanosif. Bractea petalica ad basin florum
in singulis ramulis ^{si quis epist. Anos} ^{petalis} ~~subdichotoma~~ ^{subdichotoma} subdecimpositis
cum floribus ^{neg.} ~~subdichotoma~~ ^{subdichotoma} subdecimpositis
indivariata ^{neg.} ~~subdichotoma~~ ^{subdichotoma} subdecimpositis

affinis *Corryata*

adulescentia supra glaberrima, subtus in nervis et venis subprelovente
juniora utraque adpressa pulvere sordida cinereo venoso subrugosa
petioli 3-4 pollicari compresso tetragon

pubescenti ^{3-4 pollicari} ^{compresso tetragon}
Racemus terminalis compressus, pedunculi terminalis saepe terni
unico multo brevior interdum gemini et loco pedunculi tertii

ramus uterque et crescit. pedunculi ultimi apice ^{petalis} ~~subdichotoma~~ ^{subdichotoma} subdecimpositi
lesiles, secundi 3-4 in singulo ramo quinto sessilis ^{subdichotoma} ~~subdichotoma~~ ^{subdichotoma} subdecimpositis
Rohr no 102. sub nomine M. racemosa e Martynia

Petala oblonga
Hamula decem corolla longiora

Ludaea

Aubl. fl. Gu: Vat 364

Habitat in Guana herb Guaf. et Ruel

Stipites Vetrogoni: Ervum simpliciter pennatum, ternatum et bipun
^{nerium omnes fertiles}
foliola et pinnulae alterae: pinnulae petiolatae rhomboides, aut sic rotunda
ae, semipollicares.

Epistola Belkensis

Rehania