

Linociera

10.

Gen. pl no 1709. Swartz. Fl. Ind. Occid. p. 49

Cal 4-dentatus. Cor 4-petala. Anth petala 2. opposita basi connec  
tenfer. Bacca 2 locularis; loculis 2-permis, exsuda.

L.

~~25~~  
~~24~~

Linociera.

Cal 4-dentatus. Cor 4-petala. Anth sessilis; petala  
2. opposita basi connedentes. Bacca exsucca, 2 locularis.  
loculis 1-permis. Swartz.

~~non Chococum colligand. et forte cum ea conjugenda~~

floribus paniculatis, floribus pedicellatis, petalis reflexis  
Ciguatrina 1 Linociera paniculis terminalibus  
foliis lanceolatis,

Thouinia Ciguatrina; foliis lanceolatis, paniculis terminalibus, flori-  
bus erectis. Swartz. prodr. 15

Linociera Ciguatrina. Swartz. fl. ind. occid. 1. p. 49.

Habitat in fruticetis, apud Jamaica, Hispaniola. 5

Rami teretes, glabri, cinerei, punctis elevatis adspersi. Folia bi-tri-  
polliticaria, apice obtusa, basin versus attenuata, coccinea, nitida.

Pedunculi <sup>partiales</sup> bis-ter dichotomi. Bractea minuta, ad basin pedicellorum.

Calycis dentes ovati. Petala alba, linearia, obtusa, concava, ~~fla~~  
reflexa, decidua. Filamenta sive ulla. Swartz. l. c.

Ramus teres glabrus

*Swartz. fl. ind. occid. 1. p. 49.*

caulescens Gunnera caulescens foliis squelobis, racemo terminali.  
Pank e caulescens Molina Hist. Nat. Chil. p. 120  
Habitat Chili

non vid.

latifolia. 2 ~~Linnaeus~~ <sup>paniculatis floribus. paniculatis</sup>  
~~Domingensis~~ <sup>racemis terminalibus atillatis</sup> ~~Chionanthus~~ <sup>pedicellis trifloris</sup>  
nifloribus sessilibus, petalis lanceolatis obtusis, foliis ellipticis acuminatis  
Chionanthus Domingensis; foliis ovatis utrinque glabris, panicula terminali  
subcymosa, calycibus lobis, <sup>lamarck</sup> ~~illust. 30. t. 1. p. 30~~  
Habitat in St Domingo Desfontaine & in Antillis Richard 5

Rami cortice cineres, apice in versus alternatim compressi. Solia ~~subulata~~  
<sup>et in pollicaria</sup> ~~elliptica~~ <sup>lanicolata</sup>, acuminata, acumine obtuso, subovata, avenia  
tenuissimis transversim demum nervosa, laevia, utrinque glabra. Petali  
basi fascii, superne uti certa subtus parvulae purpurascens. Pedunculi  
foliis breviores: parvales patentissimi. Bractea subulata, villosa in carina  
Calycer oculo armato, <sup>marginem</sup> vix villosa. Petala unguicularia, utrinque ejusdem  
latitudini, plana, <sup>obtusis</sup> ~~antheris oblonga~~. Intra oblonga, majoritudine pisi. Nunc

libularis.

A ligustrina differt foliis multo latioribus, <sup>minime nitidis</sup> firmioribus, acumina-  
tis. Petalis latioribus. An eadem cum Chion. incruscata Swartzii? ~~petalis~~

# vero non da-  
rda.

~~Habitat Rami teretes, solia subulata, glaberrima, integerrima, obtusa, pubes-~~  
~~centa.~~  
~~Genus facie pisi perquam similis Chionantho, An loculamentum unicum~~  
~~in variis Chionanthis exaristis naturalitate? nullum~~



3 *Linociera*  
cotinifolia ~~Chionanthus~~ racemis subcompositis, foliis glab. pedicellis trifloris floribus  
pedicellatis, petalis linearibus, foliis glab. subtus villosis  
Arbustula zeylanica, cotini foliis subtus lanugine villosis, floribus albis  
cuculli modo laciniatis Plukh alm. 44. t. 241. f. 4.

Arbor zeylanica, cotini foliis subtus lanugine villosis Burm. Zeyl. 31.  
Chionanthus zeylanica; foliis ovatis subtus villosis, drupis obsvatis. L.  
Lamant. Mont. illust. p. 30. t. 9. f. 2.

Chionanthus cotinifolia; panicula terminali trichotoma, pedunculis  
trifloris, foliis obtusis Wied. spec. pl. t. f. 47.  
Habitat in Zeylona 5

Salde affinis sequenti

fragrans Olea foliis lanceolatis serratis, pedunculis lateralibus aggregatis unifloris.  
S.V. p. 57. Thunb. Gap. 10. S.V. p. 57. Tab. 2. S.V. p. 57.

Mart. K. f. J. V. Amph. Aman. Ex. f. f. v. p. 844.

Habitat in Japonia

Rami et ramuli <sup>interdum interdum</sup> trichotomi, obsoleta tetragoni glabri cortice  
cinereo. Folia <sup>interdum interdum</sup> petiolata opposita attenuata, basi acute  
margine subreflexo, parallelis nervosa obscure venosa supra  
nitida subtus pallidiora digitalia. Pedunculi axillares  
6. 8. unguiculares,

Osmanthus Laureus.

purpurea 4  
Zeylanica 4

~~Linaria~~ ~~Chionanthus~~ racemis subsimplicibus, foliis obovatis calycibusque glabris  
pedicellis trifloris, <sup>nerviis</sup> ~~glabris~~ ~~sepalis~~ ~~sepalibus~~, petalis canaliculatis foliis subovatis la-  
ganiethagar s. arbor bovina Herm. Zeyl. 33. 23 (Clypeibusque glabris

Chionanthus zeylanica; pedunculis paniculatis multifloris Lin. Fl.  
Zeyl. 14. ~~Syst. Veg. 97~~ Syst. Veg. 97

Thouinia nutans Lin. Suppl. 89.

Chionanthus purpurea. foliis ellipticis, glaberrimis, floribus purpu-  
reis nutantibus <sup>Cammerl.</sup> ~~La Mart.~~ illust. 1. p. 30 24.

Habitat in Zeylona

Rami cortice cinereo, adpersi punctis elevatis nigri acutibus, <sup>supreme cinereis</sup> ~~asperis~~ ~~venis~~,  
supreme altero latere compressi, ~~sub~~ foliis dilatati. Folia breviter petiolata  
<sup>acuta</sup>  
~~obovata~~, <sup>obovata</sup>, vel ovalia, basi parum attenuata, acumine brevi terminata,  
obscura venosa, lœvia, ~~stimpis glabris~~, pubes minimè nitida, subcoriacea.  
Racemi laterales et terminales, solitarii, oppositi, foliis breviori, plerum  
que simplicibus, pedicellis trifloris, cernuis. Bractea brevis, linearis, <sup>ad basin sin-</sup>  
guli pedicelli. Petala breviora quam in reliquis speciebus, ~~et~~ ~~tra~~ ~~crassiusculis~~,  
et rorsum sensim angustiora. Anthera lineares, petalo dimidio breviores.

# utraque



emarginata Olea foliis obovatis emarginatis, panicula terminali pauciflora  
La mant illustr. pag 2 tab. 8. fig. 2.

Habitat in madagascaria Herb. Desfont. 7. usque 5

arbor 40 pedali  
Rami glabri, cortice cinereo. Olea opposita, petiolata, bipollicaria, obovata,  
obtusissima, emarginata, conacca, arena, <sup>citrus glabra</sup> chersia, lora, nitida, integerrima. Dupa edu  
lis, magnitudine fere nucis juglandis

flavicans Limociera paniculis axillaribus, petalis ovatis, foliis obovatis  
obtusis +

Thouinia flavicans; laciniis corollae ovatis Willden. herb. mans. 3.

Habitat in --- +

Paniculo opposito: pedicelli apice incrassati. Bractea duae,  
minuta, vides ad dichotomiam paniculo et sub apice pedicel-  
lorum. Calycis foliola ovata triangularia. Corolla coriacea,  
flavescens: petala erecta, concava. Filamenta nulla: An-  
thera basium non laciniarum inserta: Willden. l. c.

Quamvis fructus mihi ultimum trium mihi ignotus, potius  
Limociera generis esse quam Chionanthi inde judicis est  
antheris sessilibus basi conneis petalis conneis huius judi-

CO  
genus ~~est~~ Chionanthi affinis, et nonne nonnulla  
huc pertinent?

Habitus. Arbores. Rami teretes. Folia petiolata, opposita,  
integerima, obtusa, submarginata.

*Chorophylla* V.

Habitat in hinc herb. Throum

Caulis erectus, striatus; uti tota plantaglaber. Folia petiolata, pennata, 2-3  
juga opposita: foliola sesquialtera oblonga, uncinata, utrinque utraque dante, semi-  
pollicaria, alterna remota. Pedunculi axillares, superae, folio parum brevis  
vel. Flores in apice parvis.