

Macaglia

alba Macaglia folliculis lobatis

Habitat in Guiana ab inventore Pichard botanico condiscipulo suo
meum inter pl quam plurimas rarissimas plantas America com-
municata. 5.

Arbor. Rami teretes; superne, uti folia subtus pedunculi et calycis
e pulvere subtilissimo cinerei. Folia petiolata, 3 pinnata, tri- & quadrifida
cuneata, undique ejusdem latitudinis, obtusissima, retuso, integerrima, supra
glabra, ^{glabra} laevissima, nitida; subtus oblique multinerviis linearibus, nervis
si, ^{glabra} interpositis venis simplicibus, tenuissimis obscuris. Pedunculi termi-
nales, plures, compressi, uti patulae; versus apicem sensim dilatati
apice subditotomi. Flores in apice pedicellorum, aggregati, subsessiles.
Bractea parva ad basin calycis. Flores parvi. Calyx campanulatus, quin-
quefidus; laciniis linearibus-oblongis, obtusis. Corolla tubulosa, ^{glabra} calyce duplo lon-
gior, quinquefida: laciniis patulis, ante expansionem spiritaliter contortis,
explicito erecto patentibus. Filamenta brevissima, tubo corollae adnata: An-
thera globosa. Germen superum, compresso-obovatum. Stylus unicus. Colli-
culus orbiculatus, ^{centipetans} acuminatus, plano compressus, orbiculatus, acuminatus brevi

terminatus, laxis, diametro tripollucari. Semina lenticularia, bracteata, via polli-
 cania, marg. valvuli parallela, altera sutura affixa, margine lat membranaceo
 tenui lat ^{latit} ~~latit~~ ^{magis} ~~magis~~ ^{adine} ~~adine~~ valvularum latitudine cincta, extus medio parum
 convexa, ^{inter} ~~inter~~ ^{ad} ~~ad~~ ^{insertionem} ~~insertionem~~ funiculi umbilicali parum concava, mar-
 gine emarguratione parva incisae, cinerascens, oculo armato ³⁴⁶ porulosa.
 Affinitatem quadam habet cum Allamanda, differt vero corollis, semi-
 nibus laxe non biserialis valvulis minoribus

~~formos. foliis pubescentibus, corollam gallo barbato, calycibus trifloris, caulis
 frutescente & herb. Sup. nov. p. 41. d. 21.
 fovea reniformis glomerat. fol. p. 15. d. 2.
 et alibi in p. 5.
 Ramis teretes glabris. Calyx heterostachyus hypochlorici orthostachyae
 notae figura glabra venosa. tubus a velli terminatus
 campanulatus. Calyx pubescens. Corolla pulchre ^{colorata} ~~pubescente~~
 lobis calycis oris acuti.~~

Tuberculata Maraglia folliis tuberculatis

Habitat in Guiana Aubard 5

Alta Macaglia folliculis levibus

Habitat in Guiana Richard 5

~~Arbor~~ ~~fraxinea~~ ~~terrestris~~, superne, et folia subtus, pedunculata et calyces
pulvere sublimissimo cinerari. Folia petiolata, sparsa, bi- sive tripollicaria
antiqua fere eorum
oblonga, obtusissima

tuberculata Macaglia

*ridui germanici Rumph. amb. C. Tab. 45. f. 1. Calyx superius
 uti corolla similis Oldenlandis, Tetradia monogyna. baccis
 oblongis, locis luteo aurantia, disperma semina linearis
 oblonga, utraque acuta, dura
 Ruvia sylvestris Burm-gard. Dumetorum vide Rehd 4*

*Cataracta V. rarisima Vermiculata flexuosa
 Cane colata serrata Kraft Herb. pag. 3.
~~Abd. K. in nov. Zeelandia 5~~*

Xra Xera