

Maivocnemum

Corolla subampullata. Capsula biloculari; valvula lateribus externe  
dehiscentibus. Semina umbriata Sw.

1. jamaicense M. <sup>panicula</sup> paniculata <sup>antheribus</sup> aphyllis, foliis lanceolatis, ovatis
2. candidissimum M. <sup>panicula</sup> paniculata <sup>antheribus</sup> inchoatis, foliis subrotundis, foliis ovatis
3. coccineum M. <sup>panicula</sup> paniculata <sup>antheribus</sup> foliis coccineis ellipticis, foliis lanceolatis  
ellipticis

4. intercoros celloso. Dissepimentum contrarium. Semina minima margine  
obsoleto cincta. Receptacula obovata, dimboformia, callosa fl. penes

2. pug 40

corolla infundibuliformis tubus parum curvatus. Fauces dilatatae  
Stamina basi villosa. Antheris basi bifidis. Capsula truncata umbri-  
ciata utroque sulcata longitudinaliter dehiscenti valvula duplicata

1, jamaicensis Macrocremum Am. Acad. s. p. 413\*. L. V. pag. 212. Sw. obj. pag. 60.  
Macrocremum arboreum, foliis ovatis oppositis, racemis sustentaculis  
longis infidentibus Brown. Jam. 165.

Habitat in Jamaica &

Inter ramos compressis inaequalibus. Folia opposita, lanceolato-ovata, subre-  
violata, integerrima, teria. Panicula trichotoma, dichotoma, vix foliis  
longiora haec Am. l. c.

arbuscula caule ramoso teri. rami longi laxi teretes, ciliati. Folia  
approximata, opposita magna, oblonga acumine brevi, nervis  
integerrima utrinque glabra. Peduncululi axillares, oppositi  
foliis longiores, glabri, tripartiti trichotomi flore subsessiles  
secundi.

Coccineum Macranemum racemi foliis coloratis ellipticis, foliis lanceolato-ellipticis  
Symb. bot. 2. pag. 30. Tab. 29.

Folia supra subarenia, subtus venis subsimplibus inter nervos.

Racemus bipedalis erectus. Pedunculus communis angulatus, glaber,  
fuscus: partiales inferi oppositi.

Folium florale semper unicum in superioribus quandoque 2 vel 3 in  
inferis

a. flos b. corolla aperta c. calyx cum pistillo d. pistillum e. capsula f.  
eudem transverse secta

Venerum Macrocnemum

Macrocnemum venosum foliis oblongis venosis linearibus; nervo  
venis que pubescentibus; racemis terminalibus; floribus sessilibus.  
Fl. Penn. 2. pag. 49. Tab. 190. fig. 6.

Habitat in Peninsula nemoribus inis calidis &  
Caulis frutescens, teretis, exilis, glaber. Rami teneri, obsolete tetra-  
goni. Folia opposita, petiolata, oblonga, acuta, integerrima, sene-  
scima, supra lineata, subtus reticulata, dorso senu que pubescentia,  
tripalmata. Stipula interfoliaceae, bipartita: laciniis lanceo-  
latis, acutissimis, adpresso, pediculis duplo longioribus; membranaceis,  
striatis, pubescentes. Racemi terminales bracteati, multiflori  
compositi, pubescentes. Peduncululi bractea sub pediculis  
seder sessiles. Calyx minimus. Corolla parva alba fl. Per

folia oblongo ovata

*Corymbosum Macrocnemum*

*Macrocnemum Corymbosum*, foliis ovato-oblongis basi excavatis laciniatis sessilibus pluriatis coriaceis, corymbis magnis fl. Peruv. 2 pag 40  
Tab 109.

Habitat in Andium collibus et in catervis calidis 3

Caulis <sup>perier</sup> fruticosus, quadrangularis, simplicis, erecti, inferius nudus,  
superne foliosus, teretes, saepe granulis adspersi, cinereo-fusco variegato  
Folia opposita horizontalia, sessilia, pedicula sulcata, ampla, ovato-  
oblonga, basi <sup>subcordata</sup> excavata, <sup>subacuminata</sup> caninata, pluriata, laciniata, integerrima, coriacea  
supra nitida, venosissima. Stipula interfoliacea, bipartita: lacina  
lanceolata, adpresso-membranacea, lineata. Corymbi terminales  
magni, bracteati, <sup>mitrotomi</sup> multiflori. Bractea minima, lanceolata sub  
pedunculis et pedicellis. Calyx purpurascens, basi plerumque gra-  
nulosis, coriaceus. Corolla crassa, intus alba, extus oblique pur-  
purea, fauce villosa. Capsula fusca-purpurascens, tandem  
atris fl. Per.

*Microcarpon macronemum*

*Macronemum microcarpum* foliis oblongis obtuse acuminatis & subtus  
pubescentibus; racemis terminalibus; floribus congestis & sessilibus fl.  
Peruv. 2. pag 49. tab 100. fig. a

Habitat in Penuria nemoribus in calidis  
Caulis fruticosus, quadrangulis, Rami fusci: juniores foliosi, vix striati,  
t, obsolete angulati, obscure purpurei. Folia opposita, petiolata, paten-  
tia, oblonga, nonnulla oblongo-ovata, obtuse acuminata, integri-  
ma, supra glabra, nitida, & subtus pubescentia, venosa: venis infle-  
ctis. Petioli breves, pubescentes. Stipulae supra axillares, interfolia-  
riae bipartitae. <sup>concolorata</sup> lacinae acute, petioli dimidio breviores; adpro-  
ssa membranacea. <sup>racemi</sup> racemi terminalis; compositi. Forpalmanes. Punc-  
tuli bracteati sub singulo bractea. <sup>144</sup> Flores congesti sessiles; bracteis ma-  
joribus distincti. Calyx parvus, & quidentatus. Corolla alba, calycis gub-  
major: lacinae reflexae. Capsula turbinata, parva. Semina inveni-  
tissima fulvescentia fl Per.

2 Candidissimum Macrocnemum <sup>corymbi</sup> panicula trichotoma. foliis subrotundis,  
foliis ovatis

Plantam rarisima in vicinia St. Marthae legit ab von  
Rohr.

Arbor ramis teretibus, glabris, oppositis, superne articulatis,  
compressis, subfoliis dilatatis.

Folia petiolata, bipollicaria adultra, opposita, ovata, alu-  
minata, obtusa, venosa, integerrima, supra glabra,  
subtus pallidiora, venis villosa.

Pedunculi ex apice ramorum, terni, anticipites, superne  
dilatati, bi trifidi. Soliola duo parva sub prima  
divisione pedunculorum. Peduncululi pariter tri-  
ra peduncululi communi. Pedicelli triflori. Flores  
laterales subpedicellati, intermediis sessilibus, pollicari

Solum petiolatum, obovatum vel subrotundum, albu-  
membranaceum, venoso-reticulatum candidissimum  
a margine calycis lateralium corymbi  
duo in singulo corymbo

Capsula oblonga, subcompressa, bisulcata, singula valvula linea  
elevata  
Semina ut in antecedente



Colobineum Mairocinemum <sup>racemum</sup> paniculato foliis coloratis ellipticis, foliis

lanceolato-ellipticis

7<sup>a</sup> 4<sup>a</sup> insula Trinitatis Legit Dr von Kober.

Arbor ramis pilosis.

Bolia pedicelata, pedalis, basi vix lat palmam lata, basi angustata, extrorsum latiora, membranacea, integerrima supra glabra, subtus costa pilosa.

~~Pedunculata~~ ~~in racemosis~~ pedalis, pedunculata, oppositi

Racemus terminalis, pedalis, pedunculatus & corymbis parvis oppositis pedunculatis. <sup>compertus</sup> pedunculus <sup>nautilus</sup> trifidus, pedicelli laterales quinqueflori, intermediis trifidis. Bractea ad

basi pedunculorum parvulum ovata, acuminata, concava, ciliata: pedicellorum lanceolata hirta.

Calyx perianthium monophyllum superiorum quinque sedum: lobis <sup>tergum</sup> <sup>pericarpium</sup> rotundatis. Foliolum petiolatum ellipticum, perspicuum, venoso-reticulatum, oppositum, basi acutum glabrum & ~~marginibus~~ <sup>marginibus</sup> calycis in singulis corymbis ~~et~~ <sup>et</sup> ~~marginibus~~ <sup>marginibus</sup> lateralibus corymbi ut in mesependa frondosa

arabicus Pistis suffruticosus stipulatus, foliis alternis lanceolatis planis lobatis  
Len. s. v. pag. 100

Helianthemum creticum Linano folio flore croceis Tournef. Corol.

Decumbens basi ramorum elongati  
demum apice <sup>inter</sup> fructus ~~vna~~ <sup>vna</sup> ~~fructus~~ cinereo <sup>vna</sup> ~~fructus~~ rufescenti. Ramuli pollicari alterni

caulis apertus <sup>fructus</sup> <sup>ramulorum</sup> <sup>compertis</sup> <sup>rectioribus</sup>  
ramuli <sup>sub</sup> <sup>stipulis</sup> <sup>obtusiuscula</sup> <sup>obtusiuscula</sup> <sup>rigidiuscula</sup> <sup>glabra</sup> <sup>are</sup>  
floriferi terminalia <sup>obtusiuscula</sup> <sup>linearia</sup> <sup>ultimi</sup> <sup>terminali</sup> <sup>cinere</sup> <sup>stipulis</sup>  
et latere <sup>ovatis</sup> <sup>minuti</sup>. Clavula remota <sup>lobis</sup> <sup>pubescentia</sup>, stipulis

ad apicem <sup>lanceolatis</sup> <sup>terminalibus</sup>  
et absque ramulis Pedunculis <sup>paucis</sup> <sup>vel</sup> <sup>quatuor</sup> <sup>solitariis</sup> <sup>versus</sup> <sup>apicem</sup> <sup>polli</sup>  
inferne appropinquata <sup>caulis</sup> <sup>et</sup> <sup>sepe</sup> <sup>ramis</sup> <sup>crassius</sup> <sup>sub</sup> <sup>ramis</sup> <sup>polli</sup>  
membra angustiora <sup>his</sup> <sup>membranatis</sup> <sup>extrema</sup> <sup>lanceolata</sup>

stipulis minoribus  
superioribus distantibus  
stipulis majoribus

Pedunculi oppositi foliis versus apicem tantum pubescentes

#

Corolla mono petala, infundibili formae, quinquesida. laciniis  
 lanceolatis obtusis, glabris. tubus calycis duplo longior  
Stamina filamenta quinque in fauce tubi, longitudine corollae,  
 intus pilosa. antherae lineares, extra corollam  
Pistillum germen inferum turbinatum, angulatum, desutum.  
Stylus in crassatus longitudine corollae. Stigmata duo obtu-  
 sa, reflexa  
Capsula obovata, biloculari, polysperma  
Semina membranacea, subulata, <sup>imbricata</sup> receptaculo lineari affixa

(An macrocarpi generis cum corolla non campanulata? in pla-  
 nis convenit id eoque potius huc retuli. ~~et missenda differt quanta~~  
~~potius quam a differt modo haec nota.~~ ~~et missenda a differt~~

712

Arabis suffruticosa: Spicatae in albae  
 Corollae long. v. long. 500  
 florum

pallida Ruellia ped. foliis petiolatis undulato-crenatis, <sup>margine lobis) floribus</sup> ~~pedunculatis~~ (axillaribus  
unifloris solitariis subtulibus

Ruellia strepens; flore magno violaceo; foliis obtusis Cosm. Desimp.  
pag. 114.

Caulis lissus obscure tetragonus, <sup>secco</sup> ~~laevissimus~~ in junctis latus crenatus  
Folia petiolata, pollicaria, ovato-oblonga, obtusiuscula, pallide ~~viridi~~

Pedunculi <sup>axillares</sup> ~~axillares~~, solitarii, <sup>uniflori</sup> ~~oppositi~~, brevissimi, <sup>calice lanceolato utriusque</sup> ~~calice lanceolato utriusque~~  
fol. ad basin calycis ciliato, calyce longior.

Corolla tubus calycis triplo longior superne amplectus quadruplo  
latis pollicaris

Differt. a R. strepens calycibus longitudine tubi.