

Mari-gravia

1. umbellata M.

Thlasium Louren- hui pertinere videtur

mari-gravioidei

Marigravia

Marigravia umbellata, foliis ovato-oblongis acuminatis venosis

Marigravia scandens, fructu radice in posito Plum. Ind. 173.

Marigravia umbellata Lin. Syst. Veget.

Lema Solanum folis elliptico lanceolatis supra scaberrimis subtus ~~lanceolatis~~
~~base Cyma terminalitate pedunculato. Tomentosi~~
~~Habitat in Cayenna. v. Boiss. no 54. 5.~~
~~inermis Ramis scaberrimis ut petioli folia subtus; pedunculis calycis ^{concoloris}~~
~~que ^{est} ⁱⁿ ^{impetigione} ^{aspera} ⁱⁿ ^{canis} . ^{valida} ^{folia} ^{base} ⁱⁿ ^{petiolum} ^{attenuata}~~
~~et ⁱⁿ ^{sparsis} ^{frequentibus} ^{acutis} ^{areolaris} ^{nervosa} ^{supra} ^{secundum}~~
~~nervos lineata & tuberculis minutis numerosis scaberrimis subtus~~
~~striatula. Pedunculis ^{terminalibus} cum ^{ramis} ^{ulceris} ^{erectis}~~
~~scaber ³⁻⁴ ^{pollicari} ^{cyma} ² ^{sterilifera} . Flores parvi ^{lacini}~~
~~calyce ^{involucris} ^{ovato} .~~
~~Corolla profunde ⁵ ^{fida} .~~

Mangrovia conacea, foliis ellipticis obtusis coriaceis arenis

Habitat in Cayenna von Probr. 5

Rami glabri, cortice cinereo, punctis raris et vatis adspersi, inferne vese-
tes, superne subtergioni.

Folia breviter petiolata, quadri-vel quinque pollicaria, utrinque gla-
bra, laeva, margine parum reflexa.

Pedicelli versus summitatem pedunculati communi, verticillati, septemde-
cim vel octodecim, petentissimi, bipollicares, aequales, vese-tes-
versus florem incrassati, et punctis frequentissimis sparsis tuber-
culati. Flores ascendentes

(De absentia vel appropria corporum utricularium in ista specie esse asseverat Ryan, cum in unico specimine, quod inter reliqua Probrii invenit deessent. Si autem ex vestigiis judicare licet mihi verosimile videtur, quod adferentiaque adsunt in hac specie ac in altero

umbellata *Marcgravia* Lin Sp. p. 719, Jacq. herb. Tab. 96. Riv. obs. pag. 205 &

Marcgravia scandens, fructu radiati profuso Plum gen. 7. w. 173.

Phyllitidis scandenti affinis major, folio crasso subrotundo Sloan
herb. v. p. 74. t. 20. fig. 1.

Marcgravia scandens, foliis caulinis subrotundis ad margines glandu-
latis, ramorum integris ovatis alternis diffusis profusis, floribus
umbellatis terminalibus Brown. jam. 244. t. 26.

Habitat in America calidiorie &

non peroptica, sed incitis more quarendam felicem radi-
cani. Ad ultior caulis frutescens, radiculis arbori in quibusdam
ad haerens. Rami superne, lateri penduli. Folia planda
juniori radice antea subrotunda, subsimplex leviter emar-
ginata, margine integra punctis glandulosa. ramo-
rum diffusis alternis breviter petiolata, ovata
lanceolata l. ovata, lanceolata, consistens, gla-
bra. Umbello nutante ped. uniculis simplicibus uni-
flori. Glandula utriusque in centro umbello pyle-
nabulo oblongo, ventricosa, galeiformis, conato-
su.