

Marynicarpus

Flor. Penn. prodr. p. 77. 133.

Cal. 4-5 partitus. Cor. 0. Stigma peltatum. Drupe nucleos <sup>1</sup> sper

mo

69  
71

Marynicarpus. Cal. 4-5 partitus, superus. Cor. 0. Stigma peltatum.  
Drupe nucleos 1-spermo.

calyx superus.

setosus, Margynicarpus

Empetrum pinnatum; foliis pennatis, folioli linearibus subulatis, petiolo com-  
muni basi dilatato vaginali, floribus divisis La Martk. Dict. bot. 1. p.  
567.

Anisotrum barbatum; folioli linearibus subulatis apice barbati, flori-  
bus axillaribus. La Martk. illust. bot. 1. p. 77. ~~177.~~

Margynicarpus setosus; foliis impari pennatis: folioli linearibus,  
floribus axillaribus sessilibus Flor. pers. 1. p. 20. t. 8. f. 2.

Margynicarpus setosus; foliis apice barbati, floribus hermaphrodi-  
tis opulatis Wild. in act. nov. act. Soc. amic. hist. nat. Berol. 3. p.  
437.

Margynicarpus laevis; folioli apice lobatis, floribus divisis corollatis  
Wild. in nov. act. Soc. am. hist. nat. Berol. 3. 437.

Habitat in Brasilia, Pam et collibus aridis Chili

caulis suffruticosus, ramosissimus; rami diffusi, teretes: ramuli  
frequentissimi uti rami superne basi petiolorum dilatato mem-  
branaceo vaginali, <sup>facto</sup> foliis in ramulis, confertis, pedicellatis, pinnatis  
setis

