

Melastoma  
Wilsonii

x x x

herba Melast. foliis denticulatis quinque nervis ovato-lanceolatis caule hispido L. S. V. p. 404

Artuscula jamaicensis quinquenervi, minutissime denticulatis foliis et caule pubescente  
Tiburi Plum. alm. 265. fig. 1.

Grosfulana fructu non spinosa, malabathrifolius longa et rufa lanugine hispida,  
fructu majore Conleto Sloan hist. 2. p. 85. t. 197. fig. 2. Raii dendr. 74.

Grosfulana plantaginifolia angustiore hispida Plum. spec. 141.

Habitat in America meridionali t.

Flores in foliorum alis breviter pedunculati. Linn.

herba? herb. Surinam

Rami teretibus, glaberrimis. Folia petiolatis, bipollicariis, oblonga ovata, utraque  
acuta, denticulata, 5que nervis, marginibusque subtus celestibus pilis  
superius glabra nitida; nervi inter medios in petiolo desinentes, laterales parum  
superius basi. Synonym. Ploandi adjectum erat a Turc. arab. Diversa videtur

herba herb. L. Martii

Rami teretes, et punctis numerosis exasperati, superius imbricati, uti petioli. Margine  
foliorum, pedunculis et calycibus et pilis rigidulis hispidi. Folia petiolata, bipollicaria,  
ovato-lanceolata, angusta, triplicinervi, proster nervos marginales, attenuatos, serratos  
brunneis pediculis, supra tuberculato scabra, nervis punctis fetiferis, pectus varians to  
mentosa, in adultis fetis sanonibus, fetis purpureis centis. Pedunculi ex aqua cum ra



trulo, terni, dichotomi, incani, folio breviori. Floerpedicellati. Pedunculis parvula-  
peltis aei. Calycis lacinae sive lineares.

Diversa specimina ab herba in Collectione Lunani

*Setofa*  
*municata* Melastomo foliis ovato lanceolatis integerrimis, ramulis <sup>lucis</sup> ~~municatis~~ punctatis <sup>municatis</sup> ~~municatis~~  
pediculis <sup>pediculis</sup> ~~apice~~ <sup>calyx</sup> ~~apice~~ <sup>bracteisque</sup> ~~setosis~~ <sup>setosis</sup>

Habitata

Rami obscure tetragoni cortice cinereo, superne ad apicem punctis elevatis in  
quibus videntur setae rigidae sparse, articulati, <sup>articulati</sup> ~~apice~~ <sup>apice</sup> ~~setis~~ <sup>setis</sup> lon-  
gioribus rigidioribus. Folia opposita, triveria bipollicaria, nitida  
strigosa tamen non scabra subtus pallidiora nuda nervi tantum  
strigosi. Petiolus supra planus margine setis longis inaequalibus  
ciliatus. ~~Flora~~ <sup>racemi</sup> ~~racemi~~ <sup>racemi</sup> terminales pauci flori. Pedicelli breves  
uni- sive biflori basi barbati. Calyx subglobosus magnitudine  
nucis compositae. Petalis longis patulis strigosis laciniis lanceolatis  
Hamula decem

accidit ad Grosulariam plantaginis folio angustiore hispido Plum  
Sc. 141. spec. 18.

et Mel. hista differt floribus terminalibus



Kolo senčca Mel.

Oleandra, Trinesis

Melastoma triflorum, foliis elliptico-lanceolatis integerrimis, ramis petiolis calycibusque  
strigosis, floribus axillaribus subpedunculatis ternis.

Habitat in Cambi Forsyth 4.

Rami obtuse tetragoni, superne, uti petioli et calyces terti, setis frequentibus ad  
pressis.

Folia petiolata, opposita, tripolliciana vel ultra, utrinque attenuata, acuta, tri-  
nervi, proter nervi marginales, supra et infra in nervi et venis, setis  
adpressa setis adpressis flabescentibus ramorum parum angustioribus, vix ta-  
men scabra, subtus pallidiora ~~et~~ vicia.

Petioli unguiculares, tertiusculi.

Florae axillares, plerumque tres in singula axilla, brevissime pedunculati.

Calyx turbinatus, obscurè quadridentati.

Petalae quatuor, ovatae.

Stamina octo, corolla breviora. Antherae breves, lineari-oblongae, obtusae, purpureae,

dorso fusco-purpurascens; <sup>staminibus</sup>

(Differt a Meli fasciculata ut vides ex specimine ab amico optato, muso: foliis <sup>catombus</sup> ~~longioribus~~;  
super scabra, subtus pilis mollibus, floribus infra foliaceis longius pedunculatis pleribus  
calycibus minoribus glabris

Mel.







Levigata Mel. <sup>ovato lanceolatis acuminatis</sup> foliis integerrimis quinque nervis ~~ovato oblongis~~ <sup>dentatis</sup> ~~marginibus~~ <sup>marginibus</sup> ~~lin. S. V. p. 404~~

~~Grosularia fructu arbor non spinosa, malabathri folio maximo glabra et foliorum  
dentata Sloan. hist. p. 95. t. 197. fig. 1. Raci dendr. 26.~~

~~Grosularia plantagineis folio lato, fructu minimo concolor Plum. spec. 100. t. 146~~

Melastoma fructuosa minor, foliis veniibus ovatis, racemis terminalibus Brown  
fam. 219.

Habitat in America  
Folia tenuiora et minus confertia. Racemi terminaliter compressi. Flores  
minuti hinc.

Melastoma levigata foliis integerrimis quinque nervis ovato oblongis  
~~acutius~~ <sup>dentatis</sup> ~~marginibus~~ <sup>marginibus</sup> ~~lin. S. V. p. 404~~ <sup>foliis</sup>  
herb. ven. Aubl. fl. Gu. 1. p. 412. t. 159. <sup>per se non purascenti</sup> 6

~~rami~~ <sup>rami</sup> ~~obfuse~~ <sup>obfuse</sup> ~~tetragoni~~ <sup>tetragoni</sup> ~~glabra~~ <sup>glabra</sup> ~~folia~~ <sup>folia</sup> ~~petiolata~~ <sup>petiolata</sup> ~~tri~~ <sup>tri</sup> ~~vel~~ <sup>vel</sup> ~~polliniana~~ <sup>polliniana</sup>  
~~remoto raro loco~~ <sup>remoto raro loco</sup> ~~obfusa~~ <sup>obfusa</sup> ~~dentulata~~ <sup>dentulata</sup> ~~tri~~ <sup>tri</sup> ~~nervia~~ <sup>nervia</sup> ~~proter~~ <sup>proter</sup> ~~nervos~~ <sup>nervos</sup> ~~marginales~~ <sup>marginales</sup> ~~nervosa~~ <sup>nervosa</sup> ~~avenosa~~ <sup>avenosa</sup>  
~~dentium~~ <sup>dentium</sup> ~~reticulata~~ <sup>reticulata</sup> ~~subrugosa~~ <sup>subrugosa</sup> ~~ut~~ <sup>ut</sup> ~~naque~~ <sup>naque</sup> ~~glabra~~ <sup>glabra</sup> ~~Racemi~~ <sup>Racemi</sup> ~~terminaliter~~ <sup>terminaliter</sup> ~~dendr.~~ <sup>dendr.</sup>  
~~nervi~~ <sup>nervi</sup> ~~subtus~~ <sup>subtus</sup> ~~obfusa~~ <sup>obfusa</sup> ~~propti~~ <sup>propti</sup> ~~pulverulentis~~ <sup>pulverulentis</sup> ~~Pedunculi~~ <sup>Pedunculi</sup> ~~patrici~~ <sup>patrici</sup> ~~remoti~~ <sup>remoti</sup> ~~si~~ <sup>si</sup> ~~si~~ <sup>si</sup> ~~for~~ <sup>for</sup> ~~mes~~ <sup>mes</sup> ~~unguicula~~ <sup>unguicula</sup>  
~~pulverulentis~~ <sup>pulverulentis</sup> ~~coctum~~ <sup>coctum</sup> ~~res~~ <sup>res</sup> ~~calycis~~ <sup>calycis</sup> ~~minuti~~ <sup>minuti</sup> ~~dentati~~ <sup>dentati</sup> ~~farinosi~~ <sup>farinosi</sup> ~~fructus~~ <sup>fructus</sup> ~~glaber~~ <sup>glaber</sup> ~~magnitudine~~ <sup>magnitudine</sup> ~~virgini~~ <sup>virgini</sup> ~~excitus~~ <sup>excitus</sup>  
~~subtus pallidum~~ <sup>subtus pallidum</sup> ~~his~~ <sup>his</sup> ~~cordandi~~ <sup>cordandi</sup>

Pedunculi tres in apice ramorum vix semper, intermedii longius tetragonis  
incluliferi laterales patentibus compressi omnes pulvere ferrugines 25.



rami divaricati <sup>inter</sup> collibus <sup>paucis</sup> flexu  
nodales oppositi remoti biparti. Flores sessiles in ramis pedunculorum  
nodalium simpliciter prius fundi cum flore sessili in divaricatione  
parvi, distantes. Bractea ~~parva~~ <sup>parva</sup> duo ad basin pedunculorum  
nodalium pedunculorum et florum. Petala linearia oblonga

remorum melastoma <sup>integerrimis & inerviis</sup> foliis lanceolatis, floribus terminalibus subternis, calycibus <sup>mollibus</sup> pelosis. ramis <sup>capitis</sup>

Habitat in India orientali ms

Theriacali Rhæd. Mal. 4. Tab. 44

Habitat in India t<sup>p</sup>

Rami obscure tetragoni scaberrimi ferruginei punctis elevatis frigidis. Folia opposita  
pictolata, lanceolata, trinerviâ, scabra pubescentia frigidis variis, subtus in nervis magis  
conspicuis, ozulo armato sive feno oculo armato, pollicaria subtus pallidiora. Flores  
in apice ramorum, quatuor vel terni oppositi, brevis pedunculati foliolo lanceolato  
utrinque ad basin pedicellorum. Calyx campanulatus cæcis villosus inæqualis mollis  
intus glaber laciniis lanceolatis filamenta <sup>duorum</sup> petala unguiculata  
lamina rotundata  
scabi scabitur e lycium operâ offert.



*Octandora melastomo* foliis integerrimis ovatis glabris margine hispida  
Lia fl. Zeyl. 173 \* Jff. et. 213. L. V. p. 404.  
Habitat in India China





*Calyx* campanulatus, villosus tomentosus ferrugineo ~~laevigato~~. qui quiescit in Calicis  
laciniolatis dorso villosus tomentosus. Calicibus glabris margine ciliatis  
Corolla major quam in reliquis Petalis obovatis ~~subrotundis~~. <sup>subrotundis</sup> plicis longis  
subtilibus filamenta 10 basi et apice glaberrima medio hirsutis fimo, Anthera  
longitudine filamentorum subulata cum incurva ~~f~~ medio plicis  
stylus longitudine filamentorum ~~stigma~~ subacutum. Germen ovatum  
decem plicatum

Capillaris Mel. foliis subdentulatis ovato lanceolatis attenuatis glabris; pedunculis  
capillaribus trifloris axillaribus Schw. prodr. p. 71.

Habitat in Jamaica 5

Tolia trinervis. Flores octandri. Pedunculi oppositi foliis brevioribus.



*Antella* Mel. folus margine hirsuto pilosis, pedunculis brevissimis axillaribus trifloris

Schw. prod. p. 72.

Habitat in Jamaica

Folia trinervia. Flores octandri. Folia ovato-lanceolata, sub serrulato  
ciliatis acuta trinervia: Nervis subtus hirsutis supra glabra. Pedun-  
culis axillaribus subtrifloris ramis hirsutis

microphylla Mel folio subdenticulatis ovatis obtusis superne hispida in fra tomentosis  
pedunculis trifidis axillaribus Thw. prodr. p. 72.  
- Habitat in India occidentali.  
Folia trinervis. Flores octandri

non ed.



*micranthus* Mel. foliis Denticulatis oblongis acutis glabris; racemis axillaribus  
reclinatis, floribus acuminatis Schw. prodr. p. 71.

Habitat in Jamaica h.

Folia triplex <sup>herbaria</sup> seores octandri. Racemis decompositis

purpurascens Mel. foliis integris oblongis acutis striatis glaberrimis; racemis laterali-  
bus patulis floribus distinctis Schw. prodr p. 71.

Habitat in Jamaica &

Folia trinervia. Flores octandri  
perenne



mbens Mel. foliis denticulatis ovato lanceolatis acuminatis glaberrimis ramis  
petiolisque coloratis; racemo terminali floribus confertis divicis Sitw. prodr.

p. 71.

Melastoma P. Brown. Jam. 219.

Habitat in Jamaica &

Solia trinervis. Flores octandri

alpina Mel. foliis integris ovatis coriaceis glabris; pedunculis simpliciter trichotomis,  
floribus distinctis Schw. prodr. p. 71.  
Habitat in Italia occidentali &c.  
Tolia trinevia. Flores octandri



Saxifragae Mel. foliis integris ovatis acutis scabris ramis floriferis, floribus brevi-  
ter pedunculatis confertis Schw. prodr. p. 71.  
Habitat in Jamaica  
Solia trinervis. J. Ceres octandri

Melastoma lateriflora (~~Catalpa~~) foliis obovatis ~~triplicinervis~~ acuminatis setaceo subser-  
tis glabris, pedunculis infra foliaciis aggregatis unifloris.

Habitat versus summitates montium in fovea montiserrat Ryan 5 8 2 1  
Caulis bi-sive tripedali. Rami inferne teretes, superne tetragoni, glabri, dicto  
toni, purpureascentes.

Folia superne in rami, petiolata, opposita, bi-sive tripolliciana, oblonga, basi  
attenuata, apice acuminata, <sup>sub</sup>conica, margine parum reflexa, serrata  
serraturis minutissimis, serratis, apice fere brevi sermonati, margine folia  
d<sup>+</sup>pressis, utroque glabra, subtus pallida, scabruscula.

# triplicinervis

Petiolus pollicaris, supra subcanaliculatus, subtus convexus.

Pedunculi infra folia, aggregati, sex vel septem, breves, uniflori. Corei parvi.

Calyx truncatus, oculo armato subtrigosus.

Petala quatuor, parva, ovata, attenuata.

Stamina octo, longitudine petalorum. Adhaerentia rufa, exterior margine pallidiori, oblonga  
obtusata, longitudine fereamentorum 9

Stylus Staminibus longior. Stigma obtusum subcapitatum

Bacca magnitudine seminis Coriandri, globosa, oculo armato prehis adpressa rationibus  
strogosis adpressis.



Fernynea  
Ventricillata

~~aloso - subtomentosa, sepibus dupl. lobis, fl. <sup>supernis</sup> fasciatis, sepibus  
Melastoma, lutea, foliis lanceolato-ovatis denticulatis, floribus axil-  
lariis, sepibus subventricillatis~~

~~Habitat in India orientali. Dr. von Rohr.~~

~~Primi obtuse tetragoni, <sup>subferre</sup> <sup>refescentibus</sup> <sup>junioris</sup> <sup>curvis</sup> <sup>apice</sup>  
pallidioribus, <sup>griseo</sup> <sup>refescentes</sup>~~

~~Folia <sup>subopposita</sup> <sup>tripollicaria</sup>, <sup>utroque</sup> <sup>obtusiuscula</sup>,  
subtus <sup>sub</sup> <sup>tripollicaria</sup> <sup>utroque</sup> <sup>obtusiuscula</sup>, <sup>quaque</sup> <sup>nervia</sup> <sup>supra</sup> <sup>fasciatis</sup>, <sup>subtus</sup>  
villosa, <sup>sub</sup> <sup>ferre</sup> <sup>griseo</sup> <sup>flavescentes</sup>  
tomentosa, <sup>molli</sup>~~

seri quinque proter  
marginales

seri  
villi flavescentibus  
curvis

~~Floris axillaris, <sup>sub</sup> <sup>ferre</sup> <sup>griseo</sup> <sup>flavescentes</sup>  
triquetris utrinque, <sup>junior</sup> <sup>curvis</sup>~~

Petala quatuor oblonga.

Stamena 8. Longitudine corollae

<sup>arant</sup>  
glabrata Mel. foliis <sup>minutis</sup> subdentatis ovatis acutis glabris coriaceis; pedunculis  
terminalibus solitariis unifloris Schw. prodr. pag. 71.

Habitat in Jamaica

Folia trinervia. Flores octandri. Folia trinervia ovata acumi-  
nata ad insertionem nervorum lateralium utroque fasciculus.  
pedunculi plerumque solitarii interdum terni



*Hirsuta* Melastoma foliis lato-lanceolatis subferrugatis; supra subpuberis subtus hirsutis  
ram Pedunculis axillaribus trifloris hirsutis.

Melastoma *hirsuta* foliis trinerviis subdenticulatis lato-lanceolatis  
attenuatis hirsutis; pedunculis axillaribus trifloris divaricatis  
Habitat in Jamaica.

~~Paulo~~ rami petioli pedunculi et calyces hirsuti. Folia petiolate  
obtusius serrata, trinerviis. Pedunculi axillares oppositi triflori  
Flores intermedius sessili. Lacinae calycis subulatae

hirsuta Mel. foliis ovata-lanceolatis <sup>attenuatis subperfoliatis</sup> ~~acuminatis denticulatis~~ supra hirsutis subtus  
hirsutis  
melastomo hirsuta foliis trinerviis subdenticulatis lato lanceolatis atten-  
nuatis ~~hirsutis~~, pedunculis axillaribus & floribus divaricatis  
Sw. prod. pag 71.



Melastoma verticillata, foliis ovatis oblongis attenuatis denticulatis villosis supra scabris, vix  
omni axillaribus; floribus verticillatis.

Habitat in Caribis. Forst. t. 5.

Rami molliter denseque villosi, tetragoni, articulati: articuli tripollicares; sursum sen-  
sim breviores, lateribus oppositis alternatim. e margine membranaceo angusto canaliculatis

folia petiolata, opposita, quadri- vel quinque pollicaria, in axillari angusto attenuata  
triplex nervia proter nervi marginales parum supra basin orti, subtilissime denticulata,  
ciliata, utrinque pilosa villosa, praesertim subtus in costa et nervis: supra e punctis  
elevatis scabra; subtus minus punctata, minus scabra, pallidiora.

Petoli dense villosi, unguiculati, verticillati

Pedunculi ex axillis alternis, solitarii, bipollicares. Flores verticillati, ~~ferax~~ <sup>ferax</sup> in  
femore pedicellati, superiores sessiles. Pedicelli uniflori, interdum biflori,

#femina in singulis pedicellis longioris, unguiculati, inaequales.  
Co verticillati,

Calyx quadripidus, villosus: laciniis linearis-lanceolatis, brevibus, reflexis.

Petala quatuor, lanceolata, parva

Stamina octo, petalis parum longiora.

Bacca globosa, semina Sinapis triplo major.





vergata Melast. foliis integris ovato-lanceolatis acuminatis glaberrimis; ramis  
floriferis; racemis decussatis diffusis parvis Schw. prodr. 72.

Habitat in Jamaica.

Folia triplinervia. Flores octandri.

Oilandia, triplinervis

Melastoma acuminata, foliis ovatis extrorsum denticulatis acuminatis subtus incanis, corymbis terminalibus fastigiatis.

Habitat in Monserrat Ryan 5

Rami inferne teretiusculi, apicem versus tetragoni; pulverulento incanis, subfoliis dilatati.

Stem petiolata, opposita, tri- vel quinque pollicaria, saepe acuminata, corymbis <sup>paucis</sup> proxima lanceolata, inferne integerrima, extrorsum versus denticulata; supra glaberrima, laevia, triplinervis, praeter nervi marginales a basi orti; subtus ~~teretiusculi~~ incanis, venosa, venis sinuatis; ramotis, nervis adnatis, comae flavescens. juniora pulverulenta.

Secundum  
Richard  
et discolor  
Jacquin  
saepe subtus

Petiolis uti pedunculis pulverulentis, supra planis, subtus carinatis. Corymbis terminalibus, trichotomis, fastigiatis. Pedunculi universales ternis, comae uti pedunculis compressi articulati; articulis apice latioribus.

Calyx vixidi flavescens, subpulverulentus, basi globosus, superne coarctatus; limbus subexplanatus, obtuse quadrilobus.

Petalis quatuor, oblonga, flava.

Stamina octo, corolla parum longiora. antheris linearibus, dorso flava, margine albis.



Dodecandra melastoma foliis ellipticis lanciolatis glabri obsolete denticulatis,  
floribus paniculatis dodecandris

Habitat in montserrat du Ryan &

Rami <sup>inermes</sup> ~~trigonis~~, glabri

Folia petiolata, quoadi vel quinque pollicaria, patentissima, <sup>stemmat</sup> acuminata  
utrinque acuta, <sup>et basi</sup> ~~maxime~~ <sup>et utrinque glabris</sup> ~~dentata~~, subtus pallidiora: ternaria proder  
nervi marginales <sup>pubes</sup>

Panicula terminalis, erecta, rami patentes, in summo quatuor, reli  
qui oppositi, glabri trichotomi: ramulis divaricatis trifloris. Flores

ut - pedunculis communis  
articulatis basi in  
eius apice dilatatis

Calyx magnitudine feminis coriandri trineatus, leviter glaberrimus,  
lobis pallide viridis grossis viridibus: calyx

Petala oblonga lutea 5que

Stamina 12-13. Longitudine corollae anthera linearis utraque obtusa  
integra filamenta

Melastoma sessiliflora <sup>(Lam.)</sup> villosa-subtomentosa, foliis lanceolato-ovatis ~~quaque nervis~~  
denticulatis, <sup>subreticulatis</sup> floribus axillaribus sessilibus subverticillatis.

Habitat in America meridionali. var. Prostr. 5.

Rami obtuse tetragoni, et pilis densis crispis femineis tomentosi, apice griseo-~~res~~  
nufescentes.

Folia brevissime petiolata, opposita, bi-<sup>tr</sup>-five tripollicaria, utrinque obtusiuscula  
subtilissime denticulata; supra rugosula, <sup>fiatruvscula</sup> subtus villosiora, subtomentosa,  
mollia, griseo-flavescentia, nervi quinque proter marginales, <sup>et quatuor</sup>  
~~et utrinque~~ griseo-tomentosa.

Flores axillares, utrinque tres vel quatuor, sessiles, parvi

Calycis laciniis parvis quatuor, ovatis, parvis, exclusis villosis.

Petalis quatuor, oblongis.

Stamina octo, longitudine corollae. Anthera lineari-oblonga, purpurea, longitudi-  
<sup>ne tere filamentorum</sup>

Basis globosa, magnitudine seminis Conandi.



pilosa Melastoma foliis denticulatis oblongis subtus hispidis; racemis lateralibus his-  
pidis Schw. prodr. 72.

Habitat in Jamaica.

Bolia quinque nervia Flores octandri

*Stemodia*

Octandris quinque nervia

Melastoma coccinea foliis elliptico-oratis acuminatis glabris integerrimis,  
thyso terminali, ramis pedunculisque nodosis hispidis.

Melastoma coccinea: nodosa tener: foliis ovatis, angusto-acuminatis, integris, lucidis,  
quinque nervis: ramis subsessilibus ter trifloris, coccineis Act. Soc. hist. nat. Paris.

p. l. pag. 109. 2

Habitat in Montserrat. Ryan 5

Rami obscure tetragonis, <sup>ter trifloris</sup> inferne glabri, superne, uti petioli et pedunculi a pilis  
gracilibus sparsis crispis subhipidi.

Litt. N.

p. 49

Folia petiolata, opposita, tripollucaria, elliptico-orata, acuminata, integerrima  
utrinque glabra, lora, quinque nervia: supra secundum venas transversas  
lineata, soepe adpersa punctis minutis albis, prosertim juncea, subtus palli-  
de undici, venoso-reticulata, nervi venisque lutescentibus. Juniori supra  
secundum venas transversas lineata.

Petioli ciliati, basi versus glabriores et crassiores.

4 juniores

epilis brevibus  
sordidigracilibus

Thyso terminali, bracteato. Pedunculi palmari. Pedunculi partiales op-  
positi, uti pedunculi com munes ad ramificationes eras nodosi, <sup>4</sup> semel bis  
vel ternis ter trichotomi, ramo ultimo trifloro. Flores pedicellati: pedicelli  
brevi. Folium lanceolatum <sup>utrinque</sup> sub infernis ramificationibus thysi. Bractea

Petalea.

Calyx subquadrilobus, albedo-punctatus, glaber: lobi rotundati.

Petala quatuor, oblonga, glabra, unguiculata



Stamina octo.

Stylus corolla longior

Ex observata Ryani sequentia addere lubet.

Frutex mihi omnium facile pulcherrimus, quadri vel sexpedalis, summitate saepe de exaite comosa, ut Citrus aurantium viridanorum. Flores magni calicinei, octandri, in Thyro congesti, totam superficiem como vegetas.

Folia rugosa. Variat interdum flore albo.

Hic semel observandum de Melastomati, Psychotrii alisque montium indigenis, nihil certi pro charactera specifico e glabritate vel hirsutitate foliorum erudendum esse, cum perpetuo variant ~~in hirsutitate~~ <sup>vel</sup> hirsutitate viride vel in valle queto et <sup>creverunt</sup> securis <sup>in solo</sup> ~~in solo~~ <sup>in solo</sup> abrupto, unde, tempestatibusque exposito, ~~appropinquat~~ <sup>appropinquat</sup> ventorum impetum hisce regionibus familiarem exposita sunt.

nodosa lae manthi si non eadem certe tamen quam matruae affinis lex. <sup>spec. herb. Ryani</sup> tetragonis, articulati nodosi, circoscissati. Folia oblongo-ovata, utrinque glabra, nervosa, postea nervi laterales, trans versus nervosa, supra subnervosa. Pediculi trigoni. Calyces quadrilobi

discolor Mel. foliis integerrimis quinque nerviis oblongo ovatis glabris margine lobatis Linn.  
L. V. p. 404.

Melastoma floribus rotundis, foliis quinque nerviis Jacq. Amer. 130. t. 84

Arbor americana quinque nervia, comantibus flosculis, foliis amplissimis  
glabris prona parte albicantibus Plum alm. 40. t. 264. fig. 4.

Grossularia fructu non spinosa, malabathri foliis subtus niveis, fructu  
racemoso umbello modo dispositis Sloan h. 2. p. 86. t. 190. fig. 1.

Grossularia plantaginifolia apicibus candidis et falcatis Plum t. 1.  
142. fig. 1.

Habitat in America

flores in Plum. magni in Jacq. quini parvae in ceteris eadem

Specimina herbarii huius: quam maxime accedunt ad meam acuminatam  
folia cordato ovata, attenuata, 5que nervia, supra glaberrima, subtus  
canescentia, spithamea, remote denticulata. Panicula patens, uti ramis  
superioribus pubescentibus, ramis inferioribus. Capitulo vixescentibus, ad ridentem  
tate, magnitudine seminis Coriandri. forte quod huius uti acuminata  
mea eadem cum Jacq. et diverso a Plum. eam



umbrosa Mel. foliis denticulatis lato-ovatis acuminatis utrisque latis; ramulis floriferis  
fere dichotomis patentibus; ramis pediculisque hirsutis. Schw. prodr. p. 72!

Habitat in fl. Christophert

Folia sive nervia flores octandri

hispida Mel foliis subdentulatis ovatis acutis superne frigidis hirsutis subtus  
reticulato tomentosis, paniculis terminalibus divaricatis Schw. prodr. p. 72.

Habitat in Jamaica

Folia quinque nervia. Flores octandri



Corymbosa Mel. glaberrima, foliis subcordato-ovatis, in acumen angustum  
promise angustatis, cæcis nervis corymbis plentibus Verminalibus pe  
Denticulatis subcymosis Act. f. h. n. Par. 1. pag. 109  
Habitat in Amer. merid.

+ *Strandra mel.* foliis integris oblongis acuminatis basi emarginatis glabris convexis in  
filiis; racemo erecto terminali, floribus tetrandris Schw. prodr. p. 72.

Habitat in Jamaica 5

Folia trinervia. Flores ~~standi~~ ~~tetra~~ Racemo supra de composito  
bracteata. Flores parvi





*grosfa melastoma*



stellate Mel. (floribus tetrandris) foliis ovatis acutis <sup>pulverulentis</sup> subtus stellatis villosis, racemis  
terminalibus  
decemnerviis pedicellis trifloris

Habitat in India orientali <sup>supra</sup> Pottley, Setrago.

Calculi fruticosis rami teretes <sup>supra</sup> pulverulento tomentosi  
folia opposita petiolata ~~et~~ bipollicaria integerrima  
acuta uninervia utraque nervis transverso distantibus  
simplicibus supra glabra subtus pulverulento villosis  
velis subtellatis petioli breviter villosi uti racemus  
pedunculi et calycis. pedicellis 3. unifloris. bracteis 2 lon-  
geolatis basin pedicellorum et solitaria ad propeum  
florem longitudine calycis. Petala 4 lanceolata  
calyx marginis integerrimus Stamina 400 calycis  
longiora. filamenta omnia villosa stylus staminibus  
longior pedunculi et pedicelli oppositi.