

Melia

1. azadirach M. foliis bipinnatis, foliolis ovatis
2. semper virens M. foliis bipinnatis, foliolis rugosis, ovatis subseptenis
3. aradirachta M. foliis pinnatis, foliolis falcatis
4. pulverulenta M. foliis compositis decempositisque, foliolis ovatis subente
germinis, paniculis pubescentibus

1 *Ardisaech Melia* ^{folioli squarrolato ovatis}
foliis bipinnatis Lin. l. V. pag. 400

Pseudo-Sycomorus Cam. epit. 101. Tab.

Ardisaech Dod. pempt. 840.

Arbor fraxini folio, flore caeruleo Bauh. Pin. 495

~~B. *Ardisaech* sempervirens et semper florens Tournes. Fl. R. 4. 616~~

~~*Ardisaechta indica*, foliis ramosis minoribus, flore albo subcaeruleo
purpurascente majore Com. hort. l. p. 147. t. 78. Flor. Zeig. 162.~~

~~Habitat in Syria. β in Ceylona γ in Anpula δ in Cruis Pflanz. 5~~

~~Arboreus. Caele glaber Rami glaberrimi. Folia alterna bipinnata, pedita
Cui pennis quadrijugis suboppositis. Panulis oppositis alternis que
peticolatis tri-vel 4jugis, ovatis acuminatis, pollicantibus, grosse
ferratis caeternis subconfluentibus. Panicula axillaris~~

~~d α vari et altem foliis pennatis pennis tripartitis vel ternatis incisifs
V. foliis simpliciter pennatis.~~

pulvinata Melia

sempervirens M. foliis bipinnatis, foliolis oppositis pinnatis subseptenis Sw.
prodr. p. 67.

Azadirachta sempervirens et semperflorens Tournef. 4. R. 4. 616.
Azadirachta indica, foliis ramis minoribus, flore albo
subimbrato purpurascente majore Bern. hort. 1. p. 147. & 76. haeft
2. p. 167
et habitat in Zeylone

an varietas V azadirach

non vide

7 ^{pulverulenta}
~~Corymbosa~~ Melia foliis compositis decompositis que, foliolis ovatis subinteyeminis
 paniculis ^{pubescentibus}
 Habitat in ^{India Orientali, tunc}
 Arbor ramis ^{griseo} fusis purpurascentibus luteis, cortice rufo, tuberculis
 non parvis ferrugineis adpressis, cicatricibus ^{trilobis} trilobis
 foliis ad apicem ramorum congesta petiolata, pedalis, pennata ^{3-4. parvum} vel
 pennis ternatis. ^{extremum} Penna opposita petiolata ovata, ^{simpliciter} elliptica
 acuminata, ^{es} inteyemina vel obscure crenata, remotissima, pollicari
 non glabra, ^{et adnatis ad angulum} nervosa pulvere lanigosa. Pediculi flavescentes
 Pedunculi ^{et apice} axillares, foliis brevioribus fastigiati.
 Pedunculi pedales oppositi: uti pedicelli, pulverulenti fasti-
 giati in Corymbum hemisphaericum diametro bipollicari.
 Flores minores quam in antecedentibus Calyx et corolla pubescentes
 canescentes. Arbus neitani glaber

#2 ^{capa} petiolorum
 et pedunculorum

Catus folii

2 *Aradurachta Melia* foliis pennatis foliolis ^{falciatis} ~~in~~ ~~equilatis~~ ~~in~~ ~~his~~
Astor India, fraxino similis oleo fructu Bauh. Pin 416
Ani Bepou Rheed. Malab 4. p. 107. t. 52.

Olea Malabarica, fraxini, folio Pluk. alm. 26y. t. 247. fig. 1.
Aradurachta indica, folio fraxini Breyn. Ic. 2 ~~45~~ 15. p. 15. t. 1.

Aradurachfolius falciato serratis Burm. zeyl. 40 t. 15

~~Habitata in India~~

Melia foliis pennatis Lin. Fl. Zeyl. 161. * S. V. p. 400

Habitata in India ~~Jace~~

Tota glabra. Folio superne in ramo ~~congesta~~ ^{ola}, plus quam pedalia
pennata cum impari 5-9 juga. ^{inter nervi, or. profeta} Folia alterna ^{longior} brevis pome
petiolata remota, bipollicaria extrosi fensu ^{longior} minor lan
scolata, acuminata a tero latere angustiora breviora
altero fensu ^{longior} basi obliqua a basi fere usque ad apicem
serrata dentata ^{addimus} Pannculi axillares semipedalis minus
petiolatus ^{si ens} ^{supreme} ^{planus} ^{sub margini} ^{natus} ^{superne} ^{ramo} ^{quam in} ^{antecedente} ^{pedunculi} ^{partiale} ^{retro} ^{si}
mifiliformis.

+ a medio dentato
serrata

petiolatus ^{si ens} ^{supreme} ^{planus} ^{sub margini} ^{natus} ^{superne}

Melia: foliis pinnatis, foliolis ovatis dentato-incisi

ⁱⁿ
~~Cum Sapula fl. Cnicis~~

Habitat in Sapula fl. Cnicis 4.

Forte varietas Aredamit. differt foliis simpliciter. ab Aradiracta foliolis
ovatis.

Dubra M. foliis bipinnatis: pennulis acutis serratis; Terminali majore lacin

Def. bot. 7. p. 364

Habitat

non vidi