

Melicocca

♀ Calyx 4-partitus. Petala 4. tandem reflexa. Stigma subretractum 3 angu-
lari. Drupa corticosa

♂ Cal. Corol. hermaphroditi. Stamina 8. basi contigua.

1. *bijuga* M. foliis ^{sub} ~~uni~~ ^{bijugis}, petiolis alatis, racemus ^{compositus} ~~glabrus~~ ^{granulatus}
2. *nitida* M. foliis uni ^{bijugis}, petiolis nudis, floribus ^{tomentosis} ~~paniculatis~~

2 nitida

Melicocca folius uni f. bijugi petiolis nudis

Habitata India occidentali

Rami confertissime cinereo punctati. ^{pollicari in f. uni f. bijugi in unigugi et in paniculatis} ~~folia~~ ^{folia} petiolata alterna

Petiolus ^{subro nigra} purpurascens. ^{genuissime tenota} foliola lanceolata elliptica subsessilia

late vnda, venosa obtusiuscula submarginata, glaberrima

^{folia unigugi} ~~folia unigugi~~, bi f. tripolliaria. Panicula terminalis

longitudine foliorum ~~bracteata~~, calycis pubescentis
pedunculi parciales
et pedicelli alterni

differt a bijuga petiolo ^{et folioli} brevioribus foliolis minoribus obtusioribus plerumque emarginatis, paniculis minoribus calycibus tomentosis incanis
Rami confertissime cinereo punctati

bijuga Melicocca Lin. S. V. pag. 362.

Habitat in India occid. In v. Rohr et insula St. Anicis & West I.

Arbor. Rami teretes, purpureo-fentes, glabri: juniores pallidi, angulati.
Folia pedicelata, alterna, bijuga. Foliola breviter pedicelata, opposita, tri-
quadripollicaria, intima minora, elliptica, obtusiuscula, integerrima
utrinque minus glabra, venosa, tenera.

Pedunculi supra planiusculi.

Racemi terminalis, compositi, laxi, patentissimi. Pedunculi pariter
inferne, alterni, tri vel quadripollicares, filiformes, approxima-
ti: pedicelli copiosi, capillares, alterni, distantes, semper quicun-
que, uniflori

Flores polygami, dioici

(an mea species a Jacq. et Lin. diversa, vel forte quod hermaphroditum
tantum vidit Jacq.)

bijuga Melicocota neivolis alatis.
Nux americana *foliis alatis bifidis* Com. hort. 1. p. 107. t. 94.
Nux americana *costa foliorum appendicibus acuta* Pluk alm. 265.
t. 207. fig. 4.

Melicocotus foliis ut plurimum *bijugatis ovatis, per pennas alatis*
Dispositus Brown. jam. 210.

Habitat in America meridionali *racemulose angulati* 6: f. 3. *juva*
Rami *costae cinereo conspice digit. minoris.* *Solia alterna petiolata*
abrupte pennata, petiolus marginatus. Foliole subpetiolata

unicolata- elliptica, ceteriorima, obtusifolia 6: f. 3. *polliam*
vel ulera angulata ceteriorima *ultra duo* *ultra duo*
interiora minor, opposita interdum unicum *racemulose* *racemi compositi*
Pedunculus et pedicelli *pubescentes. Flores paniculati*
mianti glabri. *Drupe magnitudine Cerasi* *Rami paniculati*
alterna

netiosi et pedunculi
st. rati

Solia ramorum fructiferorum *vix alata bijuga, juniora alata*
H *et pedicelli quadrangulo, interdum alternis. sic diversa videtur si ramis*
alterni glabri prouti ramuli juniora vel adulescentia in herbano habentur

varia est secundum Richard