

# Mamecyclon

1. capitellatum M. foliis ovatis obtusiusculis, capitulis axillaribus pedunculatis
2. fragrans M. foliis ellipticis obtusiusculis, floribus subsymptosis
3. nudiflorum M. foliis ovatis acuminatis, floribus subsymptosis

Capitellatum Memecylon foliis ovatis obtusifoliis capitulis axillaribus pedunculatis  
Cornus sylvestris, foliis ovatis colorem argenteis, floiulis ad folionem alas  
globosis Burm. Zeyl. 76. t. 30.  
Memecylon capitellatum foliis ovatis Lin. Flor. Zeyl. 1364. V. p. 362.

Habitat in Zeylona;

Rami ~~teretes~~ <sup>reticulata opposita</sup> cinerascetes. Colea coracea avenia pollicaria po-  
glabra integerrima, ~~pediculi~~ <sup>pediculi</sup> oppositi solitarii breves capitulo florum  
terminati. Calyx campanulatus integer

Rami cortice griseo, glabri, subdilatome dupi, inferne teretes  
superne tetragon: utriculis apice panum crispantibus. Folia subsimplex  
oblonga, glabra, pubescentia obtusa, integerrima, trinervia, simplicia  
ter venosa. Pedunculi latentes, aggregati, breves

Coniacum Memecylon corymbis axillaribus foliis ellipticis coriaceis ovatis

Habitat in India orientali a Dale sub nomine myrti mesum  
Rami teretes, glabri. Folia petiolata bis. tripollicaria obtusa vel acuta  
integerrima. Corymbi parvi

an varietas Memec. fragrans

Royen Conium Lin. l. V. pag. 270

Selecatur ~~in~~ ~~systemate~~ ~~vegetabilium~~ cum eadem planta sit ac Caucalis  
daucoides

~~flarescens~~ <sup>elliptico</sup> laurus foliis ovatis, subtrinerviis oppositis, umbellae axillares brevissimis.

Habitat in India orientali Hongkong

Rami teretes glabri, & inermi apice purpurascens oppositi.  
Folia petiolo conica, infusca flavo-ovata glaberrima nervi-  
tibus a basi orti obscuri superne evanescenti, reliqui obliqui  
obscuri arenaria bipollicaria acuta. Petiolus brevissimus. Peduncul-  
culi oppositi plerumque interdum gemini petiolo triplo longiores <sup>parvi</sup> ~~can-~~  
sindi caespitibus. flori plures <sup>parvi</sup> pedicellati involucrio diphylo lanceolato  
to longitudine. florum proter squama minuta ad basin singuli floris

capitellatum mameylon

videtur potius species mameylo

nudiflorum  
 macrophyllum memecylon foliis ovatis acuminatis, floribus subtymosis  
 memecylon grande foliis acuminatis. Retz obs. 24 reg. 26  
 Habitat in sylvis zeylanicis hanc infra folia cyma opposita remota & cicatrix petioli dicitur  
 23 ut <sup>usque</sup> ~~ut~~ <sup>umbellae</sup> ~~umbellae~~ compositae radius universalibus 3-5: partiales plures, radius universalis  
 adducendi <sup>usque</sup> ~~usque~~ <sup>intermedii</sup> ~~intermedii~~ soepe denum trifidus pedicellis umbellatis. folia opposita,  
 vides panem <sup>intermedii</sup> ~~intermedii~~ petiolata, palmata, ovata acuminata, Coris semper coriacea nervosa  
 brevioris <sup>rami</sup> ~~rami~~

grande Retz

2 fragrans Memecylon foliis ellipticis oppositifoliis, floribus subsymptosis in profolis  
Cormus Zeylanica sylvestris altera Morakaha dicta Burm. Zeyl. p. 76  
Tab. 31.

Habita in India orientali

Similis precedenti Rar ad pedicelli distincti nec flores in capitulum  
contracti Rami floriferi aphylli nisi inferne aphylli floriferi  
Pediculi vix semiunguiculari apposti interdum solitarii saepe  
gemini et terni utraque pediculi partibus breves interdum vespis  
apicem per inferos verticillati et terminales umbellati vel etiam in  
umbellula composita folio basi acutissimo

Et Causha or Causha - chedda Malab. her. Pat. mus. herb. Vaillant

