

Mimosa
x foliis simplicibus

Simplifolia M. ^{cauliculis} foliis simplicibus, junioribus duplicato-pinnatisque
Memosa heterophylla foliis aliis simplicissimis foliatis rigidis aliis duplicato
pinnatis ~~caulis~~ ~~erectis~~
Memosa simplicifolia inermis, arborescens, foliis ovatis integerrimis nervosis
obtusis, spicis globosis pedunculatis Lin. Suppl. p. 436. 41. V. p. 913

Memosa simplex foliis simplicibus septem nerviis ovatis Sparm.
in Act. Ups. vol. 3. p. 195

Habitat in Ampula Tanna ^{amicorum, nova Hebrides, nova Caledonia} & ~~St. Paul. Bourbon~~ 5

Rami angulati per spurae scentes a foliis obovatis magnis glabris
integerrimis alternis absque costa longitudinali nervosa ner-
vis main nervosis, brevis rosine petiolata. Stipula nulla. Spica
globosa axillares pedunculata, parvo legumen compressum
incurvatum longitudine foliorum, torulofum

montanum Rumph. orb 3. t. 81.
Mangium Rumphii huius forte spectat

Lineari Mimosa foliis lineari bus, capitulis globosis axillaribus racemosis

^{inermis}
Ex horto Celsi Parisiensis et herb. Fusf.

Rami tenues, glabri, e linea a petiolo decurrente angulati. Folia parva
glabra, vix pollicaria, mucronata. Petiolus brevissimus. Racemi longi
vix inae florum. Capitula pedicellata, magis nitiditate seminis Conandri.
Flores flavi.

obliqua memosa in arboribus petiolis planis lineari lanceolatis oblongis
nervis foliiformibus, spicis axillaribus globosis La Mart
Tab. Gour n. D hist. natur. no 3. pag. 109. Tab. 5.

Habitu

Folia simplicia simplicia oblonga ovata, pollicaria marginata, glabra
Rami e linea decurrente angulati, superne compressi. Racemi
axillares, longitudine folii. Flores 3-4 sepius in apice pedicelli
Calyces glabri herb. Mus.

conf. ~~memosa suaveolens~~ Lin. Transact.

non vid.

Cyperus scopellatus umbellis ~~sub~~ ^{apice} aphyllis; umbellula radiis alternis ramosis com-
pactis fasciculatis aphyllis, spiculis linear-lanceolatis.

Cyperus scopellatus: Triquetris, angustifolius: radiis umbello ^{apice} fasciculato-
scopoformibus; spiculis striatis, angusto-linearibus Act. Soc. Hist.
nat. paris. P. 1. pag. 106

Habitat in martinica von Rodr.

Palmus erectus, pedalis vel ultra, acute triquetris, striatus, glaber,
utrumque gramen, glaber pallide virescens, inferne foliosus.

Folia inferna disticha, patentia, tri-^{se} vel tri-polliciana, duo vel tria in cul-
mo siquae dimidiis breviora; omnia glaberrima, rigidiuscula, striata,
coria.

Umbella subseptem radiata: radii ^{sepius} pollicares, sensim minores

Stellata Mimosa var. *mic*, foliis stellatis ternis, capsulis ovatis locis. fl. cochinch.
perj. 651.

Habitat in Cochinchina

folia lanceolata, integerrima, glabra. Corol. & calyces brev. Stamina 4.

legumen breve

at hujus generis non vidi

verticillata M. nermis, foliis verticillatis linearibus pungentibus L. Herod
Jest. Angl. Tab. 41.

Habitat in nova caelestia australi 5

non di

forte Funiperona

heterophylla M. foliis simplicibus falcatis ~~per~~ bipennatisque

Mimosa heterophylla ^{foliis} sub ~~non~~ aliis simplicissimis falcatis rigidis
aliis duplicato pennatis La Mark. Encycl.

Habitat in Insula Bourbonis &c.

Rami angulati purpurascens. petioli lineari-lanceolati ensiformes
filiacei nervi numerosi tenuissimi, glabri costa in medio
alii ^{igni} foliati alii apthylli utraque attenuati quadrati vel 5 pol-
licares, ~~foliosi superne ab ex angustioris~~. petioli pariter
alterni ab uno latere costa petioli universalis aequalis
bi v. tripollicares. foliola pinnis novempago lineares oblonga
semurgueculares arenis enervi glabra venella ^{venis} tomentoso
nigra. capitula magnitudine bacii juniperini pedunculo
trigemini

Heterophylla memoa meris foliis bipinnatis pubescentibus, petalis
omnibus planis ramulorum ¹¹ fertiliuim globatis
Dici Bpici ¹¹ globosi Le Mart. Horri. Inst. Nat.
no. 11. 291. Id. 15.

Spica globosa racemosa. Racemi axillares, pollicares. Legumina
linearia lanceolata, subfoliata herb. mus.

a simplicifolia diversa

~~missa in Smith~~

alba Mimosa foliis trijugis. foliolis alleptis acuminatis glabris, petioli aequales,
spicis axillaribus geminis oblongis.

Mimosa alba enervis, foliis pennatis trijugis, pennis aequalibus ovatis acumi
natis, petiolo submarginato Sw. prodr pag. 85.

Habitat in Cayenna In v. Robt.

Rami inferne teretes, glabri, cortice fusco, cinereo punctati, superne angulati,
apice, ^{sub ramulis,} ramis ^{ramis} subtomentosi: ramuli axillares pollicares vel bi
pollicares ^{noctae ad infundibulum confer}

Folia petiolata, alba, quaedam vel quinque pollicaria, pennata, tri
jugia. Foliola subpetiolata, ^{ramis} externa tripollicaria, interna sensim mi
nora, acuminata, basi acuta, integerrima, utraque glabra nitidissima,
fucula, venosa, subtus pallidiora

Stipulae caducae.

Petioli subtus con verti ad singularem par foliolorum marginem tereti parum
laeviores, leviter tomentosi. Glandulae ^{vel} foliales, inter singula foliola, circa
Peduncululi axillares, plerumque gemini, ^{transverse ternij, in ramulis ramosis}
quatuor, ^{quatuor} unguiculares ^{rubescent.}

Spicae oblongae, longitudine peduncululi, calices circiter viginti, conferti

Palyx subtomentosus. denticulatus denticibus obtusis

Corolla glabra, lamina unguiculari subrotunda subangulata ^{nervis}

Stamina monadelphia, numerosa: columna staminum corolla duplo longior

Juaveolens Mimosa foliis linearibus acuminatis rectis margine & adlagineis: primordia
libris pennatis, ramis triquetris Lin. f. Transact. 1. p. 253.

Habitat in Salepa nova austr.

Rami quadranguli, folia alterna ⁴⁻⁵ pollicaria. Racemi axillares

myrtifolia Memoja foliis ovato-lanceolatis oblique undulatis acuminatis
marginibus cartilagineis: primordialibus pinnatis Sen. Soc. Trans. 1. p. 252

Habitat in Valispa nova australi 1788 An. of New Holl. 4. p. 51. t. 15
Rami caulibus angulati. Folia alterna, viridi-glaucia, bipolliciana
Flores copiosi fragrantissimi, tres vel quatuor in fragulo capitulo

Hesperidula Memosa foliis ellipticis oblongis utraque margi
neque scabri, ramulis hespid. pubescentibus, capitulis
solitariis Bot. of new Holl. fasc: 4. pag 53. tab 16.
Habitat in nova Hollandia ~~Tr~~

non vid.