

Mna sum

Rapatea paludosa Aubl.

Habitat in Guiana herb. Guj.

Folia radicalia, petiolata, lanceolata, glauco-viridia, striata, loricata, attenuata, sequipedalia. Scapi compressi, ancipites, superne sensim dilatati. Involucrum diphyllum; cordatum, valde attenuatum, quodsi pollicem planum. Spica plerumque unguiculares, pedunculata, vertice acuto, umbonata, glabra. Squama oblonga, aluminata. Pedunculi unguiculares, angulati. Basi squama.

Cucullata

Propithea cucullata Richard.

Habitat in Guiana Richard herb Guaj.

Calmar subtinquetar, fractus. Involucrum lanceolatum, 5-6 pollicare, canaliculatus. Spica oblonga, involucro dimidio brevior.

1000

Rasitea paludosa Aubl.

Habitat Cayenna herb. Rich.

Scapus foliis multo brevior. Spiculae tenues, umbellatae. Folia lato-lanceolata, basi cuneata, inferne versus angustata, basi latiora, integerrima, glabra.

paludosum Mnasium.

Rapanea paludosa Aubl. fl. Gu. p. 305. Tab. 110.

Habitat in Guiana

obtusa A. foliis linearibus obtusis confertis, caule ramofo tero daci prodecumbente calycibus
centris glabris

~~Alphina foliis linearibus obtusis, calycibus trifidis Hall. hist. n. 863.~~

~~Cuscuta obtusa Cass. fl. Ped. n. 1714. tab. 64. fig. 4.~~

~~Habitat in alpicis Sabaudia &~~

~~Caules plures ramofo laxi prodecumbentes. Solia pinnatifida glabra
Peduncululi terminalis plerumque terni breves. Calycis foliola obtusis
scuta uti fossa non striata, glabra. Petala calyce vix majora
Fructus ovatus~~