

4
5

Monarda (ab tuberoso, 5-dentatus.)
Cor. nigrum: labio superiore linear filamenta invol-
sens. Sem. 4+, n. De

fstulos a 1. Monarda foliis oblongo lanceolatis cordatis villosis planis H. Kaw

1.p. 36.

Origanum fstulosum canadense Porn. canad. 13.t. 14.

Origanum Spurium flore fstuloso Riv. mon.

monarda fstulosa, capitulis terminalibus; caule obtusangulo.

Lin. Syst. Veg. 68

mollis 13. Clinopodium majus virginianum foliis minus hirsutis acutioribus
floribus fstulosis Moris. hys. 3. p. 374

Monarda mollis Lin. Am. Acad. 3. p. 399.

Habitat in Canada?

Paulis obtusa angulatus. Solia bipinnata, semata, utique punctata.

Capitulum terminale. Flore solieri foliola quatuor rotata.

Calyx pilosus. Corolla punctata. *

13. Differentia caule duplo altiore, foliis non pilosis Lin. l. c.

ana M. glabra la Marte. Diet. 4. p. 256 & illustr. t. p. 65 hao
notius pertinet?

Virginitate foliis Virginitate: lobis ovatis serratis R. P. & penne T. b. pag. 14. tab 22

fey. a

Habitat in nemoribus Panahuac C
luisita. Caulo teres, ramosus. Folia opposita, pedata. Lobis ovatis interiore
dis maiore. Pedunculi terminalis corymbosi. pedicelli squamis vel farinaceis
lati inaequales. Corolla C. Tea

oblongata. Monarda foliis oblongo-lanceolatis basi rotundatis - attenuatis
villosis planis H. Kew. t. p. 36. Wieden Spec. pl. 125.
monarda longifolia; foliis oblongo-lanceolatis serratis nudis
bulis, corollis punctatis Lamark. Dict. 4. p. 256. et illustr. 1.
p. 65.

Habitat in America Septentrionali *

Caulis purpureopurpureus, vallis rans adspersa, ad articulos frequentio-
ribus. Solia Tripolitana, attenuata, remota serrata, rare
villosa; ~~ut~~ ⁱⁿ quinq^{ue} puncta involucra cordato-ovata, attenua-
ta, integerrima basi tinctura viridi-coerulea, palea apie
villlosus. Corolla punctata *

~~feruginea C. felicis linearis~~ Languidula, margini revoluti subtus tomentoso ferrugineo
nervi pedicellis umbellatis, cuncte tomentosae.

Radicula feruginea: foliis linearibus, margine revolutis, subtus tomentosis.
umbellis floribus terminalibus tenui Cavan. c. V. pag. 27. tab 44 s. fig. 1.

Habitat ad radiculas Cordilleras in solo sope mundato.

Caulis teret datus, bipedalis erectus, valde tomentosus. Sola opposita, seplex
bipinnata, supra glabra: radimenta ramulorum in axillis. Pedicelli
terminalis brachii: pedicelli feruntur ^{linearum} foliis duobus. Calyx tomentosus. Corolla
lutea: labium superius ~~angustum~~, inferius turridum oratum. Cavan. c. c.

clinopodia 3. Monarda foliis ovato-lanceolatis, basi rotundatis in equalibus glabris Wild. Spec. pl. 1. p. 125.

Monarda foliis ovato-lanceolatis, basi rotundatis in qualibus verticillis lateralibus dichotomis cymbosio, foliis inqualibus serratis Griseb. Vig. 9.

Monarda clinopodia; floribus capitatis, foliis levibus imis serratis Lin. Spec. pl. 32.

Habitat in Virginia 4

Paulus ait Anguli, ut superne, ut: folia reticuli, folia floralia et bracteas subtili raro pilosae. Folia bipinnata et parvum ultra
ovato-oblonga, attenuata, grosse remote ~~de~~ serrata, supra via
punctata; floralia angustiora basi colorata. Capitulum Vermi=
nale, multi-florum. Bracteas exteriores ovato, integrissime purpureo
-violaceo, inter reliquias setae longitudine calycis, reliquias setaceas.
Calycis apice purpureo-violacea, glabrescens. Flores pallidi
Statura M. Didyma; ut folia angustiora, magis elongata; bracteas
purpureo-violaceas, nec ~~sub~~ ^{super} flores diandri, nec ~~Vel~~ ^{sub} ~~sub~~ dynamis *

~~Longifolia Monarda~~

~~Habitat herb. La Manche~~

~~Ramuli glaber, perenne villosus. folia 2-3 pollicaria, subcordato-lanceola-
ta, serrata, subtiliter pungitata, pubescens villosa. Capitulum terminale.
Involucrum integerrimum, villosum~~

Didyma 4. Monarda foliis ovatis glabris, capitulis verticillatis, floribus sub-

Didynamis, caule acutangulo Wild. Spec. pl. 1. p. 125.

monarda caule acute angulato, capitulis terminalibus Lin.

Hort. Cliff. 495. Pold. Novelt. 7.

Monarda didyma floribus capitatis subdidynamis, caule acut-
angulo. Lin. Syst. veg. 68.

Monarda floribus capitatis verticillatisque, caule acute an-
gulato, foliis lanceolato-serratis glabris Büttner. Kun. 226.

Trew. Ehret. 32. t. 66.

Habitat in ^{genesys} ~~tempore~~ Novaboracay.

Caules glaber. Solia bipinniciana vel ultra, serraturis mucronatis;
ad 8 per sola pilis minutissimis rari. Et impunctata. Flores super-
ne in verticillo Et capitulo terminali. Involucra lanceolato-
ovala, uti calycis Et Corollo puniceo *

~~rofmanine folia Pal. foliis linearibus integriformi margino reflexo pubes-
centi spicule glabro de Mart. T Enc. pag. 52.
Habitat in Penn.~~

nigosa 5 Monarda foliis ovato-lanceolatis cordatis glabris nigrois Hoffm.

p. 36. Welden Spec: pl. 1. p. 125.

Habitat in America boreali 4

Similis to antecedentia; at folia longiora, glabra, nigra, insula, per-
more Salvo, et florei albi Wild. l. c.

M. glabra L. Marti. Diet. 4. p. 256. illustr. 1. p. 65. huc forte spectat

M. glabra L. Marti. Diet. 4. p. 256. et illustr. 1. p. 65. huc forte
spectat

9

~~longiora~~ ~~foliis~~ ~~ovato-lanceolatis~~ ~~cordatis~~ ~~glabris~~ ~~nigrois~~

~~affinis~~ ~~Monarda~~ ~~spicata~~ ~~monspeliensis~~

~~36. W. C. 81. v. 1. p. 125~~

~~nigra~~ ~~mono~~ ~~affinis~~ ~~foliis~~ ~~ovato-lanceolatis~~ ~~cordatis~~ ~~glabris~~ ~~nigrois~~

~~Capitata Ziziphora foecit aliis terminalibus, foliis ovatis s. v. p. 67. t. Sp. p. 1 p. 24~~
~~Plinopodium fistulatum pumilum Inde orientalis oie dentalis, summo callo
flondum Plin. alm. 33. t. 164. f. 4.~~
Thymus humilis Cat. folius Bust. Cest. 3. p. 28. Tab. 51. f. 1.
Habitat in Syria, Armenia. O
Pauli erectus ramosus pubescens palmans. Folia petiolata
oppoſita integrer recta glabrescens. Pedicellus circiter 7 fer-
minales uni flori ^{lunati} foliis quatuor ovatis nervosis attenuatis
ciliatis. Calyx veluti fimbriatus laevius setaceis subequalibus

bus ovatis

~~flor caerulea elliptica lanceolata; florah~~

allopelta 6 Monarda. foliis oblongis serratis, capitulis terminalibus, calycibus
limbo barbatis +

Monarda allopelta; foliis oblongis, acute serratis: Capitulis terminali-
bus: calycibus ad limbum barbatis, corollis graciliter elongatis, car-
neis Michaux fl. bor. am. t. p. 16.

Habitat in America septentrionali

ciliata b. Monarda foliis eratis attenuatis, cuncte radicibusque hispatis,
bracteis eratis calycem opeantibus.
Clinopodium angustifolium non ramosum, flore coeruleo: labio trifido
atro purpureo maculis ornato mons. hst. 3. p. 374. 8. 11. 1. 0. f. 6. Pl. 10.
alm. 10. t. 164. f. 3.

Monarda spica interrupta involucris longitudine verticillatum can-
ceolatis Gronov. Virg. 9.

Monarda ciliata floribus verticillatis, corollis involucro longiori-
bus Lin. Syst. Veg. 69. Ex herbario

Habitat in Virginico, Carolina. Hab. Guspius

Caules ramosi; a juliis longis albido hispitoso prossimim superne. Dolia
biplicata, basin versus integririma, ~~stetigae punctata~~, villosa
nigra minutiis ad persa, Verticilli superne, duo ad quinque inferio-
remoti, nos albido hispiti. Bracteas in anastom attenuatas. Flores
nervi hispiti, coecines paucitati

Sacces aliena a reliquis. Odor mentha

~~fistulosa Monarda capiteles Verminalibus, caule obtusangulo s.v. 63. t. Specpl 32.~~

Mill. Diet. 7.122 v. 2.

~~Organum fistulofum canadense Corn. canad. 13. t. 14.~~

~~P. Monarda mollis Lm. Cred. 3. p. 399.~~

~~Clinopodium majus virginianum, foliis minus hirsutis acutioribus; floribus fistulosis~~

~~si Morf. h. 3. p. 374.~~

Habitat in Canada 4

~~Pauli Tetragonus glaber anguli obtusus superne subvillosum. folia
opposita ^{tota} juniora villoso, ferrata hypoleuca. cap. flore
in capitulo terminali collecti sunt foliis quaternis ovatis
subintegerrimis involucrati calycis petiolis florae pallide
purpureis estriis villosi oculi armato-punctati~~

~~foliis cordato lanceolatis attenuatis capitulo ^{subvilloso} vello
aliam quam sub nomine M. clinopodii accipi caule angulis acutis~~

~~Leonurus canadensis organo folio Tengst. Hoff. 1087
ulte~~

punctata ♂ Monarda foliis lanceolatis serratis, involucris floribus verticillatis
involucris ovali lanceolatis verticilli longioribus
Clinopodium angustifolium, ^{virginianum} ~~camii florid. luteo~~ maculato
fido, foliis purpureis maculis ornato. Mon. hist. 3. p. 375. S. II. t. 8.

J. 8

Clinopodium angustifolium ^{angustifol.} ~~virginianum~~, quovis verticilli duo
decim foliis rubentibus cincto. Plakn. alm. III. t. 24. f. 1.

Monarda punctata; floribus verticillatis, corollis punctatis, bracteis
coloratis Lin. Sys. Veg. Cg.

Habitat in Virginia ♂

Paulis eretis, ramosus, villoso subbianescens, obtuse angulatus.
Solia Besqui polliciana, versu medium serrata, glabra nervosa,
utriusque uti calycis punctata. Verticilli superne, duos ad quin-
que. Involucris foliis quatuor longioria, duos brevioria, basi ciliata
super violacea. Calyx defloratus dentes recurvis. Corollo hispido,
flavo, punctato, rubro punctato

Variat foliis magis minusve serratis, supremis serratis vel in
tererrimi vel serratis.

#punctatus,

~~8 punctata Monarda foliis ^{lanceolatis} vestiellatis, corollis punctatis, bracteis coloratis. V. p 68. 4
Spec. pl. pag. 32. 4. Mant. 317.~~

~~Clinopodium virginianum angustifolium lamii flore luteo maculato
Mon. h. 3. p. 375. f. 1. V. S. f. 8.~~

~~Clinopodium virginianum angustifolium, quoris vestiello duodecim
foliis nubentibus cuncto Plat. Alm. III. t. 24 f. 1.~~

Habitatia Virginia

~~romofus~~

~~Pauli erectus Villosum Tetragonum folia petiolata ferratis
aliquot in medio utriusque ut involucrum calyces corollaque
punctata glabra, sed qui pollicari. Involve sa lanceo
lata pallidiora et breviora intus purpurea integerrima
et lesta flore longiora. Corolla pilosa. vestielli 4.~~

~~foliis lanceolatis subferratis utriusque punctatis; involu
cri corolla longioribus lanceolatis coloratis~~

Habitus Paulis Tetragonos² Lebaient:
Hs (rami oppositi. folia petiolata, opposita, ^{indivisa}, (remotis cule serrata). Flores
verticilli vel capitula multiflori, involucrati. Bracts setaceo. Ralyo-
g. foliatur.

aterrimus

~~st. florae albore~~
Bellabunga Veronica racemis; foliis ovatis planis, caule repente S.V. Et. S. n. 14. Spec. p. 16. n. 14
Et. D. t. 511 Reg. P. n. 14 Lecoufle Herb. no. Poll. h. n. 10
anagallis aquatica, folia matinno Thal Sun Tuck. her. 725
anagallis aquatica Lob. f. 466.
Sun Brunsfer 16 gen. em. p. 620

3. p. 169

~~Hedysipper~~ Dost. p. 5.

~~Anagallis aquatica major folia sublundo Bauch. p. 252.~~

Anagallis aquatica, foliis rotundiore major Bauch. h. 3. p. 790

Veronica aquatica, folia sublundo Tournet J. L. H. p. 102

Anagallis aquatica Blattwelt t. 48

Veronica foliis ovatis, panicatis, glabris ex aliis racemos Hall. h. f. p. 634

B. minor non ordinatum Trag. Anagallis aquatica Trag. 187

~~Sic Bajntel non ordinatum Trag. Anagallis aquatica Bauch. p. 252. Bauch. h. 3. p. 791.~~

Anagallis aquatica minor + Bauch. p. 252. Bauch. h. 3. p. 791.

V. flore albo Blair Des. p. 25. t. Tabern. f. 710

Habitat in Europa ad seculum vix congelatae. Siberia. Amer. septentr. 24

Tot. globosa

+ Tabern. p.

~~bifida~~ ^{repens}

ter

var. flore albo Paulus repens bifidae poviun ~~bif.~~ ^{er} repens, globosior. Folia brevissime subfusiles, oblongo, obtuso serrata. Racemi laterales oppositi foliis petiolatis, subfusiles, oblongo, obtuso serrata. Racemi laterales oppositi foliis nervosis, triploblongior. Pedicelli filiformes ~~bif.~~ ^{er} bractea linearis. Calyx sp. avenie quadratis laevis duobus oblongis, acutis majoribus, corolla patente connexa. Capsula subrotunda

Moparda

1. foliolata M. foliis cordato-lanceolatis attenuatis vellosi; calycibus vellosi
2. oblongata M. foliis oblongo-lanceolatis basi rotundatis attenuatis vellosi
3. didyma M. foliis ovatis serratis attenuatis involucris calycibusque coloratis glabris.
4. denudata M. foliis ovato lanceo oblongo glaberrimus.
5. rugosa M. foliis ovato-lanceolatis cordatis glabri rugosi
6. punctata M. foliis lanceolatis sub serratis utrinque punctatis, involucris corolla basi orbiculis lanceolatis, floribus vestitellatis
7. ciliata M. foliis corollis involucris
conygombris vestitellatis

Moparda est America boreali odore menthe dicit Richardus.