

Nephtelium

1. Ceylanicum N. foliis pinnatis
2. castaneifolium N. foliis simplicibus serratis

Generis ejusdem ac Syntalea vel litsehi secundum La Billardiere

Sephelmum

Caribbean *Asythanica* Brown-jam 330. In prodr: pag 39.

But no affinis *odonifera* *fruticosa* minor etc.

Sloan hist. 2. 132. t. Ob. fig. 3.

Habitat in Jamaica &

Caulis teres ramosis superne angulatis canescens fruticosus. Folia
subpetiolata alterna, prope polliciana oblonga venosa integerrima acuta
glabra ~~juv. serrata~~ venella sericea. peduncululi axillares solitarii
quadri flori flori unicus foemineus. Capsula parva thecacea incana

Cappaeicum Nephelium Lin. Mant. 125 * S.V. pag 851

Habitat in India Orientali

Rami Veretes glabri. Folia pinnata fixa cum pateris, semper
Dalia trijuga vel bijuga. Folia laevia petiolata subopposita
Sini f. quadripollucana ^{integerrima} ~~est~~ ^{est} ~~ima~~ ^{ma} ~~mezora~~ ^{mezora} elliptica
glabra ^{tenuissima} venos reticulata. Pedunculi axil-
lares ^{et terminales} ~~solitarii~~ ^{rami} longiores ramosi rami simplicis-
mi longi alterni e caepe florum tuberculati. Flores
parvi subsessiles

Folus pinnatis

castaneifolium N. foliis simplicibus

Habitat in Bengala India orientali Siam &

Rami teretes glabri fere folia petiolo subopposita vel altera
na 6. 8 pollicari aut uncinata utraque acuta alter
nati in oblique nervos obscure venos nervi subtus purpi
rascentes, glaberrima obtuse remote serrata hirtus pal
lidiora. Petioli bipollicari. Folia tenella utraque
lanuginosa. articulum foliola duo prothulata simulata semi
unguiculata in articulis. Pedunculi axillares et
terminalis axillares simplicissimi terminalis in
topi utraque varicosos flosculi primus imbricati demum
remotiores ⁱⁿ ~~in~~ oblonga parva longitudine floris apice
patula ad squamulum florem. fructus ut in Cap
naero

confertur cum aucubus Stipula autem
omnino diversa