

Cenothera

Calyx quadrisfidus tubulatus. Petala 4. Capsula 4loculari, 4valvi. Semina nuda

1. biennis O. foliis ovato-lanceolatis, planis, caule mucronato villosa
2. grandiflora O. foliis ovato-lanceolatis, staminibus declinatis, caule
tuberoso.
3. parviflora O. foliis ovato-lanceolatis planis, caule basi subvillosa
4. undulata O. foliis linear-lanceolatis obtusis subundulatis floribus termi-
nalibus aggregatis
5. fasciculata O. caule foliisque lanceolatis petioli, floribus fasciculatis,
prope tuberoso. Capsulis oblongis acutangulis
collocatis
6. mucronata O. foliis lanceolatis villosis planis, caule villosa mucronato
7. longiflora O. foliis denticulatis, caulibus simplicibus villosis, pedalis
O. fraxibus trilobis
8. mollissima O. foliis lanceolatis molliter villosis undulatis
9. nocturna
10. sinuata O. foliis dentato-sinuatis, capsulis prismaticis

11. foliosa O. foliis lanceolatis - ovatis, inferioribus sinuato dentatis
 floribus racemosis, capsulis angulatis clavatis
12. fruticosa O. caule aereo piloso fruto, floribus racemosis capsulas
 acutangulis, foliis lanceolatis subdentatis, calycibus aristatis
13. rigidula O. foliis lanceolatis subdentatis capsulis axillaribus pedicel-
 latis acutangulis
14. humilis O. foliis lanceolatis obtusis glabris subnitidis, cauli-
 bus prostratis capsulis axillaribus acutangulis

ex
herb
mus. { Gen. ex insula Francis
Rami crassi ^{sub} oppositi. Folia opposita Racemi terminalis compositi
Onagra acedardi minima hys/soyifolia
Saulis ramifera

Oenothera grandiflora La Marche } herba mans.
Odonia hort. parif.
albicans La Marche hort Parif }

p. prostrata *Oenothera*

Habitat Commerçon herb. Guss.

Caules prostrati. Folia radicalia, uti et interdum caulina lanceolata
caulina plerumque oblonga, tomentum fere totum dependentia. Flores lutei
extus purpurei

Oenothera

Herb. Desfont.

Caulis herbarae erectus, teres, basi folia villosi. Folia sessilia, pollicum longa, canaliculata,
avenae, enerviata, dentato-serrata. Flores axillares, solitarii, sessiles.
Germina hirsuta. Corolla purpurea.

Corymbosa. Oenothera

Attabitat herb. la Mark.

Caulis ope lentis pubescens. Folia lanceolata, integerrima, glabra. Calyces solitarii in axillis supremis, canescentes, breviter pedunculati.

tetraptera Oenothera foliis. albis sub bipinnatis crispis, fructu
tetraptero Cavan. 4i vol. 3 pag 40. Tab 279
Habitat in nova Hispania &

~~novi~~

1. biennis *Oenothera foliolosa* ovato-lanceolata, planis, caule mucronato villosa
Lin. S. V. pag. 350. Fl. Dan. t. 446.

Hypoxymus virginianus Alp. exot. 325. t. 324

Lysimachia lutea corniculata Bauh. Pin. 245. 516. Mon. hist. 2. p. 277
fig. t. 11. fig. 7.

Habitat in Virginia, unde 1614. nunc vulgari spontanea in Europa et
caulis bipedali ^{erectus}, teris crispis pennis asperis, simplex. Folia
tripolliciana ^{versusque} ~~sub~~ basi acuta, remote obscure dentata glabra. Flores
sespites in axillis, supremo conferti longitudine foliorum. Germina
et tubus calycis pubescens lacini linearis lanceolata

lanceolata elliptis planis, caule submucronato
capsulis apice quadrifidis

Oenothera
2 *Grandiflora* *foliis ovato lanceolatis, flaminibus declinatis, caule fruticoso*
Bot. Kew. 1. p. 2. L'Herb. Prop. nov. Tom. 2. t. 4.
Habitat in America septentrionali ♂

non vid.

4 undulata *Oenothera foliis lanceis canaliculatis obtusis subundulatis floribus*
terminalibus corymbatis sessilibus

Habitat

Caulis herbarius ^{erectus} crassitie pennis, arboris ^{perispermis} purpurascens ^{et villosus}
foliis sessilibus ^{perispermis} ^{et villosis} basi attenuata obscure dentata. Flores
plurimi in apice caulis sessiles germen villosum oblongum mini
ne angulis compressis ciliatis, Tubus brevis pubescens, Calycis
foliola aniseta tubo ^{longiora} flores Corolla ut in parviflora

6 *municata* *Oenothera* foliis lanceolatis ^{vellofis} planis, caule piloso *municato*.

Oenothera *municata* foliis lanceolatis planis, caule purpurascente *municato*
Lin. P. V. p. 35 & Murr. in Bon. Jöb. 1775. p. 24. t. 1.

Habitat in Canada.

Caulis erigitur simplicissimus crispis pennis adpresso, pilis turberculisque minus
frangis. Folia bipollicaria, sparsa ^{oblongata} utrinque acuta subtus pilosissima
^{molliora} flori ex axillis supremis alternis confertis. Tubus et calyx villosi ^{canescenti}
Corolla magnitudine *Oen. parviflora*. Involucris unguicularis

7 Longiflora *Oenothera foliis denticulatis, caulibus simplicibus pilosis petalis di-*
stantibus vel bilobis Lin. Mant 2274 L.V. pag 350. Faug. bot. 2. p. 81.
t. 172.

Habitat in Agro bonanensi S.

Folia radicalia numerosa, lato lanceolata, denticulata, pubescentia
costa alba, oblique nervosa. Caulis semper 5, enati infra folia radi-
calia, simplicissimi ascendenti, hirti, vix pilis longis patulis
albis. Caulis centralis tenuis excrevit. Folia caulina ovato
oblonga vel lanceolata, sessilia, versus basin integerrima. Flores
axillares superiorum foliorum, germine calycis piloso. Calyx
tubo insigniter elongato, prothecanis rare pilosa, limbo
suo triplo longior foliolis linearilanceolatis. Petala obtunda
et magna altero die rubra Lin. l. c.

varietatem hujus? communiavit in Thora in vtilitate longi
refertur ~~tudine tubiformis diversa foliis unguiculatis et parum~~
~~angustioribus obtusissimis forte spontanea~~

lanceolatis

hirtis

4 mollissima *Cenothera* foliis lanceolatis molliter villosis undulatis. Lin S. V. pag 358.
Onagra bonanensi villosa, flore mutabili Dill. Elth. 297. V. 219. fig. 286.
Cenothera foliis linear-lanceolatis dentatis, floribus mediocaula
Lin Hort. upf. pag. 144.

Habitat in agro bonanensi ☉

Caulis teres pubescens ~~in~~ ⁱⁿ tota planta. Folia sessilia angusta, sesqui
pollicaria remote dentata. ~~in~~ ⁱⁿ ~~superioribus~~ ^{superioribus} attenuata
foliorum. Capsula cylindrica longitudine folii inferne
parum angustior.

*Nothura Oenothera foliis lanceolatis denticulatis subvillosis, floribus axilla-
ribus sessilibus*

Oenothera nocturna Jacq. Coll. 3. pag. 205.

Habitat in Cap. b. p. c.

Suffruticosus. Caulis bipedalis, erectus. Vires mollis, fulvus, subvillosus ab ipso
basi ramosus. Folia alternas, brevissime petiolata, lanceolata, acuta
crenato-serrata, subvillosa, quadripollicaria. Flores axillares solita-
rii sessiles. Germen quadrifidum, villosissimum. Petala rotundata
ample lutea. Capsula villosa octodentata quadrivalvis Jacq.

11 sinuata *Oenothera foliis dentato-sinuatis, caulis apice nutante* Lin. Mant. 228, ^{capsulis prismaticis Hort. kew. 2. p. 3.}
Murr. in Com. Gott. 1774. p. 44. t. 9

Lysimachia corniculata manthana, sinuatis et pubescentibus foliis
virginiana Plukh. alm. 235. t. 203. fig. 3.

Oenothera sinuata: foliis dentato-sinuatis, caulis apice nutante. Lin.
Mant. 228* l. V. pag. 358. Murr. in Com. Gott. 1774. p. 44. t. 9

Oenothera repanda Medic. in act. Palat. vol. 3. off. phys. pag. 190. Tab. P.

Habitat in Virginia O.

Caules simplex sesqui pedalis erectus apice nutans, junior pilosus, senior
glabriusculus: rami alterni. Folia sessilia, lanceolata, pollicaria
nuda, obliqua ^{subperennantia} & loco axillari sessile solitarii pilosi: calycis et
oculo ornato ^{germini} ^{germini}. Calyx acutus sub apice mucronatus. Germen ante anthe
dispersa
sin declinatum. Petala glabra obcordata longitudine foliorum
calycis. Capsula cylindrica quadrangulata, angulis planis, truncata
ata apice profunde excavato quadrifido; lobis emarginatis:
quadri-valvis Lin

rosea *Oenothera* foliis ovatis dentatis, inferiorem lyratis capsulis clavatis

Bot. New. 2. p. 3.

Oenothera rosea L'Herit. Supp. Nov. Tom. 2. t. 6.

Habitat in Pem 4

alternis

Caulis basi lignosus, prostratus vixatus. Ramis teretibus vix villosis.

Folia petiolata, utrinque acuta, glabra ^{pubescentia} ^{pubescentia} glabra, juniora

canescentia. Racemi terminalis cum ramulo ulterius est crescente

demum laterales longiores flores remotissimi 5-6. pedicellati

secundi. Capsula turbinata ^{folium lineare, ad basin juncti} ~~sub~~ compressa angulis 3 acutis quorum

duo laterales majores. Germina cana. Flores purpurei

basi interdum
dente uno vel
altero obtuso

pedicelli angulati
superius membranati

eodem 25 purp

^{superfusa}
 13 ~~florata~~ *Oenothera* ^{Differencia inde a Syst} ~~caule erecto piloso fructu, floribus racemosis, calycibus anfractis,~~ ^{resiliantibus.}
~~capsulis angulatis pedicellatis~~
 an grandiflora Hort. kew. vel potius frutescens Lin.

+ ^{Habitat} *Virginia* 4 ^{purpurascens - rari rare velle ofus}
 Caulis inferne ^{capiteolata} ~~lyriophylla~~ ^{et} ~~simplex~~ ^{stramineo acuta pubescentia} ~~et~~ ⁱⁿ ~~lobis~~ ^{dentata}
 folia ^{floribus} ~~ovata~~ ^{pedunculata} ~~et~~ ^{ex} ~~apice~~ ^{caulis} ~~et~~ ^{ramulorum} ~~superiorum~~ ^{con-}
~~uncane~~ ^{nimum} ~~purpurascens~~ ^{demum} ~~et~~ ^{racemosa} ~~et~~ ^{floribus} ~~subsessilibus~~ ⁴⁻⁵ ~~et~~ ^{calycibus} ~~anfractis~~
 an ^{minuta} ~~et~~ ^{super} ~~terminati~~ ^{germen} ~~glabrum~~ ⁱⁿ ~~semunquam~~ ^{ulare}
 acutis ^{angulis} ~~quatuor~~ ^{acutis} ~~et~~ ^{compressis}

+ ^{staphylli} ~~alters~~ ^{latere} ~~plenumque~~ ^{rum} ~~pentes~~
 + *Onagra* ~~angustifolia~~, ~~caule~~ ~~nubro~~, ~~flore~~ ~~minore~~ ~~Tournef.~~ ~~f. R. H.~~

302
Oenothera ~~florum~~ ~~calycibus~~ ~~monophyllo~~, ~~hinc~~ ~~tantum~~ ~~aperto~~ ~~gon.~~ ~~Viry.~~

43.
Oenothera ~~foliis~~ ~~laniccolatis~~, ~~floribus~~ ~~terminalibus~~ ~~paniculatis~~
 Joan hort. 193.

Oenothera ~~frutescens~~: ~~foliis~~ ~~laniccolatis~~ ~~subdentatis~~, ~~capsulis~~ ~~pedicellatis~~
 acutis ~~angulis~~, ~~racemo~~ ~~pedunculato~~ ~~Lin. Spei. pl. puz. 492 & 5. V.~~
 p. 35. D. L'Herit. ~~flor.~~ ~~nov.~~ ~~Tom. 2. Tab. 5~~

rigida

15. *pumila* *Oenothera* foliis lanceolatis obtusis glabris, suppetiolatis, caulebus pro-
stratis, capsulis acutangulis Lin. Spic. pl. pag 493. & V. pag 350
Lysimachia foliis glabra minor manana, angustibus foliis Pluk. man
123.
Lysimachia manitandua parva, foliis angustis acutis Raij sup. l.

468
Oenothera foliis radicalibus ovatis, caulinis lanceolatis obtusis, cap-
sulis bivalvis obtusis Mill. Dict. v. c. 100

Habitat in America septentrionali &

Caulis palmaris, teres, flexuosus, divaricatus, minus leviter pubescens
Folia inferiora latiora breviora, superiora longiora angustiora
integerrima. Flores axillares, solitarii, sessiles. Petala lutea, obor-
dato, lineata. Capsula obovata, obtusa, octangularis, angulis
quatuor latioribus compressis, quadri-valvis. Lin.