

Plat.

82 Plat. Cal. margo integer. Cor. 5-petala, singula duo mediante fila-
menti contractis, quinta libera. Nect. ~~folia~~ foliola 5. pinnata
Drupe. manet 1-p parva (nata Gussien)

Zeylanica 1. Olax foliis ovatis acutius, ramis angulatis

Molla holla Herm. Zeyl. 13. Burm.

Cibus stenocaria zeylanica glandifera Burm. Zeyl. 26.

Olax zeylanica Lin. fl. Zeyl. 34.

Olax zeylanica Symb. bot. 3. p. 7.

Habitat in Zeylona ^{Herzoni.}

Rami flaccidi, glabri, cortice viridi flavescenti, ut in Vesco, trans
versim tenuis vena rugosi. Folia bi- tri-pollicaria, tenuis vena nervosa,
ret venosa. Racemi simplicis vimi, quinque flori, plerumque solitarii.
Flores minores, quam in sequente.

a dissilio genere non differt, quod enim Linneo Nectanum dicitur
Susseo stamina sterilia sunt

Zealandia 1. *Clax folio ovato acuminato*

Psittacorum 2. Olas foliis ovato-oblongis, lanceolatisque subvenosis, ramis ter-
tibus.

Dissilia Fuss. gen. 260.

Dissilia psittacorum Lamark. illustr. t. p. 102. t. 20.

Habitat in insula Mauritiana, & in Commerson, in Zeylona

König 5

Rami glabri, juncti rigidi, glabri, punctis griseis adspersi. Folia
seriquipollucari, ovato-oblonga vel lanceolata, interdum ovalia, subovata, crenata,
obsoleta venosa, pallide viridia. Racemi simplicissimi, rarissimi
mae superne ramulo cauti, interdum biflori, interdum usque ad
viginti flori: pedicelli interdum pedunculo longiores. Calyx maris
brevis, integerrimus, persistens, intergerimus. Petala quinque, quatuor
duum parium mediante filamentis cohaere connata, quantum li-

brevis & imi, simpli
& imi: pedicelli
uniflori

berum. Nectaria quinque, filiformia, medio petali inserta usque lon-
gitudinis, basi macula nigra notata. Stamina tria, costella breviora. In-
na pino duplo major, calyce ad medium tecta.

Aliam habeo, foliis tenerioribus, magis venosis; ramis rubescentibus, forte mo-
do varietas. Ramis flexuosis, alternis. Folia pedicellata,
Habitat arboribus semper viventes: Ramis flexuosis, alternis. Folia pedicellata,
costella, integerrima, obtusiuscula, glaberrima, lona. Racemi axillares.

volubilis Hippocratea Lin. S. V. p. 83

Poa scandens fructu gemino subrotundo Plum. gen. P. G. 80.
Beyuco pendulus, floribus paniculatis Coefl. 41. 314.

Hippocratea scandens Jacq. Aust. t. 4

Habitat in America Australi ~~et~~ ~~habitat~~

volubilis H. medemii corymbosis, foliis ovato lanceolatis serratis Sw. pag. 17.

ramis tereti-angustiori ramulosis ramali oppositi ^{obtusae vel obtusae} foliosi et florigeri. Folia
ovata vel lanceolata nitida ^{obtusae vel obtusae} ciliolata opposita 3 pollicia
na ^{extensa in umbrosis} geminalia. Corymbi axillares oppositi rari
terminalis. Capsula glabra Jacq.

non longe, ramis tetragonis glabris, super femi quae ceteris tomentosi. Folia petiolata oppositi
tri polliciano ovato-oblonga ^{in quinque glebas, nitida, obscure} ^{tenues demum parvas} ^{superficientia}

retrofoli per ^{lofi}, dichotomi, cum pedicelli duobus brevissimis in ^{duo} ^{tomem}. Flores
parvas ceptis ^{parvis tomentis} ^{hab. Guj.} in Lin. Archid a Cojessa

Frugum planta diversa a Plumierii

~~non~~