

Oroonium

aquaticum *Orridium foliis lanceolatis levibus* ^{subnerviis floribus confertis} *Lin. I. V.*
Anum fluctans, pene nudo *Wanest. Virg. 1924. Catess. Carol. 1. p. 82. & 82.*
Anum folio nervi ovato *Gronov. virg. 113.*
Orridium aquaticum *Cemon: Acad. 3. p. 17. & 1. fig. 3. * Kalm d. 3. p. 84.*
Lin. L. V. p. 1. spec. pl. p. 263
Habitat in Virginia, Canada paludibus, scaturiginibus &
folia lato lanceolata, integerrima, glaberrima, petiolata, subtlespi-
me nervosa, & Scapus teres longus, pallidus punctis fasciis. Spadix
pollicis aut teres

2 japonicum *Orethium foliosum* venosum ^{scapis foliis brevioribus} Thunb. fl. jap. pag. 144 * p. 340
Kuro et Kirzo, vulg. ornitho. Kompp. Am. ex. V. p. 785.

+ Habitat in Japonia

Folium radiceale solitarium, ensiforme; inferne attenuatum, convolutum;

- Venis pluribus longitudinalibus venosum, integrum, glabrum, duo pollicari

latum bipedale. Scapus 4-eres, glaber, ensibus digitalis vel palmari

Spica oblonga, pollicari

ab *O. aquatico* differit folus longioribus, venis plurimis elevatis

scapo breviori

+ *Orethium foliosum* venosum Thunb. fl. jap. pag. 144. *

Ref. Linn. S. V. p. 340.

Jamaica für Trichter

3 submersum *Oxordium* foliis lanceolatis utrinque attenuatis ^{7-nerviis} scapo longiore
scapo longioribus, floribus distinctis
Habitat in India orientali

~~Radix submersa~~ *Solanum* pedalis et ultra, brevis petiolata, undulata, tenera, nervis
septem longis adnatis, transversim parallele venosa; venis
vix ramosis. Scapus erectus foliis multo longior, superne crassior
Spica bipollucaris. Flores alterni, remotiusculi & distincti
differt a reliquis quod duplo major, foliis brevis petiolata
ratione longitudinis folii bipollucaris.