

Panicum

ventricosum Pan. panicula ramosa, glumis ventricosis obtusis nervosis
pubescentibus, culmo basi repente La Mark T. Eur. pag 173
Habitat in Andra orientali.

eodem cum curvato credit La Markius Divessem videtur
flexuosum Retzii fide herb. La Mark.

Coffetium Pan. panicula capillari late ramosa, glumis ~~atris~~ vixi
acuminatis, culmo filiformi La Mart. T. Em. pag 173
Habitat in Amer meridionali

byflorum Pan panicula capellani flexuosa, flentibus geminis, vaginis longis
dinaliter ciliatis La Mark T. Enc: pag 174.

Habitat in Tripula Francie

an breifolium Lin. Cal. 1 valvi valvula interiore deficiente

brevifolium P.

paniculatum, foliorum vaginis longitudinaliter ciliatis. Lin
Spec. pl. puz 974

~~Panicum miliaceum vinde, foliis lato brevibus, panicula capillari, femine albo Sloan. hist. 1. p. 115. t. 72. fig. 3.~~

Panicum miliaceum latiore folio, maderappatanum Pluken-
etm. 176. t. 189. fig. 4.

Habitat in India

culmi repentes, demum ascendentes geniculati geniculis fuscis

colis ovatis ^{usque sub quadratum} acutis vel lanceolatis striatis frequentia polliciana

supra orulo armato vel ultra basi denticulata denticulis pilo terminali vaginis striatis

pernatis punctis glabris excepto margine inferiore articulo breviter sursum vel
pubescenti ^{sub} panicula ramis raris capillari-
bus sensim longioribus

glo flosculi parvi ore latis villosuli oblongi altius latius con-
vexius gibbosi distantes valvula oblonga nervosa breviter
turbiformi

panicula capillari flosculosi gibbosi, foliis ovatis lanceolatis
et basi denticulatis ciliatis culmo repente

radice anni Petru hui refer diversum foliis magis lanceolatis et
flosculis glabris

radicans P. paniculatum, culmo ramoso radicante, foliorum basi vaginisque
longitudinaliter ciliatis Retz off. 4. p. 104
Habitat ⁱⁿ ad China ad Canton

ca a Pan: brevifolia non diversa

puberula subaequalibus
pungens Pan. panicula patente ramis simplicibus glumis bifloris; folia involu-
ti pungentibus ^{depressis} hakenas
Habitat in ~~Asia~~ ^{India} Forsk

Culmi ramosi rigidi glabri. folia tri-⁴ angulicarpa patentia rigida
glabra Panicula terminalis ^{rami} ~~remoti~~ ^{approximata} duo per intervallo ^{remota}
subsimplicis glabri flexuosi glauculi pedicellati ^{seu foliis} ^{perisperm.} ^{perisperm.} Val-
vula ovata striata subaequali. flori alteri hermaproditis semina
cava
gramen huius ~~est~~ ^{est} tam simile ex India occidentali habui ut nulla
nota ^{distinctionis} ~~distinctionis~~ ^{diagnosticum}

Conglomerum Solanum inermis, foliis ellipticis integris ~~attenuatis~~ ^{subtus} subtomentosis, racemis lateralibus, laciniis corollae lineariibus: spicatis

Habitat Capanno In v. Proho.

Indes

Rami teretes, inferne glabri, superne a ^{velis} ~~pube~~ minutis stellatis fere ^{et} fomentosis

Folia petiolata, alterna, remota, tri vel quadri pollicari basi obtusa, subequalia, apice attenuata, acuta, integra, ^{inferioribus} supra a ^{velis} ~~pube~~ stellatis et oculo armato videndis scabriuscula, subtus vellosa fere fomentosa

4 adultorum
fere glabra
obscuri venosa

profertim juniora
Petioles pollicares, inermes.

leniter linguas recurvati: demum secto ^{res}

Pedunculis superne, lateralibus, brevibus. Pedicellis pluribus, alteris, 8-10. un-

guiculis: Calycis fomentosi, inermis, uti corolla extus: Dentibus quinque, subulatis, corolla quadruplo brevioribus.

Corolla fere pollicari, profunde quinquefida: Lacinis ^{crenatis} Canaliculatis, attenuatis, intus glabris, a

(Facies sol. obscuri, at totam inermis)

pallens

~~axillare~~

Paniculi terminalibus axillaribusque subsimplicibus, flospulis
lanceolatis attenuatis, foliis lato lanceolatis

Habitat in Montserrat

Pedunculi 2 vel 3 ex ^{fope} singula axilla folio duplo longioribus
glumis bifloris

Aplud aegypti Aubl. fl. Guian.

Panicum pallens panicula composita ovata, ramis confertis
erectis, spiculis ovatis subulatis, culmo duro geniculato,
foliis ovato-lanceolatis, vaginis collo marginisque ciliatis
W-prodr pag 23

sub nomine P. commelinae

reliqua vide in ter americana sub nomine Pan commelinae
~~quod non diversum~~ deserventer a Richards

Longiflora Gent

^{Capellan}
viscosum Pan. panicula composita pedunculis flopidisque glutinosi
Habitat in India occidentali?
Culmus ramosus tenuis. Folia plana bipollicina angusta glabra
vagina circum collum villosa. Panicula terminalis rara pedun-
culi flexus capillares. flopidi oblongi

Hermas

1. gigantea H. foliis ellipticis ^{ret. latis} supra lanuginosis subtus albis tomentoso albis
obsolete crenatis.
2. depauperata H. foliis cordato-oblongis, semiamplexicaulis subtentatis
supra glabris subtus tomentosis.
3. capitata H. H. foliis subcordatis obtusis petiolato obtuse serratis subtus
cauleque tomentoso.
4. quinqueidentata H. foliis ovatis quinqueidentatis subtus tomentosis, umbella
solitaria, caule glabro.
5. ciliata H. foliis ovatis ciliatis subtus tomentosis, umbellis pluribus, caule
glabro.

Diffructum P. Panícula subsimplici corymbata, flo. valvula glumae biflorae foliis
lanicolatis diffructis scabris

Habitat in India orientali.

Caubis infractis ac basi ad paniculam secti vaginis foliorum glabris
marginibus ciliatis. ^{crassitie pennae columbinae} Folii articuli inferiores pollicares superiores per
sum majores. Folia lanicolata diffructa sessilia sesqui polliciana
inter nodos et vaginis longiora glabra a ^{sum} puncti minuti eleva-
ti biculo armato videntis scabris glabra vix glaucescentia
panicula bipollicari glomeruli gemini pedicellati globosi
pedunculati scabris. valvula glabra non obtuse st nata non nervosa
acutifera nulla. valvula corollina ore lenti villosa

affinis *Stenandri* uti auctori

~~ar millonii *Stenandri* foliis obtusis cum acuminis ^{sum} l. v. long. 229
strenandri ar millonii foliis obtusis cum acuminis ^{sum} l. v. long. 229~~

arboresca. ~~Faiquinia arboresca, ramis inferioribus quaternis verticilla-
tis ceteris superioribus dichotomis~~
~~Habitat in ~~monte~~ vespis summitatei montium Montserrat~~

~~In Ripariis~~

~~Similis Faiquinia armillari ut facile eadem dicesse differet
modo caule arborescens altissimo nec humili, ramis ad se
vix illis nodosis. Rami tetragonis subangulati~~

~~Folii uncinati coronati bipollicari glaberrimi
arenarii & nervi subtus pallida emarginata. Racemi
ex apice ramorum ⁱⁿ pediculis ⁱⁿ fasciculis angulati apice
tam vespis crassiores~~

isphaemoides P. panicula erecta contracta, calycibus bifloris polygamis acutis
Culmo simpliciter, foliis distichis rigidis Reti. obs. p. 17.

Habitat in udis et ad margines stagnorum Malabar

Culmi erecti pedales sp. simpliciter. Folia inferiora opposita
superiora alterna primum convoluta deinde patula, rigida
superiora pilosa tuberculata, serrata 3 pollicaria pallide ~~rubra~~
albido viridia uti totum gramen ^{appulsi} margine tomentoso
per rami panicula longi flexuosi ^{appulsi} flouescent. flori plerum
quo solitarii interdum gemini altero pedicellato. Salsula sciss
fiora breviflora truncata.

affinis colorato ei que proxima

convolutam. panicula patens ~~capitulis~~ bifloris, culmo ramoso foliis con-
volutis valvula accipitrona membranacea truncata

Habitat in Arabia, India orientali, China.

Culmus erectus bipedalis ramosus. usque ad paniculam vaginis foliolorum
lucis ^{geniculi fusis punctis} vagina amplo glabro striato apice utrinque macula fla-
vescens, margine ciliata. Folia in pollicina convoluta ^{inferius} interdum
opposita, parva vel albida, patentia glabra. Panicula
terminalis erecta palmari vel ultra. ramis patentibus
haud multum ramulosis ramis filiformibus scabris glabris
florescens ^{longius pedicellatis, pallide rufis}
affine panicula colorata et ⁴phemide

numidicum Pan. panicula ramis subsimplicibus racemosis, glu-
mi ovato-acuti lobis pedicellis coloratis La Mart. T. Enci:
pag. 172.
Habitat in numida

~~as Panicum coloratum~~

colordum Pan. panicula patenta, flaminibus pestilisque coloratis, culmo
ramoso Lin. Mant. 30, Gaicq. Meji. v. 2. p. 365.

Habitat in Barbana Egypto

Culmi pedales erecti inferne ramosi vaginati albi. Soca
con soluta subulata inferne approximata rigida subpergenta
patentia pallida vincta fructa uti vix ad spem punci
tis minutis elevatis inter frons villosa. folia duo in culmo
erectio va remota panicula tenuis floruli oblongi
glabri ^{calycibus} ^{subtiliori} ^{et} ^{per} ^{fertile} ^{alter}
nervoso ~~flor~~ ^{floruli} ^{per} ^{fertile} ^{alter}
culm albor luxior folii plani longioribus panicula major
caetior valvula unispora ovata

sternatum Pan. panicula oblonga glumis marginibus glabris
Dibus pulchre striatis La Mart. T. Enc. pag. 172
Habitat in Carolina.

non vid.

mit Dum Pan. panicula ramosa subvolacea glomeri obliqui fructi depressi
ei semine nitido La Mart. V. Enc. pag. 172.
Habitat in Carolina

non vid.

Love Panicum panicula taxa subflacida glumi oblongi corba
foliorum vagini glabris La Marti T. Enc. pag. 172.
~~Habitat in Domingo~~
β glumi obscure striatis
Habitat in Domingo Guspula Francis β. Ex India

Panicum guineense guinea grm. polygamum Swartz

non vidi

Caffronum Pan. panicula patente, glumis bifloris glabris, vaginis pilosis
Habitat in Cap. b. p. ^{b. p.}
Culmus erectus bipedalis ~~inferne~~ ramosis inferne ^{inferne} vaginis foliorum
folia graminea pilosissima profertur ^{marginibus}. Vaginae pilosae
sternatae. Panicula parva. Flosculi magni salubritate ^{solitane vel gemini} alyxina
rindi purpurascens aut glabra nervosa. fo. pedicelli unguicula
rei capillares rami panicula angulati villosi.

an *Dactyloctenium* Thunb. *Dactyloctenium* a Pan. *Caffronum* Retzi

Yacquinia

Yacquinia arborea foliis cuneiformibus, ramis ad ramificationibus aequalibus: in
senioribus verticillato quaternis: superioribus dichotomis

Habitat in Montserrat

Rami tetragoni, glabri, cicutriati.

Folia in petiolum brevissimum attenuata, bipolliciana, concava, glaberrima,
supra laevissima arenata, subtus pallida et obscure venosa, punctata,
emarginata cum acumine brevissimo

Racemi terminales, interdum gemini, bipollicares et ultra, Flores circa
ter quindecim in singulo. Pedunculi angulati. Pedicelli unguiculati
alterni, sursum sensim incrassati. Flores in Yacqu. armellari

A Yacquinia armellari differt tantummodo altitudine et praecipue nodis
nullis ad eandem ramum, coeternum ^{in omnibus omnibus omnino} ~~similiter~~ ~~et~~ ~~specie~~
~~in eadem~~ Cum protenus igitur discrimina separantur differunt,
nunquam eas ~~esse~~ ~~specie~~ ~~diversas~~ sed potius varietatis vel varietatis
loci vel aetatis esse credidissim. Teste

Desudum Pan. panicula patente floribus solitatis, glumis glabris apice purpureis
Thunb. prod. pag. 19
Habitat in Cap. b. p. i.

6

non vid.

miliaceum ^P panicula laxa flaccida, foliorum vaginis hirs, glumis mucoso
natis nervosis. Linn. h. v.

Melium semine lacteo et albo

B — — — semine nigro Bauh. pin. 26 Theatr. 502.

Habitat in India

Pulvis eris sesqui pedali et ultra crassitie penno asperno vagina
sub. folia usque lata, raris pilos nervosa striata, q^d lora margini
scabra superiora culmo altior, exilis patentia vagina inter strias
elevato punctata pili puncti piliferi. Panicula spithamea pedun-
culi glabri valvula exterior ovata membranacea. antenae ovata
acuminata nervosa glabra
simili miliaceo et omnia robustiora

brachyca
~~var. nigra~~

hil. brachyca

Pan. panicula laxa, ramis inferioribus subtricollatis
foliis ~~lanceolatis~~ ^{vegetis} ~~vegetis~~ vaginisque glabris caule ramisque ~~remantibus~~ ^{fructibus}

Habitat in Brasilia Thonin

Gramen sesquipedale erectum crassiusculis empositis. folia
2-3 pollicaria margine scabra ore velopula. Panicula elongata
pedunculata uti glomerata remota recta. valvula calcarina ter
ovata glauca infescenti leve

effusus P. panicula diffusa, flosculis remotis glumis bifloris vaginis punctis
in geminis corbis
Habitat in Ind. occident. an?

Culmi bipedales erecti crassitie pennae columbinae. vagina glabra
strigata superne inter strias puncta elevata. folia plana longe
glabra margine scabra. plae Panicula pedalis diffusa. flosculi
gemini pedicellati ramosi glaucis virentibus oblongi valvula
calycino ter exterior ovata 2^o nervosa.

Caniculata L. caule fruticoso sabauileato ^{integro nervibus} foliis caniculatis basi revolutis racemis
terminalibus lateralibusque ^{caulis} ~~placulis~~ ^{parvis} ~~linearibus~~
habitat in India occidentali 5

Habitu L. ignei et forte non nisi varietas. Rami teretes superne villosi
aculei paucissimi aciculares purpurascens. Folia juniora utraque
pilis stellatis leviter ~~se~~ villosa adultiora fere glabra, b. f. tripolliciana
reticulata. Racemi terminalis et laterales sequipollicares calyx
minutus. Corolla parva per quinque partita laciniis angustissimis. Hac et
magnitudine femini Conardii.

notatum Pan. panicula patente: axillis notatis, foliis canaliculatis ciliatis

Petz. obs. 4. p. 18.

Habitat in Sumatra

non vidi

pubulatus

acuminatus P. panicula filiformi composita florepulis lamiscolatis acu
minatis attenuatis, vaginis punctatis pilosis

Habitat in India

folia angusta conduplicata. florepuli nervosi acuti glaucis
raspenti fusi valvula acesporea ovata acuminata

glutinosa
polygamum

panicula composita patente, ramis flexuosis, spiculis
pedicellatis distantibus glutinosi, Culmo erecto simplici, foliis

Catambus Sw. prodr. pag. 24
gramen miliaicum Sylvaticum maximum femine albo

Ploar hist. 1. 114. v. 71. fig. 3.

Habitat in Jamaica.

Definitio
toto
ad Pan.
obliquum peduncul
basi altero latere
Cathora et rotund
Data

Culmus erectus, glaber tenuis lavis nitidus vaginis foliorum vestitus. Folia
pessilia pollicem lata attenuata sive polliciana nervosa, tenuissime
fenestrata lora margini vix vix fiabra collo villosa. Vaginae stri-
atae glabrae Panicula subterminali, peduncululi inferiores remotiores
summi breves ramulati subulsi flores globosi subulula externa
nava striata apice villosa
an haec eadem ac Swartzii? nimis affinis videntur obzardis unguis
Catholum arundinaceum glutinosum

obliquum P. folus ovato lanceolatis basi obliquis nervosis. Panicula
pyramidata stricta flosculis globosis

Habitat in India occidentali

Culmibus tres vaginatis vaginis ad collum tomentosis. ^{rigidis} folia ultra
pollicem late attenuata uti vaginis glabris, 4-5 pollicianis
nervi paniculae breves angulati inferiores remotiores superiores contra
tiores

^{adun}
Canigraefum P. panicula patens, foliis glaucis subglobosis, foliis lato
lanceolatis scabri vaginis villosis, culmo erecto
Habitat in India occidentali

an lanatum Swartz

Culmi crassitie penno asperino simplicis, Vaginis vaginis foliorum
stratis, glabri. Folia pithamea, ^{base equalia} ultra pollicem lata, multi
nervia strata, retrospum strata, glabra. Vaginis bi-³ tripollicariis
se molliter villosis propestris marginis. Panicula semipedali
pedunculis ^{subscissis} flectuosis. glauculis gemini remoti pedicellati pedicel-
lus alter longior, magni. Valvula accessoria ovata strata obliqua
marginis submentosa, duo ovato-subrotunda nervosa nitida

Canigraefum lanatum panicula composita erecta glabra, spiculis ovatis
culmo ramoso, foliis ovato-lanceolatis pubescentibus, vaginis lanatis
Sw. prodr. pag. 24.

Dipsermum Pan panicula composita capillari patente, seminibus
geminis, foliis ascendenteis glaberrimis La Mart. - T. Linc. pag 173
Habitat in Amer: meridionali

non vidi

Cayenne *Panicum panicula* oblonga patente summi divaricatis & agnisi
respirati, culmo ramoso La Mark T. Enci: pag. 173.
Habitata Cayenna

non vid.

Capillare

Trichodes P panicula ramosissima patente, rami ramulique subdivisi
capillatibus, culmo declinato, geniculato, foliis ovato-lanceo-
latis glaberrimis Sw. prodr pag. 24.
Gromen mellei cum viride, foliis ciliatis brevibus, panicula capilla-
ta Journal de Bot. par. 1. p. 86. 115. t. 72. fig. 3.

Habitat in Jamaica, Hispaniola.

Culmus tenuis ramosus geniculatus geniculis fuscis, folia ^{sub}ova-
to lanceolata sesquipedalium glabra margine scabra. vagina
glabra striata margine ciliato. Panicula diffusa flospice
parvi nervosi alteri gibbosi cymbiformis
saede affinis brevifolia ut limitem ut invenitur.

capilla et cum Lu Mont. fide herb. Mont.

amudonacium P panicula coarctata patente, ramis ramulisque fructibus
capellanibus, spiculis subtendis, culmo subdiviso geniculato,
foliis lato lanceolatis acuminatis rigidis Sw. prodr. pag. 227.
Habitat in Jamaica.

non vid.

polygamum
Amaranthaceae P. panicula ^{capellan} composita pendent, flosculis polygamis,
Culmo articulis hirsutis Sw. prodr. pag. 24.

~~Holcus~~ 2. Brown. Jan. 366.

Habitat in India occidentali.

non vidi.

Coarctation
suadentia

Dactyloides P. Panicula coarctata, elongata, flosculis subulatis secundis
^{culmo erecto}

Habitat in India orientali Hongk.

Facies: ~~Dact. glomerati modo parva~~ ^{arundinis vel agrostidis} Caulmus bipedalis erectus; glaber,
stratus ter, simplex, geniculis fuscis. Folia plana longa, semi
pollicem lata, margine scabra. Ligula brevis membranacea,
^{cresta} ~~margine rugoso margine fuscata~~. Panicula pedalis primum con
^{tricta demum ramis nonnullis parum patentibus inferioribus et}
^{ramis} remotioribus. Rachidi filiformes scabri, glabri ramuli brevissimi secundis
5-6 floris. Valvula glabra strata, lanceolata attenuata, acies, spon
parva ovata. Corolla longitudine calycis angusta

agrostoides P.

inter amer; quere. Desfont

~~Polygonum P~~

~~Panicum polygonum fructu remotis, glomerulis periclysis racheo. Desch. obs. Polygonum
Retz. obs. 4. p. 14.~~

antidotale R. panicula nutante, calycibus bifloris polygamis acutis, culms erecto
angustissimo ramosissimo Retz. off. 4. p. 17. 4
Habitat in Hortis malabarorum

manus ramosum altum ramosissimum culmo & tereti ^{inspice per penam antennis} folia longissima pallida
virentia. Panicula sesquipedalis valvula nervosa

miliare la Marti fide herb la Marti
descript. vna

miliare affine

fleunosum Pan. panicula capillari patente, pedunculis flexuosis, calycibus
ovatis, geniculorum barba reflexa Retz. obs. 3. p. 9.
Habitat

non vid.

Capillare ρ panicula capillari erecta patente, foliorum vaginis hirsutis

in S. V.

Milium indicum, panicula sparsa erecta Tournef. F. R. H. 515.

Panicum miliaecum v. inde, foliis latis barbuis, panicula capillacea Swan herbar.
1. p. 115. T. 72. fig. 3.

Habitat in Virginia, Jamaica \odot

Culmus erectus sesquipedalis vaginis foliorum ^{articulis superioribus villosis} setis folia longa
plana nervosa ^{marginata} punctata ^{marginata pubescentibus nervis} glabra. Vagina striata confertim
elevato punctata punctis filiformibus. Panicula pedalis et ultra
ramosissima diffusa ramis capillanibus pubescentibus. flosculi
parvi oblongi glabri aculi valvulae lobis exterioribus ovatis interioribus
oblongis

capillare Pan. panicula capillari superne expansa, glumis acumina-
tis, vaginis hirs. La Mart. T. En. p. 173.
Habitat in Virginia

~~Trichodes Sw. fide herb. Rich.~~

capillare La Mart. Icon 82. fig. 3. glumis obtusis & stragulis
est Trichodes Sw. fide herb. Rich.

~~Trichodes Sw.~~

virgatum Panicum panicula virgata, glumis acuminatis lobus extrema dexte
parte L. L. V. pag. 107.

Panicum paniculatum, glumis acutis Gron. Voy. 133.

Habitat in Virginia
gramen altissimum panicula diffusissima

non vidi

turgidum P. paniculae spiculis obesi triflori polygami glomerum vaginis longis
2 D. uterque ventricosi, caule frutescente Forst. fl. arab. pag. 100.

mibi non notum

forte inter paniculae b. varietate

tumidum - P. pliculis rugidis culmo repente Kunz Natuf. 23. p. 205
Habitatio India orientali

michi haec notum Dubii credens

~~aspera panicula ramosa summa patentissima, pedunculis asperis
spiculis decemfloris, vaginis foliorum articulis hirsutis 4 ang.
herb. 3-7. 56. L.V. p. 114.~~

palmifolium P. foliis plicatis, culmo erecto Kunz in Nat. Hist. 23. p. 200.
Habitat in India orientali.

non vid.

resicula variegata

Sugars - *P. paniculata* celerima maritima, culmo decumbente König in Naturf.

23. pag. 209.

Habitat in India orientali

non vidi

an inter pan. compof.

maculatum P. fraxino, caule arundinaceo Aubl. fl. Gu. pag.

non rdi

Desmodium

Congifolia Rhexia pilosa, foliis lanceolatis integerrimis petiolatis ^{longiusculis} in nervis; nervis
lateralibus bipartitis, pedunculis axillaribus dichotomis

Habitat in America meridionali Dr. J. Roth.

Herbaceum Caules laevi, debiles, angulati, et pilis densis adpressis cineres-flavescentibus
remota ^{longiusculis} pollicem lata, prostrata
foliis petiolata, opposita, tri vel quadripollicaria, lanceolata, altera
tertia, acuta, integerrima, supra adpressa pilis nonnullis adpressis, subt
rindica, subaequali, pallida, petiolo ^{longiusculo} basi
remota ^{longiusculo} basi, laterales in duos ramos parallelos divisi

Longi tudine petioli
particuli longitudine
pediculi oppositi
terminaliter
pedunculum unversum
aequantis, pediculi fere
uniusculares

Petiolus petiolus pollicaris pilis et pilis pilosis cineres flavescens
Pedunculi axillares, bis bifidi, cum flore pedicellato in singula dichoto
mia, pilosi, cinerei. Colla opposita ad primam dichotomiam.

Calycis oblongi, gibbo hinc; laciniis quaeque, fibrillatis, intus glabris
coloratis

Petala oblonga subobovata, calyce dimidio longiora
capsula subglobosa apice hinc calyce breviori
Preseptatula quinque lunata superne libera hinc

a Rhex. herb. hispida Act. de la Soc. Phil. Natur. de Paris.

Tom. 1. pag. 160. Differt foliis petiolatis, pedunculis trifloris

no 15 Dr. J. Roth

manisatum P. panicula patente, floribus solitariis mucronatis, culmo radicante
adspicente Retz. prod. obj. 4. p. 10
Habitat in India orientali

~~Canatum panicula composita erecta glabra, spiculis ovatis, culmo
ramoso, foliis ovato-lanceolatis pubescentibus, vaginis lanuginoso
sorbis Sw. prodr. pag. 24.
Habitu in Jamaica~~

non nisi eadem ac lanuginosum nostrum

foliatum P. folii cordati Aubl. fl. Gu. pag.
miliun, cordifolium folii cordati Plum. Catal.
Habitat in Guiana

non vidi

geniculatum P.

Tenax Panicum macrophyllum racemo magno sepli composito
multifloro, aristis longis tenacissimis flavescientibus. Act.
Soc. hist. n. Par. 1. pag. 106.
Habitat in Amer. merid.