

# Pastinaca

1. lucida P. foliis radicalibus cordatis lobatis, caulinis ternatis quinatisque, ramis obovatis simplicibus cordatis
2. opoponax P. foliis <sup>bi</sup> pinnatis: foliolis <sup>ovatis</sup> cordatis oblongis basi antica brevioribus, petiolis ~~fragilibus~~ serratis petiolisque glabris
4. fabiva P. foliis pinnatis: foliolis <sup>radicalibus plerumque</sup> individualibus ovatis in ipso summo crenatis
3. thuronium P. foliis pinnatis, foliolis cordatis basi antica crenatis petiolis hispida

*Pastinaca pimpinellifolia* foliis pennatis, foliolis inciso serratis, inferioribus  
subrotundis, superioribus oblongis Mart. in Schrad. Journ. f. d. b. 16. pag. 458.

*Tordylium umbelliferum* remotis, foliis duplicato-pennatis, pennis inciso tomentosis  
Müll. Fr. ed. germ. tab. 26.

*Pastinaca orientalis* foliis eleganter incisio Tournef. Corp. 23. Buxb. Cent.  
3. p. 17. tab. 27.

Habitat in germanis circa Buxb. Bulath 4



lucida P. foliis simplicibus cordatis lobatis lucidis acute crenatis Lin' Manx  
5. Pl. + L. V. p. 190 - Jacq. hort. 2. tab. 199.

Pastinaca folio quasi libanotidis latifolia Boerh. huyob. 1. p. 67.

Habitat in Europa australi & angulatis

Caulis sesquipedalis fide atrox ramosus stridus altitu  
dine fere hominis. Rami numerosi paniculati. Folia radi  
calia cordata vix lobata, ~~reticulata~~ ~~frabra~~ venosa

caulina prostrata ternata foliolis lateralibus sesquipedibus  
oblique ovatis impari petiolato cordato trilobo quinque  
terminale loboque secunda quinato pennata summa simplicibus  
maximam lobato ovata, lobata. Ramea omnia ovata omnia  
acute crenata umbella numerosissima <sup>ramulorum</sup> terminales  
et axillares ramulorum. Semen orbiculatum

lucida P. foliis radicalibus cordatis lobatis, caulinis ternatis  
quinatisque, rameis obovatis Jouan ill. p. 19. tab. 11. 12.

opoponax P. foliis bipinnatis folioli ovato cordati serrati  
petiolis que glabris

Panax castaneum Bauh pin. 156

Panax heracleum Monf. hist. 3. p. 315. f. 9. v. 17. fig. 2.

Habitat in Italia Sicilia

Caulis altus fistulosus subtilissime striatus, ramis

Folia petiolata bipinnata. Foliola petiolata

Foliola 5 que penna tripartita extrema simplex reliqua petiolata  
super foliola inferiora 5 penna bifida in penna interiora <sup>in penna interiora</sup> ternata exteriora  
superiora 5 penna bifida in penna interiora ternata exteriora  
ultima ternata bitoba. Omnia ovata latere exteriori breviter  
rotata utraque acuta grosse serrata terminali  
Petioli tripartiti <sup>in</sup> subtus convexi, articulati organici  
cili crispantibus

Rami bipinnati floriferi oppositi folium utrumque  
pinnatum supremum simplex  
umbrella ampla in summo utraque amplexata  
ta floris plures abortientes



planta a Lasepitiis Chironio Volo cælo diversa quia ut  
 hanc jam monuit ad Lasep. Chiron. neq. synon. ad Lasep.  
 opop. allata cum Lasep. Chiron. conveniant. Differt  
 etiam a Ch. umbellatis vel ramis umbelliferis non  
 verticillatis compositione foliorum quo bipinnata  
 ut monitum habet differentia Lin. cum in Chironio  
 foliola ternaria pennata sequa foliorum hirsu-  
 tum differt de eoque species omnino ~~comparanda~~

Lichen digitis  
 vel potius  
 pastinacis  
 germanicis  
 affinis Panax  
 live Ephedra  
 costis flore  
 caute Krauch  
 hist. 3. p. 2. p. 156

~~Chironio~~ foliola etiam nonnulla sub um-  
 bellis et umbellulis adsunt in Chironio quo  
 plane desunt in ~~Oppoponace~~. Gouerni opop.  
 esse hanc esse Chironi. neq. ~~Oppoponace~~ con-  
 agnovit et speciem hanc ad eum misit  
 opp. neque hanc habitare neq. ~~Moy~~  
 pelet.

erronee yttur Magnoliam false Panax costi-  
 num cum Panace perquam dod. conjungit  
 et Bauhini h. figura ab epi diversa patetur

~~Chrysanthemum~~  
~~maum~~

*Lappaceum* foliis oblique cordatis petiolis hirsutis Len Spec. p. 350 \* S. V. p. 202.

*Sagapenum eximatum* Ges. hor.

*Panae es peregrinum* Dod. pempt. 309.

*Panax pastinaca* folio Bauh. Pin. 156

*Panax heracleum* Monf. h. 3. p. 315 / 2. t. 17. fig. 1.

Habitat Montpelii

Folia integra, cordata, lobo superiore brevior, crenata, obtusa. Petioli valde hirsuti. ~~4~~ Involucrum parvum, tam universale quam partiale; hinc diversa a *pastinaca* Costina Len Sp. 1. 1.

*Pastinaca oppositifolia* Linn. ~~alt. Gouan~~ ~~alt. Gouan~~ <sup>len</sup> *alt. Pastinaca* Rhomus cadens



Cheroneum E

~~Opoponax~~

P. foliis pinnatis: foliolis basi antica excisis <sup>cordatis</sup> ~~len~~ <sup>acutis petioli</sup> ~~len~~ <sup>strigosi</sup> ~~len~~ <sup>hirsuti</sup>

~~Panax rostratum~~ Bauh. pin. 156.

~~Panax Heracleum~~ Moench. hist. 3. p. 15. f. 9. t. 17. fig. 2.

Habitat in Italia, Sicilia

Caulis altitudine hominis ramosis crispis digitifloris  
basi testusquamis membranaceis scariosis lene nitens  
folia pinnata simplicissima cordata acute crenata  
demum ternata quinquata folia infima caulis bipin-

nae perinata foliolis <sup>cordatis</sup> ovatis oblongis crenatis venosis septem  
num dod. perspic. excepto terminali obtusi altero tal laterali  
309 altero latere breviora, tripollicana, ad basin

~~Panax pastinacae~~ folia simplicia petioli omnes strigosi uti vagina  
foliorum Bauh. ped rami floriferi inferiores solitarii demum oppositi  
pin 156. si demum verticillati folium lanceolatum septe

~~Panax Heracleum~~ ad basi ramorum, rami iterum verticillato rami  
Moench. hist. 3. foli. Semina omnino Pastinacia  
p. 15. f. 9. t. 17. fig. 1.

~~Laserpitium Cheroneum~~ huius refert Juan & David  
rapite & videtur  
Laserpitium Cheroneum len foliolis oblique cordatis  
petioli hirsuti len. hist. pl. pag. 368

Sativa P. foliis simpliciter pennatis Lin. S. V.

Pastinaca sylvestris latifolia Bauh. pin. 155.

β Pastinaca sativa latifolia Bauh. pin. 155.

Habitat in Europa in fronsionibus moderatis et pasuis.

Caulis sex pedalis. <sup>infrum per utroque pinnarum</sup> Pinnas pinnules oblique ovatae, basi cartilagineae  
rimo perenne ~~perenne~~ crispae.

vetenus Describenda et differentia enedatur



Puff-Beech Bark