

Pinoa

nitida Pansa foliis oblongis congestis nitidis, racemis terminalibus
Lour. fl. Cochinch. pag 172.

Habitat in Cochinchina 5

Caulis fruticosus 4 pedalis erectus ramosis. folia ovata-oblonga
integerrima. plana margine reflexa. Cal. 2 phyllis. —
Corolla alba campanulata. Stylus lateralis Capsula ovata

non vidi

an huius generis

Scandens Pensa foliis cordatis, pedunculis bifloris, caule scandente. Lour
fl. Cochinch. pag 73.

Habitata Cochinchina &

Caulis fruticosus ramosis. Folia lata acuminata, integerrima.
Pedunculi longi axillares. Corolla lobis campanulata plicata.
Cappula 4-lobulata. Calyx magnus siccus persistens.

non vid. an hujus generis

1 Jarcocolla ^{Thombeo} foliis ovatis planis, calycibus ciliatis, foliis majoribus Lin. Spec. pl. p. 162 +
Thymali Myrsinitis specie abusive Othoneica, subrotundis foliis,
Stoechadis arabica squamato capitulo duobus lacrymam pendens Plukn
mant. 103. t. 446. fig. 6

Habitat in Othoneica

romuli Tetragoni

Rami tuberculati ^{avenia} ^{glabra} mucati a capite foliorum. Folia ut et acuta
patentia ^{avenia} ^{glabra} minima nec quadrifariam imbricata ut in squa
mose. Flores terminalis brevissime pedunculatis. Limbi lacini
ovato obtusifido basi ^{veo} linearis colorata

corollis obtusis laciniis limbi ovatis obtusi
bracteis linearibus

mutatur differentia
differentia Jarcocolla pro Syst. Veg. potius ad Squammosam spectat

2 *glauca* P foliis rhombo-oratis, bracteis cuneatis coloratis Lin. Mant. 199. t. V. p. 1524
Pennisia sanicollis foliis oppositis rhombicis acutis planis subpetiolaribus patentibus, foliolis calycinis lineari-bus Berg Cap. p. 35
Habitat in Cap. b. plerq;

Indica ramopisum. Rami superne a lineis decurrentibus tetragonibus
Folia bracteata, ^{subrhombica} acuta, ^{apiculata} patula, ~~subrhombica~~ ^{obovata} ^{obovata} lobata. Flores
fasciculati, terminati, purpurei. Bracteis foliis minoribus,
cuneatis, acutis purpureis. Similis *sanicolla* flores duplo
minori

Longiflora Pen. folus rhomboides acutis, floribus quadrifidis sub Longiflora
Meebury. T. 51. fig. 2.
Habitat in C. 6. p. 1.

non vidi

Cneorum Pansa. foliis linearibus lanceolatis serratis, floribus congestis
caputatis, bracteis foliis minoribus La Mart. T. En. pag. 317
Pansa cneorum Meerb. tab. 51. fig. 2.
Habitat in C. b. p. 5

non vidi

3 Squamosa P foliis rhombis cuneiformibus carnos. ^{Folius flegalibus ciliatis glutinos}
Habitata ^{Lin. Spei: pl. p. 162 *} ~~Obsequia~~

~~Pensa~~ tetragona foliis rhombis imbricatis patentibus, tubis florum
longis, foliis laeycians obovatis ^{Benjia Pl. Cap. p. 36.}
Habitata C. b. ^{Spei.}

Rami obscure tetragoni, mucicatis, ^{scabi. conf. e foliis deciduis} ~~latiniatis~~. Folia sessilia qua
glandula ^{fulva} ~~est~~ ⁱⁿ ~~angulo~~ ^{supra} ~~folium~~ ^{quadrifida} ~~imbricata~~, rhombea, glandulosa, conica are
na int. egeronia. ^{suprema} ~~inferiora~~ ^{majori} ~~marginem~~
cilia ^{setuico} ~~ciliata~~, glutinosa. ^{colorata} ~~corolla~~ ⁵ ~~lobis~~ ^{terminalibus}
aggregati bracteis longioribus ^{lobis} ~~lacinis~~ ^{ovata} ~~acuta~~
obtusis

tomentosa Penna florum lateralibus, foliis ovatis tomentosis truncato-
pinnatis. Cap.

pag. 30.

Habitat in Cap. b. p. i.

non vidi

4 mucronata P foliis cordatis acuminatis Len. S. V. p. 154

Encia africana, unedoni flore amplo, foliis cordiformibus in acumen
desinentibus Raj-dendr. 97.

Habitata in Othiopia ^{verticillati tres vel quatuor}
rami e capi foliorum scabri. ^{semin. pucillana} folia parva congestissima patentis
forma superne ^{adnatis} subreflexa basi ^{separ. adnatis} connata in alio remotum
vix retiolata. Flores ad apices ramorum axillares subsusp.
les foliis parum longiores laciniis ovatis acutis pilis

Vaccinia africana foliis decussata cordiformibus Petrus. 692. herb
mus Vaillant

5 Fruticulosa foliis oblongiusculis obtusis, bracteis orbiculatis acutis ^{Len. Hypt. p. 121.}
L. V. p. 154

Habitat in C. b. Speitz sub

Fruticulus ramifus, dichotomus ^{ramis} ~~ramis~~ ^{superne tetragonis} nudatis. Folia
opposita subpetiolata ovalia, ^{glabra} ~~conspicua~~ ^{remota}
Folia floralia parva rhombica orbiculata, acuta. Flores
in apicibus ramorum, subsolitarij, ~~bracteisque~~ ^{bracteisque} pedunculati
acutiusculi. Bractea nulla

6 myrtoides P foliis lanceolatis Lin. Suppl. p. 122. * S. V. p. 154

Habitat in Q. b. spei 7

Rami teretes, superne obscure tetragoni. folia opposita, sessilia, confer-
ta, ovata, subtus vix nervosa & locis terminalibus, subfoliariis
cincti bracteis viridibus, acutis, sub quibus folia duo ^{naevy} ~~ovata~~ ^{subulata}
la parva ad basin floris, calyx acutus viridis.

varietatem foliis subulatis juniperi subtus carinata. floris oppositi
breve pedicellatis in ultimis axillis. laciniis corollae acuti.

7 Cateniflora P foliis ovatis floribus sessilibus lateralibus Lin. Suppl. pag. 122. * L. V. p. 154

Habitat in C. b. Speitz

Rami rubentes saepe terni fastigiati verti foliis ovatis
globis conaevis obtusis, sessilibus interstitiis longiora. Flores
axillares in ramis ultimis, ultimis foliis parum longioris
Corolla laciniis lanceolatis iniquetro latere interiori
cannato canna basi incip.

laciniis corollae lanceolatis

S. marginata ^{sessilibus} foliis cordatis marginatis, floribus lateralibus *Lin. Mant. 199 + L. V. p. 154*

Habitat in cap. b. p. u.

Totus fere ramus interdum verno glabri vindi flavescens.
folia sessilia opposita, ^{interdum ternis} glabra, nitida, pallide vinda late
gemina margine crassiora, magnitudine basi inferne remotio
ra, superne magis approximata. Flores axillares, sessiles folio
breviores, bini vel oppositi laciniis ovatis acuminatis

acuminatis

— laciniis corollae ovatis