

Perkulana

1. glabra P. foliis oblongis acuminatis glabris
2. tomentosa P. foliis cordatis ramisque tomentosis
3. japonica P. foliis cordatis glabris, umbellis simplicibus laciniis corollatis
4. <sup>purpurascens</sup> cordata P. foliis cordatis glabris acuminatis, laciniis corollae linearibus  
oblongis, umbellis proleferis

glabra Pergularia foliis oblongis acuminatis glabris  
8 Co. pergularis Rumphii amb. 5 p. 51. tab. 29. fig. 2.  
P. glab<sup>7</sup> foliis ovatis glabris Mant. 53. l. V. p. 256.

Habitat in Seylone mesotrope ty

asales Detumer  
V. 25

Solubilis glaber tenuis. Ramis teretes articulati. Folia petiolata oppo-  
sita patentissima oblonga vel elliptica, acuminata, interspersa veno-  
sa. Petioli basi ad caulem ciliati. Panicula panum supra axillis foliorum  
longitudine folii foliola lanceolata glabra ad basin pedicellorum. Calyx  
glaber foliolis ovatis. Corolla glabra tubo calycis panum longior laciniis  
limbi rotundati.

Corolla facie Pergularia sed nectarium nullum antheris cum sty-  
mate <sup>non</sup> ovato acuto connatis, sagittato filamenta brevissima stylus uni-  
cus deoque forte alius generis

2<sup>3</sup> purpurea Perularia Perularia foliis cordatis glabris, acuminatis, laciniis corollae linea  
n-oblongis, umbellis proliferis  
Aflicta foliis cordatis acuminatis, floribus terminalibus Burm.  
Ind. pag. 72. Tab. 27. fig. 2.

Habitat in India orientali, China. Desit Dr. Burmann  
Volubilis Rami <sup>tenues</sup> filiformes, cinerei, ad lentem villosi.  
Solia opposita, petiolata, glabra, nervosa, venosa, bypollia anis,  
superiora minora.

Petiolus longitudine foliorum.  
Umbella caudales, pedunculata, foliis brevior. Pedunculus petio-  
lo crassior, rigidus. Pedicelli filiformes, unguiculares, pubescentes  
uti Calycis.

Corolla infundibuliformis, unguiculares. Tauro villosa, interna  
cum etiam limbis ~~aque~~ parum villosa.

Affinis Japanica Thunb. et  
Umbella prolifera nec simplicis ut in P. japonica, laciniis lim-  
bi lineares nec ovata, an satis distincta?

Perularia odoratissima foliis cordatis glabriusculis meentibus  
genitalibusque corolla brevioribus, fignate conis obtu-  
to Smith &c. prosp. 16. t. 16  
Cynanch. odoratissimum Loewius Cochin  
166

Scabra Matth. ~~Lin.~~ L. V. p. 210. Spec. pl. p. 1161.

Matthiola folio aspero subrotundo, fructu nigricante Plum. gen

16. ii. 1773. fig. 2.

rugosa G. foliis subcordatis ovatis acuto subtus tomentosis superne glabris, floribus  
Habitata in America <sup>in hexandris his prodr. pag 59</sup> <sup>in</sup> <sup>triacate</sup>

Rami teretes oppositi inferne glabri, superne vellosi. Solia pe-

tiolata, versus apicem ramulorum congesti, opposita, ovato-oblonga

interdum versus basin angustiora, quicquid parvo ter minata

in breviora, supra <sup>apicem</sup> ~~habent~~ <sup>subtus</sup> ~~habent~~ <sup>minuti elevatis</sup> ~~subtus~~ <sup>minuti elevatis</sup>

venoso reticulato rugosa, vellis minutis finis adpressa, bis-  
tripolliciana. ~~Stipula~~ ~~subulata~~ ~~decidua~~.

Peduncululi axillares oppositi compressi, vellosi, perquam pini

versus apicem dilatati bis vel ter dichotomi. Flores interdum

in capitulo subsessiles. Flores in singula dichotomia uti reliqui

sessiles bractea subulata decidua ad stamificationes

et flores calyx unicolatus oblique dentatus. Tubus floris

sesqui polliciani velloso subsericeo molli. ~~Tractus~~

magnitudine nasi vellosus ~~sericeo~~ <sup>sericeo</sup> ~~sericeo~~ <sup>sericeo</sup> ~~sericeo~~ <sup>sericeo</sup>

Japonica P. foliis cordatis glabris Thunb. fl. jap. p. 1114 N. p. 256

Habitat in Japonia China

Caulis volubili, teres, glaber, simplex. Folia petiolata, opposita, ~~alt~~  
~~acuminata~~ <sup>acuta</sup> integra, nervosa, ~~venis~~ ~~fere~~ ~~bipollicari~~ ~~ca.~~  
~~pollicaria~~ Petioli unguiculati. Umbellae laterales <sup>simpliciter</sup> ~~per~~ ~~interdum~~  
~~proliferi~~ pedunculati, pedunculus petio duplo ~~causosus~~ ~~ca.~~  
caulis calycis lacinae ~~hirs.~~ limbi lacinae ovatae <sup>obtusae</sup> ~~inter~~  
villosae.

umbellis simplicibus, lacinae corollae ovatae

inter villosae

tomentosa P. foliis cordatis tomentosi Lin-Mant. 53. \* L.V. p. 256.

*Repleta cordata, foliis petiolatis cordatis reniformibus, caule ramis  
apice volubili. Forst. 2. sept. p. 49.*

Habitat in Arabia

Rami adtermi volubiles villosi tomentosi uti tota planta

Folia <sup>lanuginosa</sup> petiolata opposita ~~in cord~~ unguiculata, crassifolia  
aerea cerasia latitudine et longitudine aequalia, suprema  
minora attenuata. umbella laterales simplicis. folliculi  
duo ovati attenuati utriusque mucositate.

*Edulis* *Pergularia* foliis ovatis acuminatis glabris, caule herbaceo  
*Pergularia* *Edulis* foliis ovatis integris glabris, caule herbaceo  
volubili Thunb. prodr. fl. cap. pag. 30.  
Habitat in Cap. b. pei.

*Senecio Perulani* foliis ovato oblongis, pedunculis multifloris, caule  
suffruticoso volubili Lour. fl. Cochinch. pag 169.

Habitat in China

Caules teres, teres. Folia obtusa, integerrima, glabra. Pedunculi  
terminales et laterales. Corolla lutea laciniis sive laciniis  
folliculis oblongis

non vidi



Dwanicata *Pergularia* foliis lato lanceolatis undulatis, caule cespitoso, pedunculis  
cis bifloris Dwanicatis Lour. fl. Choct. pag. 167.

Habitat in China.

Frutex pubescens; ramis multis herbis. Folia integerrima, incurva  
glabra, opposita. Flores lutei, ~~per~~ rubra adpressi. Calycis foliola sub  
lato. Corolla subhypocrateriformis tubo crasso. folliculi uti oblongi  
monogyra uti omnes *Pergularia*

*P. dworakata* Lawrence  
*prosp.*