

Pharus

1 latifolius P.

latifolius Pharus An. I. V. pag. 255.

In mont. ferat legit Dr. Ryan.

Culmus erectus, bipedalis, simplex, brachiis rectis vaginis gemi-
nis longis striatis, diphyllis, brachiis: articulis inferioribus tri-
pollicariis.

Folia radicalia 3. vel 4. culmi 2. petiolata, spithamea, lanicola
ob-elliptica, pollicari brevia lata, integerrima, acuminata.

basiaque acuta, multistriata, subtus fibrinifera.

Petiolica apice
vaginae, cana
lobulata

Panicula terminalis, erecta, spithamea. Pedunculi et
pedicelli compressi, villosi. Flores sessiles, sex alterni; un-
guiculares, steriles scilicet.

Glume calycinae lanicolatae, subaequales, glabrae, fusco-pur-
purascentes.

Corolla valvulae calycis longior, linearis, acuta; exterior
involuta, inferne striata glabra, superne villosa.

ciliatus ^{Phorus} panicula subramosa; calycibus apetalis ciliatis muticis. Retz. off.

s. p. 23. *

Habitat ad margines stagnorum

latifolius Pharus In. V. pag. 059. Arn. Acad. S. p. 409 &

Graemen avenaceum sylvaticum, foliis latissimis locustis
longis non anfractis, glumis pediculis Moen. hist. 1. p. 116. t. 73.
fig. 2.

Pharus foliis nervosis obtusis pagina superiora coelum respicienti
bur Brown. Jan. 344. t. 30. fig. 2.

Habitat in Jamaica

Graemen sesquipedale et ultra Culmus simplex erectus fructus
folia per ~~capitulum~~ vaginata. Folia petiolata glaberrima

frithamou lato lanceolata attenuata integerrima laevia

~~fructus p. r. r. r.~~
Panícula terminalis patens. glomeruli alterni breviter pedun-
culato

Cappulaceus Ph. culmo ramoso, seminibus pilis rigidis ad apicem
hertis Aubl. fl. guian. 2. p. 859
Habitat in Guiana

non vid.

ciliatus *P. panicula* fulvamofo, calycibus apetalis ciliatis mutuis
Retz - obj. s. pag 29
Habitat in India orient.

nisi ignotus

Cappulaceus P. culmo ramoso, femine pelis nigris ad apicem hirs. Cuba.
fl. guan-2-pag-859
Habitata Guayana

non vidi