

Pirramnia

♂ Calyx tripartitus Corolla 3. petala

♀ Calyx et Cor. maris. Bacca biculana

1. antidesma P.

Pseudobrosium La Mothe, luc forte prestat




a



b

*Pirramnia antisejma*





antidesma Piramnia Sw. prod: pag. 27

Berberis fruticosa, racemosa, fraxini folio alato, fructu nigro dipyreno

Roan. hist. 2. p. 101. tab. 200. fig. 2.

Habitat in Montserrat de Ryan 5

Rami teretes, inferne glabri, cortice cineres, punctis elevatis adpressis, superne flavescenti-tovinosi.

Folia petiolata, alterna, semipedalia, plerumque pinnata cum impari rarisime cum pari, unicum tum terminale, tri vel quadrijugis. Folia ter petiolata, opposita vel alterna: interiora ovata vel subcordata bipolliciana, intima minoris: exteriora elliptica, majora, omnia acuminata, obtusiuscula, integerrima, utrinque glabra, venosa, supra nitida, subtus pallidiora.

Stipulae supra planiusculae, oculo armato villosae: partiales breves, crassiusculae, fuscae

Racemus terminalis, compressus, laxus. <sup>puthamous</sup> Pedunculi partiales inferne patentes, alterni, flavido-cinerascentes, uti pedunculus communis compressus: pedicelli <sup>breves</sup> compressi, in arbore floemineo pauciores, aggregati plures simul aggregati, distant fasciculi distantes;

Calyx quinquefidus: laciniis ovatis

Corolla: Petala quinque, linearia, calyce longiora.

Stamina filamenta quinque frequentissime ex observatione S. Ryan. Anthera

didyma

a) flor masculus magnitudine naturali b) auctus



antidesma Pirramnia Sw-prodr. p. 27.

Herbena fruticosa, racemosa, lacinia folio alato, fructu  
nigro depresso Sloan h. 2. p. 108. t. 200. fig. 2.

Habitat in Jamaica, Domingo, <sup>in</sup> Montserrat & Ryan &  
rami superne lentic tomentosi canescenti.

Folia petiolata alterna, pennata cum impari trijuga. foliola petiolata  
alterna, bipollicaria et ultra interiora <sup>acuminata</sup> ~~ovata~~ exteriora oblongo ovata  
utroque acuminata utroque acuta, integerrima, glabra venosa supra nitida  
membranacea.

Petioles quinque pollicares glabri: pariterque superius purpurascens

Racemi terminaliter basi ramosi diffusi: flores sparsi per intervalla  
aggregati parvi breviter pedicellati

Stamina 5 que semper observant de Ryan