

# Pisonia

1. inermis P. caule inermi
2. alternifolia P. foliis petiolatis alternis suboppositisque lanceolato-ellipticis  
- spinis brevibus recurvis
3. aculeata P. spinis axillaribus setigeris fissis, foliis ovato-oblongis  
- <sup>longis</sup> <sup>stipulis</sup>
4. nigricans P. inermis, foliis ovatis acuminatis, floribus cymosis erectis,  
fructibus baccatis
5. subcordata P. foliis cordato-subrotundis, fructibus fixis sublatatis pentagonis,  
angulis apice mucronatis
6. coriacea P. inermis foliis lanceolato-ovatis, pedunculis terminalibus  
laxis, floribus nutantibus, fructibus baccatis

Pisonia uti Boerhaavia gaudet potius calyce infero quam superno  
quod ulterius examinandum Richard

*Lemnitomentosa* Pisona

Habitat in Antillis. herb. Surinami apud Gussone no 770  
Rami dichotomi, teretes, apice villosi. Folia petiolata, opposita, bipollicia  
nerviata, acuminata, utrinque acuta, basi altero latere parum angustiora  
et subbreviora, obsolete simpliciter venosa, subtomentosa. Flores non  
vidi



*Labiifolia* *Pisonia* foliis ovatis subrotundisque

Habitat in insula franco Commerçon herb. Guss.

Rami dichotomi, glabri, ciliati, crassius panno cygneo, cortice cinereo. Folia versus apicem, approximata, 2-4 pollicaria, ovata, oblonga, vel subrotunda, obtusa, saepe emarginata, glabra, remote nervosa, subtus obscure subtilissime venosa et villosula. Corymbus terminalis diomphalis. Calycis tubulati, villosuli.

*Infolcata Pisonia*

Habitat in Hipula Francie Commeson herb. Jusf.

Rami ternati, glabri, angulati. Sola ternata, raro 4<sup>na</sup> petiolata, 3-4 pollicaria,  
oblonga, lata, integerrima, utraque obtusa, glabra, subtus simpliciter venosa. Petio-  
li semipollicares. Pedunculis ternatis, longitudine foliorum: partibus umbellati,  
quinque flores glabri.



*Collina P. inermis* foliis lanceolato-ovatis, pedunculis terminali-  
bus laxis, floribus nutantibus fructibus baccatis Sw. prodr.  
pag. 60.

Habitat in Hispania

nigricans P. inermis, foliis ovatis acuminatis, floribus cymosis erectis,  
fructibus baccatis Sw. prod. pag. 60.

*Pisonia inermis* Gaertn

Habitat in Samaria 5



vermis P caule vermi ten L. V. ~~Quinqu~~ ~~lul~~  
Kata-kata walli Rheid. mal. 7. p. 338. 17.  
Pisonia malabarica non pinofa Ann. m. herb. 502.  
Habitat in India

michi ignot

*Alternanthera* <sup>lanceolata</sup> *Rafin.* foliis petiolatis alternis sub oppositisque ellipticis, super spinis  
brevibus sciuvis

Habitat in India orientali *Frantz*

Rami teretes *caulis* superne villosus ~~terres~~ cinereus. Folia petiolata  
inferne alterna superne opposita *la* subcordata *eu* via nervosa bipolli-  
cana *et* *ultra*, oculo armato subvillosa. *ae* *spina* *varicosi* superne  
saepe nulli inferne axillari *via* *femi* unguiculata. Pedunculi *at* *illa*  
*res* *et* *gemma* *diffusa* foliis breviores *floris* corymbosi



undata Rafida foliis pennatis undulatis, floribus trigynis tetragynisque Lin. S.V. 9034 40  
Rafida decursiva; foliis decursive pennatis; foliis subsqualibus; floribus trigynis  
Dorfl. Catal. pl. aegypt. pag. 66.

aculeata P. spinis axillaribus patentissimis An. 5. V. Gaegu. Am.

Pisonia aculeata, fructu glutinoso et racemoso Plum. gen. 7. v. 227. p. 1.

Rhamnus: Lycium Trogaeo Tamaicaphus dictum Plukn. alm. 318.

tab. 100. fig. 2.

Palium affinis arbor spinosa, flore herbaceo pentapetaloides, fructu sicco  
nudo canaliculato. Lappaceo Sloan. hist. 2. p. 25. T. 167.

Habitat in America meridionali &

Rami teretes ~~superne cindus vellosi cinerei~~. Spinis duo interdum  
foliis in axillis foliorum unguiculari recta subulata, interdum  
~~brevis recurva~~. <sup>opposita vel</sup> ~~ovata~~ basi acuta  
nervosa avenia bypolliana et altera acumine brevi obtus. Peduni  
culi axillares foliis oppositi foliariis interdum gemini in altero  
axilla tomentosi, e gemma <sup>folio</sup> ~~conspicua~~ cinerei reticoli  
fructus oblongus basi parum angustior

recti foliis ovato-oblongo



negrivans *P. inermis* foliis ovatis acuminatis, floribus cymosis erectis, fructibus  
baccatis Sw. prodr. p. 60

*Pisonia inermis* Jacq. hist. 275.

Habitat in Jamaica et Insula St. Crucis Pflug 5

Rami glabri. Folia petiolata, opposita glabra integerrima nervosa  
Cyma terminalis. filamenta corollae longiora. Bractea tres ad basin  
fructus. Embus brevi glaber cylindricus laxis coronatus margine integro

————— lobus.

Subcordata P. inermis, foliis cordato-subrotundis, fructibus siccis sublacinati pentagonis,  
angulis apice mucronatis Sw. prodr. pag 60.

Habitat in Antigua

Folia lata palmata glabra, nervosa integerrima obtusa. <sup>Cyma</sup> ~~conspicua~~  
~~est~~ Terminalis. ~~fructus~~ fructus unguiculati stratis angulis apice  
quinque mucronatis



Coccinea P. inermis foliis lanceolato-ovatis, pedunculis terminalibus caulis, floribus  
mutantibus, fructibus baccatis Sw. prodr. pag 60.

Habitat in Hispaniola.

Rami glabri flexuosi folia petiolata subtus pallidiora & pallida  
cania

elliptico lanceolatis

~~non red~~

Antiquarian