

Poa

plumosa Poa paniculata diffusa, calycibus setifloris, valvulis corollae
interioribus linearibus recurvis dorso ciliatis Retz off. 4. p 20
Habitat Tranquebano

michi ignota

barbata ^P panicula patula capillari, spiculis sexfloris oblongis, glu-
mis glabris caninatis Thunb. fl. jap. p. 1. v. pag. 114.
Habitat in Japonia

non vidi

amboinensis Poa panicula coarctata secunda culmo tereti Mart. 557. * L. V. p.

¹¹⁵
Phoainⁿ amboinica montana Rumph. amb. 6. p. 19. t. 7. fig. 03.

Habitat in India

Similema Poe compressa, sed culmus teres. Flores paniculati, verticillati. Flores quinque flori. Anthera rubra. Pistilla alba. C. M. L. C.

Distans Poa panicula ramis subdivisis, floribus ex spiculis quinquefloris, flospiculis
distantibus obtusis. Lin. S. V. pag. 115 Mant. 32. *

Poa divaricata; panicula ramis geminatis divaricatis, spiculis subquae
drifloris. Jouran. fl. & illustr. pag. 4. tab. 2. fig. 1.

Habitat in Europa australi et Barbana ad littora maris
Caesari plures diffusi ^{pro parte} binodes simplicis ^{vel ramosi} saporis foliorum tertri
folia angusta corduplicata. ^{bipollicidna} ligula membrana lanata
sed unculi radiales oppositi 2-4. Tripartiti capillares pedicelli exten
sum inmersi breves glumis cordatis superne apice membrana
iceo

— pedicellis glavatis

*Pennisetum P. panicula patula capellan, spiculi quinqueflori ob
longi glumis glabris Thunb. fl. jap. pag. P. V. pag. 114
Habitat in Japonia*

non vidi

anceps P. panicula diffusa, spiculis & flori acuti basi pubescentibus.
culmo ancepsit. Herp. prod. pag. 9.

Habitat. in nova Zelanda Fabricius glabro

Calami tenues ^{strati} geniculis fuscis folia angusta. Rami panicula
pubescentillata remoti capillares subimplexibus flori rari
longior spicula compressa glumae nervosa.

expansa. Panicula divaricata, ramis geminatis, spiculis peltatis inflexis. Quercus
illustr. tab. 2. fig. 1

pratensis P panicula diffusa, spiculis quoque floribus glabris culmo erecto.
Secreti Lin. S. V.

Gramen pratense paniculatum majus (latiore folio) Barck-puni 2] Sicut
gram. 177.

Habitat in Europa prae fertilissimis

in herbario desiccatur

Poa panicula laxa basi verticillata, spiculis raris subsquifloris, conellis
Lambert: La Mart. T. Enci: pag. 103
an *Poa flava* L.
Habitat in Virginia

flava ^P panicula diffusa, spiculis ovato-oblongis nigris L. L.
ray
Gramen praterse majus virginianum Pet. mus. 239.
Habitat in Virginia

nichi ignota

Anata Poo panicula diffusa capillari, spiculis glabris subsquellonibus, corollis
exquisite striatis La Mart. T. Enc. pag 103

Habitat in Virginia Carolina

Calyx brevis

Festuca glabra

Erhart ~~trineria~~
~~trineria~~

1. panicula diffusa spiculis subquinquefloris glumis acutisemis
Pollut. hist. pl. palat. no 27

~~Poa sylvatica foliis planis obtusis, culmo obliquo compresso, panicula diffusa spiculis ovatis subquinquefloris~~ ~~Corchus Will. fl. delph. 2. p. 120. tab 3. specimen~~ ~~Allionum figura vero omnino recta~~
Festuca sylvatica autem magis convenit.

Fest. Poa sylvatica All. ^{noy 48} ~~autuar.~~ fide specimeni a vna Hel. m. sp.
Poa trineria Erhart

Habitat in sylvis holstria, germania et Piedemont.

^{neura} Calmar ~~erdt.~~ ~~tripedales~~ y ~~calce~~ ~~tereti~~ ~~unde~~ ~~glaucesci~~ ~~glauco~~
vires. genicula 4 or. purpurascencia. - - Solia plana
late vinda, margine scaberrima glabra. Ligula laciniata
pedunculi inferiores 2-3. capillares spiculis parvis
flavo minime vero olivaceo compresso 4-5 floris pallide vixiter glumis
lin. potius hui spectat ^{calgis} at ^{corolleno} angusta ^{innato} attenuato
quod quoque in holstria, si recte memin. vdi

* Non vero Poa
sylvatica ut
credit all.

Festuca sylvatica foliis planis acutis, panicula divaricata
spiculis quadrisfloris distantibus acutis subantheratis
Will. fl. delph. 2. noy 105. tab 2. no 8 #

capillari^s panicula laxa patentissima capillari, foliis pilosis Cul-
mo ramosis pms. Lin. Sp. pl. ^{pl.} pag. 100 ♀
Gramen paniculatum virginianum, locustis minimis Monf. hist. 3. p. 202
J. P. t. 6. pag. 33.

Poa panicula laxa erecta, fructus, spiculis erectis oblongis pms. Virg.

136
Habitat in Virginia Canada.

Panicula capillata, strictissima, patentissima ramosis pms. - In
ca ex spiculis 4-5-6. minimis. Folia ad basem sili longis rari-
culis ramosis pms.

Culmi basi ramosi infirmi interdum prothamoi sili formes interdum
altiores. folia angusta spiculis linearibus v. ind. pubescentibus. Bad 9 flori-
ci

~~non vidi.~~

~~an alumbia non vidi~~

pedunculata Poa panicula plicata nitida, glumis patulis pedunculatis subsquifloris
culmo superne albo La Mark T. Eni pag 102 tab. 45. fig. 4

Habitat

glomo tener albido nitido mutico.

Festucoides Poa panicula elongata angusta intempta, spiculis subsquello
si flosculis glabris d. partibus La Mark T. Ceni pag 102.

Habitat

confer cum P. spicata

Lanceolata Poa panicula longa angusta interrupta, spiculis glabris ^{sub} ex
flori, glumis minutis
Poa interrupta La Mart. T. Enc. pag. 105.
Habitat in India orient.

an eadem cum mea interrupta?

Salina Poa panicula diffusa: speculo septemflori linearibus glume omnibus obtusis
fimis, carinis basi repente. Pöllner fl. Palat. 1. pag. 89.
Habitat ad Salinas Germaniae
Lacini Poa maritima and vespa

in h. h. v. p. nota

Ceniza P. panicula contracta, fl. spiculis 7 floris, glomerulis basi villosis

Poa ceniza culmo inferius compresso, pedunculis geminatis, locustis
Septemfloris, corollis villosis All. fruct. fl. Pedem. pag. 40.

Habitat in monte Ceniza

Culmi plures pedales erecti glabri inferne basi fo. geniculis
subnatis folia plana glabra margine fibrifera
ligula lanceolata. Pedunculi gemini ad pagulum dentem rach.
dis 2-3 flori alter interdum uniflorus, flexuosi breves
fl. spicula pallide vides vel purpurascens valvula
calycina lanceolata margine membranacea fo. recepta
culum et fo. glumis corollino basi villosis

procumbens. P. panicula lanceolata secunda cuneolata scabra; rachis tereti; quin-
li subquinquefloris, flocculis obtusiusculis nervosis Smith. Fl. Br. p. 96.
Poa procumbens

Curtis Flor. Lond. fasc. 6. tab. 11.

Poa repens With. arrang. of Br. pl. 146. tab. 66.

Habitat in ruderalis maritimis inundatis O

Glauca. Culmi plures, prostrati; spatharum, gemulati, foliosi, globi-
folii longissime vaginata, latiuscula, acuta, plana, nervosa, scabri-
uscula. Ligula acutiuscula, ~~et~~ erosa. Panicula rigida, semiglobi-
dnica, rachis ramulisque scabra. Spicula disticha, sessilibus; lanceola-
ta, 4-5 floris, glabra. glume calycis obtusa, valde inaequalis, lan-
nata: exterior sive nervis. Flocculi obtusiusculi, squenaves, apice
membranacei, erosi: glume interior ciliata Smith. l. c.

Cynosuroides ad Poam pertinet secundum Smith. ^{calycis} Aglumis novembris, rachis flexuosa
sa complanata differt.

Poa nitida La Martk

Herb.

Label

repteur Poa

Habitat

Herb. Richardi

habes

hirsuta Poa.

Habitat in Carolina herb. Kuhn.

Culmi pedales, simplices. Sola graminea, culmo altiora. Vagina hirsuta. Pa-
-nuculi longi, verticillati, filiformes, flexuosi. Pedicelli capillares, longi.
Panicula patens, subcomposita. Glorculi glabri, calyce longiores. Spicula oblon-
ga

per Anacra Psa

Habitat in Allemani herb. Rich.

habes.

pectinata Poa

Habitat

leb. Richan.

habes

Poa

Habitat ad Senegal herb. Richard.

habeo.

Dioica Poa

Habitat ad Littora fluvii Naasha hinc Am. sept. Michaux herb. Guss.
Caulis repentes, teretes, elongati, ramosi, villosi; rami et folia in geniculis
aggregati. Vagina brevis, ^{spica masculis pedicellat.} striata, villosa, ut folia. Solia vix pollicis
na, linearia, acuta. ~~Spica~~ panicelli pauci, subramosi, brevissimi, vil-
losi. Spicula lanceolata, 15 flora. Valvula corollina externa tri-nervia
Spicula foemina copiosiora, ^{pub.} sespiter, in capitulis multifloris: flosculi pluri-
bus

Poa

Habitat in insula Barbina Commesf. herb. Græc.
Culmi plures, fere pedales, simplices

habes

Coccoloba Poa

Habitat in America boreali herb. Presp.

Folia graminea, plana. Culmus glaber. Paniculo patens. Spicula verticillata
la, ceteris, acuta, quadriflora, basi inclusit perlopha.

Poa

Habitat frequens in horto et agris montis Vides Comnes. herb. Tus.
Culmi plures, geniculati, basi rotundi, pedales

herbes

1
Caryophyllus Pea

Habitat in monte video Nömmes, herb. Auf.

Caules striatus, teres, glaber. Folia conduplicata, glabra. Panicula coarctata
palmaris, hirsuta. Spicula 6 flora. Calycis glume subulate, dorso serrate, spi-
cula breviores. Flosculi basi filis longis lanati, uti valvula exterior inferne
et marginis: valvula exterior lanceolata, attenuata, basi purpureo-roseo, apice
hyalino.

municata Poa spiculis trifloris floribus duobus ultimi margine tuberculatis setigeris

Habitat ad Serram Leonara herb. Thourni

Culmi pedales, erecti, simplices, inferne foliosi. Folia bi pollicaria, lanceolata, unquam vel parum ultra lata, nervosa, glabra, teria. Panicula simplex: pedicelli capillares, solitarii. Spicula lanceolata. Calycis gluma altera brevis. Corolla valvula lanceolata, mucronata, calycis ^{super}duplo longior, floretuli inferioris valvulae exteri teria duorum superiorum ad marginem versus apicem tuberculis 3-5. quibus infident pila retroflexa Albidis

aurea Poa

Habitat circa Peking Herb. Incarn. apud Auz.

Caesulis odorati. Culmi striati, glabri. Panicula palmata; pedicelli
capillares contracti. Calyces argentei nitidi, longitudine glomerulorum. Flo-
culi aurei, margine ciliati mucici.

caesia P. panicula diffusa. Spiculis ovatis quinquefloris, glumis lanceolatis
subsericeis liberis, ligulis brevissimis obtusis Smith. Fl. Br. pag. 103.

Habitat in Scotia 4

glauca. Culmus pedalis, erectus, teres, striatus, tenuis, basi versus
nodis et foliosus; superne nudus. Folia lanceolata, obtusiuscula, plana
dorso basi versus longata, ceteroquin scabra. Vagina longitudine foliorum
striata, subscabra. ^{Ligula} Spicula valde variabilis, aliquando brevissima, quan-
doque elongata, lanceolata, laevis, proecipue ad folia caulina. Panicula
erecta, ramosissima parum elongata, ramis semi verticillatis, elongatis angu-
latis, scabris, ramulis binatis. Spicula ovata, erecta, saepius quinqueflora
glauco albido, purpureoque varia. Gluma calycina subaequalis; lanceo-
lata, acuta, cuneata, obsolete trinervis, apicem versus scabra. Flosculi
remotiusculi, lanceolati, obtusi, apice scaniosi, erosi, dorso lanati, margi-
ne scaniosi, sericei, basi liberi; valvulis interioribus margine scabris;
sati flexuosa, villosa Smith l. t

a glauca Fl. Dan. diversam credi. quod glauca Withenry.

Panicum flexuosum P. panicula flexuosa, spiculis trifloris, glumis ovatis basi villo connexis
ligulis omnibus lanceolatis Smith. Fl. Br. pag. 101.

Habitat in alpebus Scotia. 2

Culmi plures, spathamoi, adscendentes, geniculati, subtrinores, glauci, super-
ne angulati, striati, glaucescentes. Dolia erecto-potentia, plana, angusta,
acuta, supra scabra, vaginis omnibus elongatis, latis, panum compressis, striatis,
glabris. Ligula acuta, foliorum caulinarum saporis laceni. Panicula abbrevi-
ata, ovata, subcondensata, ramis biniatis, alternatim ramosis, magis mi-
nusve basi flexuosis, superne angulatis, scabris. Spicula lata ovata
compressa, glauca, albedo et purpureo variegata, plerumque triflora.
Gluma calycina panum inaequalis, lanceolata-ovata, acuta, vix folia-
ta, carina scabra. Glorculi lanceolati, acutiusculi, margine denu-
dati, carina scabri, basi villis complicatis lanuissimis connexi. Val-
vula interior vix ciliata. Smith l. c.

P. alpina diversa, colore glauco, culmo magis folioso, foliis angustioribus, ac-
tioribus, supra scabri, vaginis longioribus, striatis omnibus confertibus, pedun-
culi flexuosis, glumis angustioribus, basi villo connexi nec liberi, neque cari-
na neque margine setoso-priceo Smith

expansa Poa panicula patentissima capillari, axillis ^{un} pedunculorum barbati,
speculis 10 flori, vaginis punctatis pilosis
Habitat in Insula Francia de Thourin

~~Cyrtus paniculatus~~
Cyperus pedale, ramosum. Palmus
Palmus vaginis foliorum texti

folia involuta, subulata, suprema plana nerva, et hinc glabra, intus
pilosa. Vagina subtilissime striata, versus marginem punctis
sparsi numerosis quibus insident pili longi. Taxis hirsuta

Ramosula longitudine Pedunculi tres ad singulum geniculatum, simplicissimi, bis- tripolli
Culmi, patentissima. Cases, superiores flexuosi, capillares; uti pedicelli, pollicares

vel sesqui pollicares, ^{plurimumque simplicissimi} alterni, remoti, uni flori.

Speculo lineares, 10 flori, compressi, Sto pallide flavi dorso di
Palyx parvus, ^{subtus punctatis apice o} purpurascens. Sto pallide flavo

dorso dilute purpurascens, apice obtuso, saturatioribus, maturi
scilicet basi gibbosi, subfusi a femine fusco translucente

[Affinis P. capillari

in Poa tenella vel punctata satis diversa

~~Roemerii Saich. panicula spicata, floribus dracenis, culmi geniculis barba
tis foliis planis Retz off: 5. p. 16.
Panicum spicatum Thunb. p.
Habitat in India orientali~~

or ^a Lin. spec: D. west

Poa littoralis

*Poëus ad Alopecurus referenda, a genere Cynopii plane diversa est ad quæ
ta fuit a Commersonio*

~~Lesleia magellanica ^{panicula} ~~spicata~~ puberula oblonga~~

~~hab. ad fratum magellan.~~

~~Culmus erectus glaberrimus
folia glabra~~

~~Panicula spicata oblonga~~

~~bractea nulla
gluma calycina oblonga villosa carinata apice margineque reflexa
apicem fusca apices divaricati villosi~~

~~Corolla valvula 1. dorso aristata, calyce brevior glabra
arista recta ~~ca~~ calyce duplo longior
in apicem membranaceum vidi inter stamina at pistilla nulla~~

Dichcha P. *Spica* D. *Sticho* terminali hinc convexula illinc plana
S. V. p. 115. *Gayu* m. p. 2. p. 74

Poa *falenoid* ex *All.* flor. *pedemonta* 220d. Tab. 91. fig. *Spica*
ovata, *locustis* pubescentibus *sex* floris obtusis *all.* flor
ped. no 2208 Tab. 91. fig. 1.

Habitat *Pedemontii*

24

non vidi

compressa P. panicula coarctata secunda, culmo obliquo com-
presso L. V. p. fl. Dan. 1. 742.

Gramen murinum radicis repente Brant. pan. 2. prodr. 2.

Gramen paniculatum, radicis repente, culmo compresso Vaill. panf.

Loisium deflexis pulchellis Schumbr. gram. 148.

Gramen paniculatum, radicis repente, culmo compresso Vaill.
panf. 91. 1. 10. fig. 5.

Habitat in Europa et America septentrionali siccis, minus, siccis

6-7 fl.

in herbis deflexis

Stolonifera Poa panicula est paucis spiculis compressis per floris culmo
stolonifero Bellard. app. ad fl. Ped. pag. 9. tab. 1.

Habitat in montosis Pedemontii &

Capitata. Culmus terminalis nudus et nodis striatus erectus
folia brevia planiuscula planiusculis subnerviis subtus striatis
Panicula divaricata: spicula inferiores longius pedunculatae binae
tertia: pedicelli subflexuosi spicula per floris Bellard.

alpina ^P panicula diffusa ramosissima, spiculis sex flori corda
tis. Lin. S. V.

Gramen alpinum paniculatum majus, panicula spicosa variegata
Scheuchz. gram. 106

^B Gramen alpinum latifolium panicula laxa foliacea, foliolis in
panicula paucioribus et magis crispis Scheuchz. gram. 112.

Habitat in alpebus Norvegiis, Helveticis

gramen subcepsitopem, spithamoum, culmus simplicis et pum
eritus binodis, diphylleis. folia plunima radicalia lineari
lanceolata vel parum ultra glabra corva ligula membra
nacea oblonga acuta panicula subdiffusa. Spicula ovata
subsex floro purpurascens vel vides valvulis apice ferdide
flavescens valvula calycina canata canna glabra e serotani
mentis oculo armato vides uti corollina margine membrana
cea. glume corollina inferiore villosa prosertim carina et mar

^{gna}
folia in ovipara angustiora convoluta
panicula subdiffusa spiculi — ovatis glabris basi villosis ^{coloratis}

aquaticum paniculata diffusum, spiculis sex floris lineamentis
L. V. p. 1134

Gramen palustre paniculatum albescentum Bauh. pin. 2. Theatr
30. Scheuchz. gram. 191.

Gramen paniculatum aquaticum latifolium Bauh. pin. 3. Monof
herb. 7. p. 201. f. D. V. b. fig. 25.

Habitat in Europa ad ripas fluminum, fluviorum &
Culmum tripe vel quadripedalis erectus glaber. folia ensiformia
pedalia, semipollicem longa glabra, marginibus serrata
Panicula pithamosa, pedunculi distantes inferne 4-5 ad
aggregata inaequalis scaberrima. spicula linearis sublanata
et sepe unguis semurquiculares valvula oblonga obtusa calycine
Cores cotillina nervosa

maritima Poa panicula diffructa sciunda ovata inferne patula, ramis subternatis, spiculis
glabris, foliis planis culmis adscendentibus Host. Kew. 1. p. 100

~~Poa maritima Hudf.~~

Gramen paniculatum maritimum vulgatum primum Ray. syn. 409

Poa maritima panicula coarctata sciunda ramosa, ramis binatis,
spiculis oblongis; flocculis obtusis, foliis involutis acutis,
culmo obliquo Texti Hudf. fl. Angl. pag 427

β. Gramen caninum maritimum paniculatum R. syn. 410

Ab habitat in littoribus maritimis Europa septentrionalis &

facit Poa aquatica et omnia minor. folia siabra angust.
hucula linearis obtusis ^{utrum} prope aqua ^{prospere} mane ^{colore} brunnea
^{prope} lente subleat ^{more}
necumino gouvani foliis involutis gaudant forte a loco
tantum loci

mantemum T

arenaria Poa panicula ramosa, spiculi verticillati sex flori, glumis obtusis
lobis margine membranaceo La Mart. F. Eni: pag. 183
P foliis angustis convolutis spiculi quadrilobis.
Habitat in montibus arenosis ?

michi ignota

Dwaniata Poa panicula *Dwaniata*, spiculis 6-floris, foliis planis striatis
Vill. fl. delph. 7 pag. 127
Habitat in montibus delphicis &
Lepidali folia plana.

non videri

Penella panicula oblonga capillari subverticillata, floribus
flexuosis minutissimis nutantibus *Lin. l. V. p. 114 & Spec.*
pl. pag. 1014

Thama pulla *Reed. mat. 12. p. 75. t. 4.*

Gramen paniculatum, Col. uti tenuissimum subrotundis *Burm.*

Zeyl 710. t. 47. fig. 3.

Gramen paniculatum ex oris malabaricis, panicula deliata
re Plukn. alm. 176. t. 300. fig. 2.

Gramen fumi *Rumph. amb. 6. t. 4. fig. 3.*

Habitat in India

Culmi filiformes geniculi inferne infracti pedales; folia
angustissima glabra. Panicula primum ^{ped ramis vix raris} contracta demum
diffusa ramis capillaribus flexuosis. Specula minuto et bfo
sa valvulis cor hyalinis corollinis utraque linea viridi marginibus
lente usque ciliatis vagina ore barbato uti et articuli basi pedun-
culorum

articuli raris barbato