

Pyrola

1. *rotundifolia* P. Ramenibus ascendentibus, pistillo declinato, scapo bracteato
2. *minor* P. Ramenibus pistillisque rectis, scapo ebracteato.
3. *secunda* P. racemo unilaterali
4. *umbellata* P. pedunculis subumbellatis, foliis lanceolatis obtusis supra cuneatis
5. *maculata* P. foliis ovato lanceolatis planis acutis, pedunculis bifloris
6. *uniflora* P. scapo unifloro

muironata Pyrola

Habitat in alpebus Carolinae Michaux herb. Fusf.

Folia radicalia, cordato ovata, glaberrima, rigidiuscula, bipollicaria, longiora quam lata, crenata, crenis mucronatis, obsolete venosa. Petiolis folio longioribus. Stipulae ad basin, lanceolatae. Scapus longitudine fere foliorum, induratus, glaber, uniflorus. Squamae nonnullae superne, acuminatae. Petala flavo viridia. Capsula 3-valvis, septis Pentameris Fusf.

Pyrola

Habitat in America boreali herb. Musci

Radices longae. Caulis gemini, infra folia angulati. ^{Lea} Folia radicalia ovata,
per petiolum non decurrentia, glabra, acuta, serrata, vix venosa. Scapi pi-
thamici, teretes, ad apicem squamis remotis, ovatis, minutis. Racemus secundus.
Styli recti

1 rotundifolia Pyrola Raminibus ascendentibus, pedicello declinato Stans breviterato
Linn. V. p. 408

^{Geor. Bot. t. 110,}
Pyrola Brunfels. Dod.

Pyrola vulgarior Cusi h. u.

Pyrola rotundifolia major Bank. Pin. 191.

Pyrola Nov-eboracensis Cold. Noveb. 99

Habitat in Europa septentrionaliore, Virginia, Brasilia &

Radix repens. Folia radicalia longe petiolata ovata vel subrotunda
pallidiana, obtusa, crassa subcoriacea, obsolete remote serrata
nervi folio glaberrima nervo venosa. Scapus erectus simplicissimus, semi-
duplo longior pedalis adpressus gemmis remotissimis lanceolatis. Racemus termi-
nalis, tri-pollicaris. Flores nuda nati

Capo ebracteato

2 minor *Pyrola* flaminibus pedicellis que rectis, ~~Capo ebracteato~~ ~~Fl. Dan. 5. 5. Gr. flor~~

Pyrola minor Raj Angl. 3. p. 363.

Pyrola foliis subrotundis, scapo utriusque racemoso genitalibus rectis Act. Holm.
1741. p. 192.

Pyrola floribus racemosis dispersis, flaminibus pedicellis que rectis L.S.V. p. 408. Fl. D. 1⁴⁵ ~~1⁴⁰~~ Gronov.
tab. 4. t. 56 ⁴ Habitata in Europa frigida.

Similis procedente at folia ~~oval~~ ^{ulata, venis minus exstantibus} ovalia obscure crenata petioli longitu-
dine foliorum, teneriora. Scapus brevior absque bracteis, flores ~~minus~~
Duplo minor

3. *Secunda Pyrola racemo unilaterali* Lin. S. V. p. 400 Fl. Dan. V. 402.

Ambrosia Dalech.

Pyrola secunda Clus. hist.

Pyrola folio mucronato serrato Bauh. Pin. 191.

Habitat in Europa borealis sylvis &

*Caulis erectus glaber simplicissimus subangulatus inferne nudus. Folia
petiolata approximata ovata, obtusa mucronata, tenuissime serrata,
unquicularia. Pedunculus terminat caulem, erectus; caule longior,
appressus squamis ovatis, obtusiusculis distantibus. Flores in racemo
scillitani secundi nutantes.*

pedunculo bracteato

4 umbellata *Pyrrola* ^{pedicellis} ~~pedunculata~~ ^{foliis lanceolatis} ~~subumbellatis~~ ^{superioribus} ~~lin. t. v. pag. 409~~

Pyrrola *tertia* *fruticans* *Clus. Pan. p. 507.*

Pyrrola *frutescens*, *arbutifolia* *flore Bauh. Pin. 191.*

Habitat in Europa, Asia, America septentrionalis sylvis.

Caulis erectus, glaber superne ramis binis purpureis ^{angulatis} fl. angulatis. Folia
lanceolata, aggregata, ^{superioribus} basi in petiolum breviter sinuata, glaberrima
supra lineata, ^{superioribus nervos} subtus lœvia pallidiora, obtusa, conacina, pollicaria
a medio ad apicem argute serrata. Pedunculus terminalis bipollicaris
erectus. Pedicelli umbellati 4. s. quæ.

5 - maculata *Pyrola* ^{olatu} foliis ovato-lanceatis pl. planis, ^{acutis} pedunculis bifloris
Pyrola manlandica minor, folio mucronato arbuti *Pest. musf. 679.*
Pyrola mariana, arbuti foliis angustioribus, trifoliata ad medium nervium
linea alba utrinque per longitudinem discurrente *Pluch. mart. 157. t. 349.*

Fig. 4.
Pyrola petiolis apice bifloris vel trifloris *Gron. Virg. 40.*
Pyrola maculata pedunculis bifloris *Lin. l. V. p. 409.*
Habitat in America septentrionalis sylvis h.

Caulis erectus palmari vel spithamovis inferne nudus superne bifidus glaber
sicut tota planta. Folia ^{breve} petiolata, aggregata, plerumque ternis ovata,
unquiculana, acuta, anguste serrata, glabra, absque lineis secundum
nervos ut in umbellata. Pedunculi ^{breve} sive medio linea alba. Pedunculus
nudus longitudine plantae, biflorus: floribus nictantibus

6 uniflora *Pyrola scapo uniflora* L. N. p. 409

Flor. San. Tab. 6

Pyrola 4. minima Clus. hist.

Pyrola rotundifolia minor Bauh. Pin. 198.

Uniflorae ampliore Monf. hist. 3. p. 505. f. 12. t. 10. fig. 2.

Habitat in Europa boreali Sylvis &

Scapus erectus, simplex, digitalis, et ultra. Folia radicalia petiolata, sub
rotunda acutiuscula inaequaliter serrata, venosa, pollicaria. Coste termi
nalis maxima in hoc genere