

4. Qualea 7

5

7 Cal. 5-phyllus, foliolo unico majore basi ^{glandula} nectaris calcarato. Cor. 1-petalus
Peric. 2-loculare, polyspermum

rosea Qualea petalo integro, nectario calycis breviora

Qualea rosea; floribus amplis, carnis Aubl. fl. Gu. 5. tab. 1.

Habitat in Guiano Sylvis

~~Asteris copulatis Aubl~~ Rami inferne teretes, superne tetragoni,
subrillosi. folia petiolata, bip. tripollicaria, rigida, oblonga,
attenuata, utrinque glabra, transversim subtilissime ^{confertim} ~~sub~~ multi-linea
sublineata, subus transversim nervosa, nervi ante marginem concurren-
ter nervumque crenatum utrinque prope marginem forman-
tes, inter nervos venoso-reticulata. Racemi terminales, erecti,
s. 6 pollicari Petiolus sive unguiculari, glandula impressa, nigra
ad basin. Racemi terminales, erecti, s. 6 pollicares: pedicelli so-
pe geminati. Calyx extus tomentosus. Petalum multo majus quam in
sequente. Germen hirsutum

~~rosea Qua lea floribus rubris petala infimis ^{ro} ~~est~~ ~~caerulea~~ La Mart.~~

~~caerulea Qu. petalis emarginatis, foliis acutis.~~

~~qualea caerulea pet. floribus parvis Aubl. fl. Guian.~~

~~i. p. 7. tab. 2.~~

~~qualea caerulea floribus inter-subcoruleis petalo infima
emarginato La Mart. Enc.~~

~~Abitat in fylis. Guiana & in herb. La Martii &~~

~~Rami teretes, glabri, purpurei center. Corolla petiolata, opposita, bif.
tripolliciana. Rami ex axillis supremis et apicis, longitudine
folii simpliciter.~~

~~Petalis 9 phyllis duo opposita, oblonga, obtusa, altero parum majore, duo intenor, ma
yo ovata, acutiuscula, altero margine rectior, altero arcuata, quantum reliqui
duplo major subrotundum, obtusum, emarginatum latitudine & longitudine
equali omnia extus villosula. Petalium unicum, magnitudine folioli
calycis etc. Germis parvum villosum, unicum. Stylis file formis
petalo brevior. Stigma obtusum. Glandula tres, erectae, obtusa ultra calycis
foliolo ^{est} parum prominens, ascendens, ca. nigrum & basi calycis enatum.~~

caerulea Qualea petalo emarginato, nec tano longitudine calycis.

Qualea caerulea floribus parvis Aubl: fl. Guian. 7. Tab. 2.

^{ex herbario}
Habitat in Guiano sylvis. Herb. la Martini 5

~~Culm. ex oppositis Aubl.~~ Rami teretes, glabri. Folia petiolata,
~~opposita~~, bi- & tripollicaria, oblonga obtusa acuminata vel atte-
nuata, utrinque glaberrima, confertissime nervis tenuissimis lineis

apice obtu-
siuscula

to, integerrima. Racemi ex axillis supremis et apice,
longitudine foliorum ~~simpliciter~~. Calyx pentaphyllus: foliola duo opposi-
ta oblonga, obtusa, altero parum majore; duo interiora magis ovata, acutius-
cula, altero marginibus rectionibus, altero ~~marginem~~ arcuato; quintum
religuis duplo majus, subrotundum, obtusum, emarginatum; omnia
crassiuscula, extus villosa. Glandula e basi calycis ad extus ad
foliolum calycinum majus exatum, teres, adscendens, cartilaginea, obtu-
sa, longitudine foliorum ^{calycis} minorum. Petalum unicum, magnitudine
folioli 5^{to} calycis, at multo tenerior, subbilobum. Germe parvum
villosum, incanum. Stylus filiformis, petalo brevior. Stigma obtu-
rum.

~~caerulea~~ Obt. Folia opposita, petiolata, integerrima. Racemi ~~simpliciter~~ vertice

nigricans Justicia species terminalibus distichis, bracteis & staminibus, foliis lanceolato-linearibus
nigricans Justicia, foliis lanceolato-linearibus obtusis
nigricans Justicia Lour. fl. Cochinch. pag. 24.

Habitat in Cochinchina
Justicia 6 pedali. Folia integerrima, crassa subsessile. Rami
et folia linearibus nigro lineata. Flores albi rubro variegati in
caudicibus terminalibus. Carol. fabrum superius integrum erectum
inferius 3 fidum, reflexum bracteis foliaceis. Lour

Habitat in Cochinchina. Justicia 6 pedali. Folia integerrima, crassa subsessile. Rami
et folia linearibus nigro lineata. Flores albi rubro variegati in
caudicibus terminalibus. Carol. fabrum superius integrum erectum
inferius 3 fidum, reflexum bracteis foliaceis. Lour