

Randis

parviflora Prandia
in domino
Habitat herb. Tusc.

Rami teretes, glabri, cortice cinereo, decussatum oppositi. Folia petiolata, saepe aggregata, angustulana, oblanceolata, obtusissima, basi acuta, in petiole, integerrima, arenaria, enervia. Spina angustulana vel semipollinaria. Bacca & Calyx per se, magnitudine pisi, calyce coronato.

parviflora Randia

Sloan. tab. 207. fig. 1.

Habitat in America herb. La Manthii

Rami rigidi, glabri, patentes, cortice griseo. Spino opposito patentes. Folia
plura, semurquiculata, oblonga, nitida, basi acuta, acuminata, enervi. Flores
axillares, solitarii, septes, parvi. Fructus magnitudine pepere

rotundifolia Brandia

Brandia rotundifolia; spinis ramisque subrectis illatis, foliis subrotunda
diis ovatisque rugosis Fl. Per. 2. pag 68.

Habitat in Pennis nemoribus.

Caulis fruticosus, ^{erectus} oxygali. Rami teretes; oppositi quaterni. Spi-
no rugido oppositi quaterni. Folia quatuor ex singulis punice
gemmateo, subrotunda ovataque, integerrima, rugosa, ut denique
pubescentia, parva, subsessili. Stipula interfoliaceae, ovatae, acutae
membranaceae, fulvae. Flores solitarii ex centro quatuor foliorum
sessiles. Corolla alba, calyce duplo longior. Bacca unilocularis
magnitudinis nucis avellanae lutescens. Pericarpium plerumque ovatum, rudi-
cantia, obscure fulva, pulpa nigra fl. Pennis

obovata Randia.

Randia obovata Spenis, oppositis, foliis obovatis acutis fl. Peruv. 2. pag
68. Tab. 220. fig. 6.

Habitat in Andium pronytis calidis 5

Pubescens. Paulis fructu osus, oxyphalis, erectus. Rami brachiati, infer-
ne floridi spinosi, superne foliosi inerme: Venae obsolete, tetrago-
ni, compressiusculi. Spina axillares, brachiato, horizontales; nigrae
Folia opposita, approximata, breviter petiolata, obovato-oblonga, acumi-
nata, integerrima, venosissima, bipalmata. Stipulae interfoliaceae,
basi connatae, subulatae, erectae, petiolis longiores, membranaceae.
Flores ^{intra axillam} ex axillis circiterium foliorum, solitarii, sessiles. Bractea
septem sub singula flore, lanceolata, acuta, imbricata, membra-
nacea, marcescentes, fulvae. Corolla alba calyce duplo longior
Baccis pubescentibus, luteo-cinereis, nucis mucatoe magnitudine
fl. Peruv

folia lanceolata oblonga basi in versus
sensum angustata