

Raphis

mon. t. nov. 1792

Cal. perianth. 1 phyllum *spidum* laciniis ovatis acutis erectis
Corolla petalo s. calyci inserta obovata erecta calyce breviora
Stamina filamenta sive nulla anthera septis s. in calyce
germen superum, hemisphaericum, glabrum. Styli duo bifidi segmenta acuta

~~Tomar~~ *Erp. Paphis*

Lucida *Raphis*

Habita in Cap. b. India Orientali 5

Rami glabri alterni e lineis a petiolo decurrentibus angulati tetragoni

Folia alter petiolata alter subopposita pollicaria obovata

^{atq;} glaberrima nitida, nigra obscure crenata.

Petioli semicirculares supra canaliculati

Stipula pubulata ad basin petioli

Pecunuli ad latius petioli teres glaber longitudine petioli tri vel qua

driflorus. floribus breviter pedicellatis

Bractea ovata acuminata ad basin calycis

Trisalis Rhapsis Lour. flor. Cochinch. pag. 553

Cuspe Rumph. amb. lib. 10. c. 8. tab. 5. fig. 1.

Habitat ad vias Cochinchinae et Siam

Palmus pedalis per nudus teres simplicissimus. folia subradicalia longi-
cula tercia. spica terminalis solitaria conica laxa. ♂ Gluma
1 floro vel rursus subulata mutico Per. 2 valvis valv. ciliatis muticis
ham. 7. ♀ sessilis in fra masculor in eodem involucro. gluma
collata ^{altera} a rista longiuscula.

non est