

Rathea

Robinsonides hinc transfer *Robinia squamata*

inermis Prothea

Habitat in Guinea Thonning 5

Erudax ^{ter. rar.} bival tripedale, ramosus: rami alterni glabri, lovis costae fusco
perpurascens. Folia petiolato, pin recta cum impari, 6 8 puga: foliola
oblonga, semipollucina, petiolato, alterno, obtus, mut ronato, pubtus glauca
avenia. Stipula ut naque linear - lanicolata, bravis. Racemi axillares
caeci, solitarii, pauciflori. Braetee ad basin calycis, lanicolatae.
Calyx monophyllus, hinc gibbus, tuberculis atro viridibus punctatus,
apicis: lacini acuti, superior bifida, manescens. Corolla flava,
venis atro viridibus. Ventellum ovatum, emarginatum, intus basil
le duobus alar comprimentibus, reflexum. Ala sexillo breviores; ora
et base obliqua, erecta, conniventes. Carina compressa, gibba, base
bifida; ungues longitudine calycis. Silamenta 10, omnia connata:
anthera porita. Germe pedicellatum, lineare, tenuis, compressum.
Stylus bubulatus. Nectarium basin margo basin germinis cingens.
Legumen pedicellatum, 3 5 articulatum. articuli striati, compressi ^{tuberculis}
monospermis, oblongis. Semina ovale, compressa, glabra.