

Salicornia

10

~~Salicornia~~

9. Salicornia. Cal. ventriculosus, integer. Petal. 0. Sem. 1. (Stam. 1-2.)

herbacea Salicornia herbacea, patula, articulis apice compressis emarginatis,
spicis axillaribus oppositis pedunculatis, squamis obtusis Willd. Spec. pl. t.
p. 23.

Salicornia Dod. pempt. 84.

Salicornia herbacea; herbacea, patula, articulis apice compressis emar-
ginato-bifidis Lin. ^{Syst.} ~~Spec. Pl.~~ 53. Basl. opusc. v. 2. p. 105. t. 10. 86. Dan. 8. 303.

biennis β . Woody jointed Glasswort. Pet. 46. Brit. t. 9. f. 4.

Salicornia erecto, foliis brevibus, cypressiformis Dill. in Raii Syn.
p. 137.

~~myosuroides~~ Salicornia ramosior procumbens, foliis brevibus purpureo-
Dill. in Raii Syn.

myosuroides γ . Salicornia myosuroides-procumbens, surculis longissimis
Dill. in Raii Syn. p. 137.

purpurascens δ . Salicornia ramosior procumbens, foliis brevibus purpu-
rascentibus Dill. in Raii Syn. p. 137.

major E. Kali geniculatum major, semper vivens Lamour. Schola bot. 37.

Habitat in Europa littoribus maritimis arenosis. C. J.

Caulis erectus vel adscendens, ramosus, comi oppositi ~~quidam~~, articu-
latis: articulis apice dilatatis, ramosis. Rami oppositi. Spica ~~capillares~~
et ter parum minores terminali, teretes, attenuatae. & longiores in singulis arti-
culo utrinque oppositi, intermedius superior. Squama calycina oblonga, ob-
tusa.

B. gaudet caule ramosiori, specie longioribus et crassioribus; & caule diffuso,
specie longioribus; & caule ramosissimo, specie parvis, purpurascensibus.
& caule ramosissimo feraciori, specie brevibus, viridibus. Smith. fl.

Angl. 1. ² ~~partes~~

Stamine ² numerata. Foster in Act. Angl. 1767. v. 57. p. 312. Quodque

sic etiam Moching in Ek. Beitr. 7. p. 178. Moching^{us} observationem ve-

ram esse: scilicet quod anthera unica primum apparet, haec evanescente & adla-

tus trianguli, haec evanescente prodit altera ad oppositum latus post aliquo-

dies, ex iteratis observationibus confirmare aucto^r proprio a-

topia asseverare queo.

india² Salicornia articulis clavatis compressis sicut truncatis bidentatis,
spicis terminalibus cylindricis solidariis obtusis Wild. in Act. Soc. Am.
hist. nat. hist. Ind. vol. 2. p. 111. t. 4. f. 2

Habitat ad littora maris in vastis planitiebus copiose prope Franque-
las Pötker.

Caulis diffusus, decumbens, rarius in alto repens, basi tantum ramosus et
multo crassior quam S. herbacea, uti tota planta glauco virescenti
rami oppositi, ramosi, articuli semirungiculares, denticulis articulorum
carinati Pötker

Spicis terminalibus, pollicares, vel ultra, crasso, teretes, e squamis
truncatis arte imbricatis.

perfoliata L. fruticosa foliis alternis uncinato orbiculatis perfoliatis

Forst. Jap. pag. 3. t. 1.

Ramis subdichotomi

Habitat in Arabia

Truncus diffusus pedalis flexuosus. Folia carnosu vix serotina
quicquidaria, congesta. Spica pedunculata axillares
et terminales semipollicares. Vires. flores terni

Caulis lignosus interdum non palmatus
foliis alternis perfoliatis orbiculatis perfoliatis ramis subdichotomi

3

perennans Salicornia herbacea patula, articulis apice compressis
quarto bifidis, spicis axillaribus ternis pedunculatis, squamis acutis
Willden. Spec. pl. 1. p. 24. †

Salicornia herbacea P. Pall. It. 4. app. no 89. f. 2. 1.

Habitat in paludibus siltionibus in Siberia ad Taitum &

Radix perennis. Caulis prostrati, tenuiores et ramosiores quam in
herbacea. Pall. l. c.

amplexicaulis Salicornia Gleis cordatis amplexicaulis

Habitat ad Littora lacus prope Bardo Tuneti vel parum ultra. Esqui
Caulis decumbens, ^{besti frutescens, palmari vel parum ultra} rami alterni, patenti simi, pollicia
res, bini semper approximati, ex omnibus axillis foliorum

Folia parva, alterna, obtusa, fuscula, subtus convexa, supra plana
uscula, amplexicaulis, in ramis junctibus imbricata

(A Salicornia perfoliata Tost. quo forte eadem cum Sal. arabia
Len. omnino diversa)

4
fruticosa ⁴ Salicornia caule erecto fruticoso; articulis ramulorum anticipitibus,
Equamis floralibus truncatis membranaceis Willd. Spec. 1. p. 24

Kali Camer. Epit 246

Salicornia fruticosa, caule erecto fruticoso Lin. Spec. pl. p. 5.

Habitat in Europa maritima. 5. articuli subquadratis

Caulis ramosissimus, rami oppositi. Spica sessiles, oppositae, un-
guiculares, obtusae.

virginica Salicornia herbacea, erecta, ramis simplicissimis Lin. ~~S. Vp.~~ ⁵
Syst. veg. 52. t.

Habitat in Virginia

Rami oppositi, indivisi, terminati spica longa, articulata. In specie
quodam Virginiana, quam nuper ante oculos habui vivam, erat
Calyx squama ovata cochlearis instar concava erecta. Stamina cer-
te duo. Pistillum superum. Lin. l. c.

Strobilacea S caules prostrato fruticoso, articulis trinatis alterna
tina spiciferis, spicis nudis brevissimis oppositis ^{decussatis} Wild.
Salicornia arborescens sine gemulis Buarb. cent. 1. p. 6
tab 10. fig 2.

Salicornia strobilacea Pall. it. 1. app. no 91. tab E

Habitat in proreptis salpini mari capuei 5.

Caulis prostratus pro rami erecti ramosi spica cylindri-

co obtusa

suffruticosa pedalis squa opposita densa cetera flores unis-

20 in 3 in quila pur

6

cruciata ^{fruticosa,} Salicornia articulis inferioribus ^{sub} globosis, superioribus cylindricis
cruciatis.

Salicornia cruciata, fruticosa: pedunculorum articuli cruciatis
caulis globosis Forsk. descr. p. 2.

Habitat Alexandria ad Salinas 5

Caulis procumbens, teres. Rami conferti, erecti, articulati: articuli
inferiores steriles, minores, & sextuplici ordine squamarum tuberculati;
superiores majores, floriferi, transverse cylindrici, squamati, floribus
sparsis, nudis ad apices squamarum, quo bullulento Forsk. bi

~~175~~ Forsk. bi

~~Tetraphylla *H.* foliis quaternis oblongis obtusi lin. Suppl. p. 81. L. V. p. 53.
Hippuris lanceolata foliis lanceolatis, inferioribus quater-
nis, saepe nonibus quinis Retz. off. 13. p. 7. t. 1.
Habitat in Suecia circa Aboam. in Finmarkia circa
Wadfoe. Caulis ^{sed ultra}
Caulis tetragonus exserto palmari folia linear
Lanceolata internodiorum longitudine saepe tri-
infima. ^{inferiora remota} ^{et non ovata. reflecta} cum versus confertiora~~



7
Strobilacea 7 Salicornia caule prostrato, articulis ternatis alternim
spiciferis, spicis nudis brevibus bimis oppositis Wed. Spec. pl. 1. p. 25
Salicornia arborescens sine geniculis Burch. Cent. 1. p. 6. f. 10 p. 2.
Salicornia Strobilacea Pall. It. 1. app. no. 91. t. 1. E.

Habitat in promptu Salis maris Caspici

Caulis suffruticosus, pedalis. Rami erecti, ramosi. Spicis sessilibus, densis
seris, cylindricis, obtusis, flores in singula circiter viginti. Pall. l. c.

Caspica & Salicornia articulis cylindricis, spicis filiformibus Wild. Geopfl. p. 26

halit. arborescens, tamarisci foliis Burm. Cent. 1. t. 11. f. 1.

Salicornia caspica Pall. A. 1. app. n. 90. t. 8. f. 2

Habitat in squalido mari Caspici et Medici.

Gigas in suo genere. Truncus fere arborescens: Articuli Ramorum articuli
cylindrici. Spicis pedunculatis, saepe 1-2 pollicares, of coris confectione cre
beris nigra brevitate. Pall. l. c.

Solcata L. foliis lineariibus alternis amplexicaulis. Decurrentibus Lin. Suppl.
p. 81. L. V. p. 53. Pall. It. 1. app. no 93. Tab 8.

Habitat in Sibiriā 5

Fructus cymosus. Brachi alterni subsimplicis. Folium
cartilagineo basi amplexicaule adnata quasi decur-
rentia.
natula

non vid.

arabica & Salicornia foliis alternis vaginatis obtusis linei dehiscentibus Lin. Suppl.

p. 81
Kal. arabum secundum genus ~~Salicornia~~ Salach: Lugd. app. p. 20.

Salicornia arabica Pall: it. l. app. n. 9. Tab. D. p. 3.

Habitat in Arabia 5

Stipulus via pedalis. Caules plures, erecti, cortice equali, rimoso albo.

Rami annui alternatim sparsi et subdivisi; articuli ovati superiore angulo

prominulis. Spicae in ramis exterminis, interdum plures, alternae,

parvae; ovatae, teretibus. Flores pauci in singula spica Pall. l. c.

† tres vel
quatuor

Scandens B. caule erecto, floribus dianthi, foliis cordatis acutis. Lin. Man. 315. f. V. p. Jacq. hort. t. tab. 4.

~~Solanum bacc. fenum americanum, fructu corymboso Plukin. alm. 349. f. 226. fig. 7.~~

~~Salicaria cuneifolia, fructu apice ^{apice} Plukin. alm. 301. f. 113. fig. 7.~~

Habitat in Jamaica 4

~~Baccharis scandens, foliis cordatis subrepandis, caule glabro scandente, floribus umbellatis verticillatisque, fructibus clavatis apice tuberculatis. Sw. obs. p. 124~~

~~apice tuberculatis~~

~~Caulis teres glaber. folia petiolata crassa subvenena rami alterni pedunculi axillares et terminales bifidi vel rami dichotomi pedicelli 6. 9. glandula in apice geminis~~

fructibus clavatis

10

Foliosa 10. Salicornia foliis linearibus alternis amplexicaulis Saururus
tibus Lin. suppl. 81.

Salicornia foliata Pall. it. 1. app. n. 93. t. 2.

Habitat in Sibiria, ad mare Caspicum. †

Trunculus pedalis: Rami annui, sparsi: articulis obovatis, su-
perius excrecentibus in foliis crassis, teretibus, obtusis, qua-
drifariam patentibus. Spicæ alternæ in foliorum axillis
vel internodiis, sessiles, sæpe uncialis, cylindricæ vel fusiformes.
Flores ^{scule} vix prominuli; areolæ tribus spathiis in an-
gulum positæ, quarum laterales tropæiæ, minores, anthers-
fera sessiles sustinent; mediæ pentagonæ, stylo centrato tri-
furcæ. An varietas Sal. arabicæ? Pall. l. c.

constant.

Classis 1.

Monadonia

monogynia

* Flores completi, inferi
+ inferi.

1. Lopezia Cal. 4-phyllus. Cor. 5-petala, inaequalis. Caps. 4-valvis, 4-locularis, polysperma
+ Boerhaavia varia ** Flores superi completi, superi.
2. Cucularia Cal. minimus, 4-lobus. Pet. 4. Duo majora, concava, altero basi corniculato: duo interiora, breviora, linearia. Anthera loculis discretis.
3. Usteria Cal. 4-dentatus: lacina unica reliquis major. Cor. infundibuliformis. Caps. ~~un~~ locularis, 2-sperma. Gemina anillata.
4. Qualea Cal. 5-phyllus; foliolo unico majore, basi glandula calcarato. Cor. 4-petala. Per. 2-locularis, polyspermum

11
perfoliata" Salicornia foliis alternis orbiculatis perfoliatis, ramis distichis

Salicornia perfoliata; fruticosa; foliis carnosis, unicolorato-orbicu-
latis, caule perfoliatis Forsk. descr. p. 3. Ic. tab. 1.

Habitat ad littora Maris rubri.

Diff. Caulis diffusus, teres, non articuletus; interdum crassitie pennis
ansens, pedalis, flexuosus. Volia conferta, vix semilungeta utaria
in juniori planta nondum frugifera vagina crassa unicolorata, circa cau-
lem decurrentia. Spicae ~~terminales~~ axillares et terminales
pedunculatae, saepe sesqui-pollicares, teretes, floribus ternationi positi

¹²
amplexicaulis Salicornia foliis cordatis amplexicaulis. Symb. bot. 2. p. 1

Habitat in Agro Turetano ©.

Caulis palmatis vel posum ultra, lignosus, decumbens, ramosis & ramosus.
Rami alterni, patentissimi perque pollicares, tibi semper approximati
si ex omnibus axillis foliorum folia parva, alterna, obtusiuscula,
subtus convexa, & supra concava planiuscula, in ramis junioribus
imbricata.

Habitus Carnoso, plerumque caule vel ramis articulatis. Flores spirato

orientale ~~Coriiform: foliis lineariibus angustis summitatibus floriferis subpa-
nuculatis pubescentibus La Mark P. Enc: pag 15
Habitat in oriente~~

non vidi