

Samara

1. *Cela S. corymbi* lateralibus, foliis obtusis
2. *conica* floribus sessilibus conglomeratis, foliis lanceolato ovatis acutis subinacuis
3. *pentandra* floribus pentandris, foliis ellipticis

156

Samolus rorarius

Cal: perianthium minimum. Strobilum folioli orati.

Corolla quadripartita ^{interdum 5, partibus} laciniis ovatis ^{ad} sicuti concavis, acutis antheris ferventibus

Stamina filamenta vis ulla. Anthera quatuor, sagitta basilari aequa, unica in singula cavitate


laciniis Stamina interdum 5.

germen superum parvum oblongum. Stylus longitudine Strobili germinis

Pericarpium duplo, magnitudini Pappi ² nigra, rugosa, monosperma.

montserrat Dr. Ryan

a ^{maxillae} flos magnitud. tubulari
 b — acutus
 c — ^{retroflexo} cum reflexo *Pedicularis foenic.*
 d petalum cum anthera
 e — anthera cum ^{nidamento} nidamento
 f nidament. pistilli
 g flos hermaphrod
 h pistillum
 i supra
 k. nec



Pedicularis foeniculifera

coniacca Samara floribus ^{sub} sessilibus conglomeratis aggregatis, foliis lanceolato-ellipticis acutis
Samara coniacca floribus sessilibus conglomeratis, foliis lanceolato-ovatis acutis
subconiacca Sw. prodr. pag. 32.

Habitat in montibus de Rya

Arbor polygama. Rami teretes, cortice fuscescente griseo, glabri, punctati, sub
dichotomi

Folia petiolata, parva, frequentia, bipollicaria, basi acuta, apice saepe ob-
tusiuscula, utraque glaberrima, supra obscure venosa lucida, subtus
pallidiora interdum arenia, vix coniacca, integerrima.
Peduncululi axillares corymbi, brevissimi.

1 Cota Samara Mant. 1994. V. p. 154

~~Mamestylon~~ umbellatum Burm. Ind. 87. ~~Zeyl. 76. t. 31.~~ Lin fl. Zeyl. 469.

Cornus sylvestris altera Zeylanica *sylvestris altera* Burm. fl. Zeyl. pag. 76. t. 31

Habitat in India Orientali

Rami purpurascens, laves. Folia alterna, petiolata, lanceolato-elliptica, ob
tusis integerrima, utriusque Coria. Cymbi laterales, axillares, vix petiolis
longiores. Flores copiosi lutei

Differt a Cornis non tantum germine supero, sed et filamentis petalo oppositi
nei alternis

Cota P. Cymbis lateralibus pedunculatis, foliis obtusis

Rami fusco-purpurascens, glabri, uti reliqua partes. Folia petiolata, alterna, pellucida
viri, lanceolato-oblonga, basi acuta, subserena integerrima. Pedunculi axillares, breves
multi flori herb. Guss.

Rami cortice cinere. Folia oblonga obscure nervosa, ^{rami} basi, pellucida pedunculi
axillares, oppositi, foliis breviora. Flores in apice congesti, plures. Germen inferum
calyce coronatum herb. Herm. Lin in biblioth. Lugdun

Hoc est Burmanni cornus etc: non vero Samara Cota Lin. oblitus forte post 40 annos plant
tam quam in herb. Herm. viderat eandem esse credidit. Specimen enim herb. Lin
gaudet foliis alternis multo angustioribus et omnino alia plant
Burm. nullo est mei. Mamestylis non vidi

2 conatae l. floribus sessilibus conglomeratis, foliis lanceolato-ovatis
acutis subcoriaceis Sw. prodr. p. 32.

Habitat in Jamaica

Indivisa alia mascula, alia foemina. Pistilli incrementum minimum in
masculis videt Richard

Diff. vide inter
amer. & hanc

3 pentandra L. ^{lobus} foliis pentandris, foliis ellipticis H. Kew. 1. p. 160.
Habitat ad Cap. b. p. 1/2

non vidi

rapanea ^{Lenora} Rapanea floribus pentandris, foliis oblongis obtusis.
Rapanea guianensis Aubl. fl. Gu: pag. Tab 46.
Habitat in Guiana

~~Rapanea guianensis~~