

# Serpicula

1. verticillata L. foliis verticillatis aculeato-<sup>Canari-</sup>seriatis.
2. callitricoides L. foliis oppositis integerrimis lanceolatis
3. repens L. foliis oppositis alternisque

repens f. foliis alternis linearibus Lin. Suppl. p. 416. Mant. 124. ~~+~~ L. v. pag.

840.

*Serpicula serpyllifolia* Burm. prodr. p. 26. ~~et caulis adspersus setis nullis vel~~  
~~caulis f. punctulato armata~~ ~~caulis f. punctulato armata~~ ~~caulis f. punctulato armata~~

Habitat in Q. bono speciei

~~caulis f. punctulato armata~~ ~~caulis f. punctulato armata~~ ~~caulis f. punctulato armata~~

Caules digitales prostrati. Folia lineari-coniungentia alana flore

margine aculei minuti axillares sespales, et minuti  
ferrata

*Laurentbergia repens* Berg. in fl. Cap

verticillata L. foliis verticillatis aculeato serratis Lin Suppl. pag. 248. 416.

J. V. p. 248.

Habitat in India orientali

<sup>Toto glabra</sup> Caulis siferms ramifus <sup>veretupulus. Rami oppositi glabra</sup>  
<sup>oblonga mucronata</sup>  
subtilissime aculeato <sup>folia quatuor vel pna sessile acuta ut semperguiculana</sup>  
<sup>in ramis internodiis confon.</sup> serrata <sup>glabra, mucronata. Pedunculi axillares, capillares, sesqui-</sup>  
pollicares uniflori

Callitricoides L. foliis oppositis integerrimis lanceolatis

Habitat in India orientali in agris

Caules filiformes simplicissimi ad gemmula radicans

Palmaria folia lanceolata sessilia opposita semunquiuila

ner <sup>angusta</sup> superiore conferta acuta. Capula axillares solitariae

subrotunda <sup>folia vero diversa</sup> marginata opposita sessiles

Habitus Callitric. autumnalis an eadem, sub nomine L.

C. verticill. ex India meso, omnes autem diversa  
de genere dubito