

Sherardia

*Gratiola* *Sheareria* foliis quaternis aequalibus Len. Spec. pag. 149\* f. V. p. 148

Habitat in Insula Adfensionis

Fit ex toosis, inaequalis, scabra: ramis obtusis tetragonis. Folia quaternae angustolanceolata, integerrima, glabra, marginibus revoluta, articulis ramorum paululum longiora, interstincta. Stipulis brevissimis, subulatis. Flores oppositi, sessiles, axillares. Calyx tetraphyllus germi inaequalis, persistens. Perleolis lanceolatis. Corolla infundibuliformis, quadrisida, alba, longitudine germanis. Antheris oblongis. Pericarpium diducum L. sp. l. c.

arvensis *Siberardia* foliis omnibus verticillatis, floribus terminalibus L.S. V. 140.

*Rubra* arvensis repens, *Conulea* Bauh. Pin. 149. pin. 334.

*Rubra* parva, flore conules se hargens. Bauh. h. 2. p. 719.

*Appendula* flore carneo, acuto folio Barrel. 9c. 541.

*Siberardia* foliis senis, lanceolatis, floribus sessilibus, umbellatis Hall. h. 734.

Habitat in arvis Europae septentrionalis O.

Caulis plures, humiles, ramosi. Rami tetragoni <sup>eterni veluti villosi</sup> glaberrimi angulis vix scabris. Folia  
verticillata, <sup>inferiora 4 perispermia</sup> sessilia, lanceolata, <sup>superiora 2 perispermia</sup> equalia <sup>ramorum adpressi</sup> ~~duo~~ <sup>duo</sup> ~~caena~~ <sup>caena</sup> margine que  
sunt retrosum <sup>supra</sup> ~~caena~~ <sup>caena</sup> serrata, <sup>supra</sup> ~~caena~~ <sup>caena</sup> ~~si abra~~. Flores in apice ramorum congesti  
sessiles, involucrio sessido.

