

Silphium

pumilum Siphium.

Habitat in Cardena herb. Ruth.

Caulis herbaceus, teres, erectus, superne canescens. Folia inferne pectinata, superiora sessilia, pollicaria, ovata, obtusa, dentata, supra cava, subtus subrepanda, subconescentia. Pedunculi superne, uniflori. Calycis foliola oblensa. Corolla medicina.

integripes Siphium

Habitat in regione Illeonor herb. Ruth.

Caulis herbaceus; tetragonus, striatus; scaber. Folia subconnata, bipollicaria et
ultra, ovato-lanceolata, acuta, supra avenia, enervia, margineque puncti-
slevati scaberrima, subtus venosa, scabriuscula. Pedunculi terminales
terni, intermedius uniflorus: laterales diphylli, triflori, adpressi foliolis pinnu-
loso-ciliatis, supra scabris. Calycis foliola ovata, acuta, reflexa, ciliata.

compactum Silphium

- Habitat in Carolina herb. Nut.

Caulis herbaceus, erectus, striatus, glaber, superne in ramos floriferos divisus. Folia petiolata, inferiora pinnata: pinnae alternae inferioribus subpetiolatae, superiores minores, confluentes, margine pilos-ciliata, venosae, dissectae: lacinae lato lanceolatae, remote inaequaliter dentato-subpinnatae; superiores pinnatifidae, remote inaequaliter grosse serratae, sub pedunculi parva, brevia, lanceolata, integerrima. Pedunculi elongati, ramosi. Calycis foliola oblonga affinis laciniato.

Nanum Salspiceum?

Habitat in monte Sidao Commesor herb. Turf.

Caulis villosus, teres. Folia sulphurea, semipollicaria, ovata, tri nervia, cin-
nis utrinque 2-3 in medio folii, interdum obscuri, acuta, ascendens, puberula
villosa. Pedunculus terminalis, solitarius, uniflorus, superne tomentoso-
incanus, elongatus. Calycis foliola pauca, rotunda; duo extrema mayo-
ra, reliqua involucentia
an huius generis, habitus convenit

Lacinatum Siphium foliis alternis pinnato-foliatis Linn. Saeser. 1. t. 1.
L. V. p. 704 - Linn. Spec. pl. pag. 1301*

Habitat in America septentrionali, Mississippi &

Caulis bis humano altitudinis, crassitie pollicis inferne loris
superne tuberculis fistulis pilisque natis albis pilis & c.

Folia alterna petiolata bipedalia ^{loba marginibus velis pubescent}
lanceolata vel ovata ^{loba marginibus velis pubescent} ^{loba marginibus velis pubescent}
~~loba marginibus velis pubescent~~ ^{loba marginibus velis pubescent} ^{loba marginibus velis pubescent}

Lacinis pinnatis

~~loba marginibus velis pubescent~~ ^{loba marginibus velis pubescent} ^{loba marginibus velis pubescent}
lanceolatis integerrimis vel uno alterove laciniis vel
natis see dentibus glauca suprema indurata. Soliola
calycis ovata ^{obtusata} acuta ciliata velis est densa longior
Cone fere intenorin

Theranthinaceum. ^{dent. serratis} foliis alternis ovatis serratis scabris: radicalibus cordatis

den. Suppl. p. 303. * l. v. p. 789

Siphium Theranthinaceum Gaerq. hort. v. 1. p. 16. t. 43.

Habitat

Caulis erectus simplex superne in pedunculatos. Divisus glaber
angulatus. Folia radicalia long. petiolata, scabripula
pithamea. ^{vertibus glandula supra} floralia ad basin pedunculorum lanceolata
foliola calycis ovata glabra long.

perfoliataim J. foliis oppositis deltoidibus petiolatis perfoliatis, caule tetragono
Linn. Spec. pl. pag. 1301.

Habitat in Mississippi &

Caulis erectus, ~~fulcatus~~ ^{radice} ~~angulatus~~ ^{et lignis cordatis} ~~homine~~ ^{albor} tetragonus folia
suis laevi. Folia ^{opposita} ~~opposita~~ ^{petiolata} ~~petiolata~~ ovata per petiolos
decurrentia ~~et~~ ^{obovata} ~~ovata~~ ^{imbricata} ~~dentata~~ ^{dentibus glandulosa}
oblonga obtusa terminatis. Petioli mediante folio decurrente
se connata. folia ^{suprema} ~~suprema~~ ^{folia} ~~folia~~ ^{se} ~~se~~ ^{subint} ~~egerrime~~
pedunculis ex axillis ^{supremis} ~~supremis~~ ^{et} ~~et ^{terminalibus} ~~terminalibus~~ ^{longatis}
simplicibus vel ^{subramosis} ~~subramosis~~ ^{foliicola} ~~foliicola~~ ^{calycis} ~~calycis~~ ^{ovata} ~~ovata~~ ^{glabra}
laevi, acuta~~

ovatis

connatum - foliis oppositis sessilibus persfoliatis, caule tereti scabro lin. Manis

p. 574. t. V. p. 709

Habitat in America 2.

Caulis altitudine hominis crassitie pollicis erectus simpliciter
tereti scaber (inferne obsolete tetragonus, scaber pilis deflexis
folia opposita connato-persfoliata; ovato-oblonga disco
concava, sessilia, basi petiolata, utraque persfoliata scabra
pyramide, vertice parva suprema floralia distincta
ter. Panicula terminali dichotoma: pedunculis pilosis
dichotomis solitariis unifloris elongatis
Calyx squamifolius: foliolis ovatis obtusiusculis lobis apice
paucis reflexis

Trilobatum L. foliis oppositis sessilibus cuneiformibus Lin. Am. ac. 2. 4. p

3074

Phryganthemum palustre repens minus odoratum, foliis cuneis trilobatis
Stoan. hist. 1. p. 262. 7. 155. fig. 1. Platan. phyt. 2450. fig. 4.

Corona folii maxima humilis, foliis cuneis trilobatis, flore luteo
Plum. frui. 10. u. 107. fig. 2.

Habitat in America

Caulis longi ~~prostratus~~ ^{repens} ter gem plicati angulati
pilis raris adpressis gemulis, saepe infractis. Folia oppo-
sita interdum subsessilia ~~oblonga~~ ^{rhombica} oblonga vel ovata
paulatimque medio lobo oblongo vel ovato acuta dentata
fiabra interdum coriacea. Pedunculus primum terminalis
dein axillaris solitarius elongatus uniflorus apice pilosus
foliola calycis oblonga pilosissima

variat quoad folia

videtur species haec

rhombica in dissepiti trilobique

Trifoliatum s. foliis ternis Linn. s. V

Chrysanthemum virginianum, foliis oppositis ternis s. quaternis adgeniculatis
Flos Monf. h. 3. p. 24. f. 6. 4. 3. fig. 60.

Habitat in Virginia 4

Caulis erectus angulatus glaber superne in pedunculatos
Dissipis folia petiolata ternis in culta interdum 4 na
inferior superiora opposita suprema alterna inter
gemma, omnia sabernna lanicolata. Foliole
calycis ovata lora glabra acuta

affinis L. foliis in visis seplibus oppositis: in fernibus alternis
Lin. l. v.

Affenfius coronis folio et facie Dill. Elth. 42. t. 37. fig. 42.

Habitat in Virginia, Carolina &

michi haud nota Specimen herbarii
Dubium

Scalium S. foliis alternis lato - lanceolatis serratis scabriuscula
tis subsessilibus Walt. Car. pag 216
Habitat in Carolina

non vidi