

Sloanea

Pat. 1. phyllus. Pericarp. ciliatum 3-6 loculare, 3-6 valve

1. Masfoni L. foliis cordato-ellipticis, stipulis linearibus: calyce quinque
partito, fecti capsula longissima
2. dentata L. foliis cordato-ovatis denticulatis, stipulis gematis
3. Jenerman L. foliis subrotundo-ovatis integerrimis, capsulis ovatis setosis
enjo. apice dehiscentibus

humilis

humilis detinetur inter amara inventus



a




c



d



b



e



f



g

Scoanea minor

minor Roanea foliis oblongo-ovatis integerrimis, stipulis integerrimis, calycibus
pentaphyllis, antheris terminalibus.

Habitat in Montserrat in Ryan &

Rami teretes cum adnatis tuberculatis, rugosi, glabri.

Folia versus apices ramorum, petiolata, alterna, spithamea et
ultra, oblongo-ovata, basi vix cordata, integerrima, acuta vel
acuminata basi terminata, glabra, supra secundum nervos lineata,
obscure venosa, subtus pallidiora, costata, venosa, venis exstantibus

Petioles bi-3. tripollicares, teretiusculo, stipula lanceolata, brevis, deici
dua integerrima.

Pedunculi infra folia, pollicares vel ultra, saepe plures, simplices vel
nova species
nunc. in litt. ramis simpliciter ramosi. Pedicelli ^{lemi} unguiculares, vix amplexiflori
Cinctus 6. superiores n. florae minores triplo fere minores quam in dentata. Bractea ova
opposita
do, concava, obtusa, brevis.


Calyx pentaphyllus: folioli aequo ovatis subsaequalibus

Corolla nulla.

Stamina filamenta copiosa, calyce longiora. antherae oblongae,
terminales.

Styllum german ovatum. Stylus filiformis, 4-5 partitus stigmata
acuta

Capula magnitudine nuclei juglandis fere, tota fetis mollibus concavo fimo
obfusa, his abrasis oblonga, tetragona, quadrilocularis



Sloanea dentata

calycis parum longiora. ^{ultra antheras in acumen producta.} Anthera filamenti ad caetera filamenta.

Germen ovatum depresso nectario impositum, vellosum, apice in stylum attenuatum
Stylus bifidus; laciniis capillaribus erectis. Stigma acutum.

(An *Sloanea dentata* Plumieri? differre videtur foliis subintegerrimis, contra
nit vero flore et stipulis, quantum ex figura Plumieri ~~videtur~~ apparet stipulae
non sunt cordato triangulari ut Venet. Descriptio Swartzii si mea eadem
ac Plumieri potius nomen sequens forte magis conveniat

dentata Sloanea foliis oblongo ovatis, stipulis lineari lanceolatis serratis,
calycibus subnovefidis, antheris lateralibus

1 *Marstoni* L. foliis cordato ellipticis, stipulis linearibus; calycis quinquepartiti
peti capitulo longissimis Sw. prod. pag. 82.

Habitat in Cambis 5

Rami teretes, glabri, apice villosi. Folia petiolata, alternis
utroque glabra, integerrima, supra secundum nervos
lineata, subtus costata oblique venosa. petiolis nudi
transversalibus striati. Pedunculis laterales ramosi pollicares

~~non~~ non vidi descriptis ad *Sinemarcesium* pertinere

2 Dentata L. foliis cordato ovatis denticulatis, stipulis serratis hinc sicut
pag. 730

Plounea amplis castaneo foliis, fructu subnatis Plum. gen. 49. u.

244.

Habitat in America meridionali

Folia ad apices ramorum petiolata semipedunculata utraque
glabra nervosa costa venosa, venis minoribus obliquis inter
maiores, remote denticata. Pedunculi ad apicem congesti
filo hypoleucana. Strobili saepe 3. 5 in singulis pedunculis
alberrimi pedicellati. bractea subulata ad basin petalorum
pedicellorum. Petiolus uti stipula pubescenti. Stipula
subulata serrata unguiculata

varietatem an speciem diversa a de Lund foliis ovalibus denticatis
basin non cordatis

3 Guineensis f. foliis subrotundo ovatis integerrimis, capsulis covatis setosis
ex apice dehiscentibus Sw. prodr. p. 82.

Stoanea Guineensis, foliis subrotundis integerrimis; fructu par
vo aculeis ⁹³⁴ ~~pepido~~
Aubl. fl. Guian. 344. t. 212.

Habitat in Cambois, Guana f
arbor folia ampla rigida, glabra saepe emarginate petiolo
ta. Sepalo oblongo, flores axillares in racemulis ^{foliolarum} pedun-
culo ad basin squamula minuta flores minores quam in
dentata Aubl.