

Sphenoclea

<sup>rotata</sup>  
Cordila ~~tubulosa~~ . Capsula biculani<sup>2</sup> polysperma circumcisa

spicata L.

Spicata Sphenoclea

Pangetti Rheed. Mal. <sup>pag 47</sup> ~~44~~. Tab. 24

Habitat in aquosis Indiæ Orientalis König

Caulis herbaceus erectus, teres pedalis sursum versus sensim attenuatus  
glaberrimus, ramifus, Rami aeterni patentes <sup>patentissima</sup> fructura caulis.

Folia petiolata, alterna, sesqui-pollicaria, canaliculata, obtusa, integer  
rima, glabra, tenera simplicitate venos

Spica terminalis ex apice caulis et ramorum, pedunculata, ovato  
oblonga, pollicari obtusa.

Sacie accidit ad Phytolaccam

*Sphenoclea* Gaertner de fruct pag. 113. Tab. 24. fig. 5

Calyx Perianthium monophyllum quinquefidum laciniis subrotundis  
Corolla monopetala, tubulosa, quinquefida <sup>longior</sup> calyce <sup>paucis</sup> longior quinquefida  
laciniis subrotundis

Stamina filamenta vix ulla, Anthera 5 versus basin corollae globosa

Pistillum germe inferum stylis breviteris stigma simplex <sup>per</sup> convexiusculum  
style parum latior

Sphenoclea

zeilanica Sphenoclea Gaertn: de fruct: 1. pag. 113. Tab. 24. fig. 5.

Gaertnera pangati. Retz. obs. 6. pag. 24.

~~Pangati~~ Pangati Retz. hort. Mal. 9. Tab. 24.

Habitat in India orientali

Caulis herbaeus erectus; uti tota planta glaber  
ramosus. Folia breviter petiolata, <sup>puberula</sup> ~~la~~ alternata lan-  
ceolata, integerrima bipolliciana. Pedunculuster-  
minales cum ramo ulteriores exirespente ungui-  
culari astate longior. flores seplei in capitulo oblon-  
go. Corolla rotata alba

Caunifolia Sapindus <sup>glabri</sup> inermis foliis pinnatis: foliolis ovato lanceolatis attenuatis  
~~per~~, petiolo simplicibus, panicula terminali.

~~Boerhaaf f. iter Rhed. Mal. 4. pag. 43. Tab. 19~~

~~Sapindus trifoliata foliis ternatis Lin. Sp. Pl. pag.~~

~~Rami teretes striati, glabri.~~

~~Folia petiolata alterna pinnata cum pan. p. tripega. Folio la  
subopposita breviter petiolata, <sup>et 5 pollicaria, infra menta</sup> ovato lanceolata vel elliptica lance  
olatis attenuate indegerrima, acutiuscula, utraque glabra,~~

~~Petiolus <sup>inter se</sup> altero breviora <sup>et 5 pollicaria, infra menta</sup> universalis teretiusculus  
et loci in racimo supra axillam propositus. Radiculi et pedicelli subpubes  
centes. Bractea ovata minuta.~~

~~Calyx, corolla stamina et anthera compacta~~

~~(Cur trifoliatam auctoribus hanc dixerint <sup>per</sup> facile non liquet, cum foli  
pinnata uti et monstrat figura et descriptio Rhedii l. 6. nomen tri  
te igitur ut pote quidem maxime incongruum mutari~~

*Spicata Sphenoclea* Mal. 11. pag. 24. Kozb. 24.  
Panzabichius

*Gaerthneria pangatti* Petr. ob. 6. pag. 24.

*Sphenoclea zeylanica* Gaertn. de fruct. pag. 113. Tab. 24. fig. 5

Habitat in aqua in India

Caulis erectus, ramosus, teres, glaber, ubi tota planta. folia petiolo  
ta, alterna, lanceolata indeguerrima obtusifolia, bipinnatifida  
ria. Pedunculus terminatus erectus, solitarius. Flores conferti  
in spica oblonga