

Stative

Catipalea Strobilifera paniculato ramifera, foliis pubescentibus
pilis fasciculato stellatis. Len. Trans: 1. p. 252.

Habitat in Siberia 2

Solia pedalia oblonga, panem undulata, mollia emarginata. Scapus
pubescens. Pedunculi teretes. Bractea opposita concavo subrami
ficationibus.

acerosa Statice. foliis linearibus subulatis rigidis pungentibus
scapo simpliciter spicato. Willden. L. C.
habit. galacia

Caulis fruticosus basi ramosus palmatis usque spitho-
mum. folia alterna sessilia conferta patentia
linearibus subulata basi parum dilatata vaginantia
margine scabra, apice pungentia, glauca. Scapus semi-
pedalis erectus floribus purpureis alternis sessilibus
bracteâ ovata acuta margine membranacea suffultis

Plantaginoides *Habia* foliis cuneatis glabris, scapis simplicibus erectis, specu Ver-
menali cylindrica.
Habitat in oriente.

Folia radicalia, cuneata integerrima, arenosa, enervata, crassa, fusca, ope lentis
tuberculata ceterum glabra. Scapus digitalis vel ^{in stamens} ~~parum vultu~~, inferne Vesic; super
versus presum crassior, striatus angulatus ^{supra supra ovata} ~~pitatus~~. Specia Vermenalis sesquipedalibus
flores compacti glomerati glomeruli ^{inferni} ~~inferni~~ duo vel tres parum remoti, bractea
ovata parva suffulti. Calyx ^{& laciniae parvae} marginibus membranaceo hyalinis, dentibus sive lacin-
is. Corolla incarnata s. fida tubo longitudine calycis, laciniis erectis oblongo obtu-
satis. Stamina sive corollae breviora, germina oblonga. Stili sive

Specia ^{longe} ~~etiam~~ plures cylindrica
folia integra vel fimbriata

Statice

Limonium maritimum fruticosum L'herit. Boiss.

Habitat herb. Tournef.

Radix superne in Caulis plures densis, basi ramosi & reliquis foliorum
squamosi. Folio pubescentia.

herb.

Statice

Limonium minus bellidifolio, flagellis foeniculaceis Boiss. in f. 2. p. 143.

Tab. 103.

Habitat herb Tournef.

habes

Statice

Linonum valentinum Statice folis capitulo glomerato
Habitat in regno Valenti. herb Tournef.

habes

Capillacea. *Statice*

Statice capillacea folio major Tournef.

Habitat in Luptonia herb. Guss.

Caules breves, herbamofi. Folia frequentissima, confertissima, filiformia, basi dilata-
tato membranacea, ferruginea, supra plana, subtus convexa, angustissima, ovata
armata villosula. Scapi pedales, erecti, glabri, teretes, laeves. Squama externa
ovata, interna oblonga, margine lato membranaceo cincta.

Stachis

Stachis lusitanica capillaceo folio major Tournef. f. v. herb.
Habitat in Lusitania herb. Mus.

habes

Salsa Stodice

Habitat in Hispania herb la nevell le Pausse

folia radicalia, lineari cuneata, scapi ramosissimi, flexuosi, dichotomi, teretes
conescentes

globularifolia *Statice*

Habitat

herb. Desfont.

Folia radicalia, cuneata, bipollicana, glabra, integerrima, acutiuscula, nervosa. Scapi teretes, 1-2 pedales, dichotome divisi, glabri. Spicae superne in dichotomis et apice, plures, pollicares vel minores. Flores solitarii, secundi, distantes. Squamae extremae ovatae, sequentes oblongae, margine membranaceae.

Statice

Limnium orientale humiliori canyophylli folio angustiore non aculeato, flore
spicato suavescente Tournef. Cor. pag

Habitat in oriente herb. Desfont

Caulis plures, brevissimi, alterni a superiore parte radice, inferne squarrosi
et squamis foliorum revolutis persistentium. Solia penipenna - Scapi digita
ler

Prathulata Habria

Habitat

Herb. Desfontaine

Folia radicalia, plurima, petiolata, uncinata, sesquipollucaria, obtusa, glauca, ob
scurè trinervia, integerrima, crassiuscula duo vel 3 in scapo petiolata. Scapus ve
res, sesquipeda^{lis}, glaber, inferne glaucius, alternatim ramosus. Squama ovata,
attenuata ~~fr~~^{sub}ramis. Flores ex apice ramulorum, secundi, alterni, distantes, p
ctari. Squama externo ovata, interiores oblonga

diffusa Stature

Inbitat

Herb. Desfont.

Radix superne divisa. Folia radicalia, linearia, glabra. Scapi plurimi, palmari, filiformes, teretes, diffusi, alternatim ramosi: rami semipollicares; distantes. Squamae ad ramos hyalinae, membranaceae, lanceolatae. Flores solitarii, pinnati, 3-4 in singulis ramulis, alterni. Squamae exteriores lanceolatae, albidae; interiores oblongae, ferrugineae, glabrae.

Descripta a Pourret in Act. Tolojanae

Statice

Limonum orientale, campophylli foliis acutissimo, floribus velut in capitulum
congestis, acaulis et calycis purpureis cetera Tournef. Cor pag 25
Habitat in Armenia herb Tournef.
Caules plures, caespitosi, digitales

habeo

Statice

Statice seu Caryophyllus maximus frutescens flore globo Gussley v. Ind. Luf.

Habitat in Lusitania herb. Tournef.

Caulis brevis, imbricatus squamis, a basi foliorum persistente, apice ramos nonnullos, crassitie penno anserina emittens; foliis vector. Folia basi in squama dilatata, rigida, 3-4 pollicaria; convexa, canaliculata, linea nervi integer omnia. Scapus et apice ramorum, spithamorum, veretes, bracte tubuloso semipollicaris ut in affinis. Stalysis squama lamina ovata, oblique Capitulum duplo major quam in armeni

Statice

Lemonium bellidii folio proliferum
Habitat herb. mus.

notes

Statice

Limonium hispanicum plantagineis foliis bullato Tourn. Suppl. v. herb.

Habitat in Hispania herb. musci

Scapus paniculatus. rami dichotomi

Statice

Limonium creticum juniperifolium Tournef. Cor.

Habitat in Creta herb. mus.

Leptogramon

habe

Statice

Limonium orientale, humile canyophylli folio angustiori non aculeato, flore
spicato, siccasse rubente Tournef. Cor.

Habitat in Oriente herb. Masf.

Caulis semipedalis, ramifus: rami sparsi, dichotomi.

Hetia

Limonium gracium bellidis folios minorum et lanceolato Tournef. Per.
Habitat in graecia herb. mus.

habes

Statice L

^{minimum}
Limonium gracum foliis hysopii hirsutis Tournef. Cor.
Habitat in Graecia herb. mus.

herb.

Statice

Limonium græcum minus ozymifolius Tournef. Cor pag.

Habitat in grecia herb. Mus.

Radix superne ramosa.

habeo.

aphylla Hartia

Habitat in Sibiriâ herb. Turf.

Scapus erectus, flexuosus, ramosissimus, teres. Rami suspensivi, dichotomi. Stipula ovata,
acuta. Bractea nulla. Spica bifida. Flores copiosi, parvi. Squama oblonga, obtusa, arida
alba, uti calyces. Calyx squesfidus.

Statice

Limonium minimum foliis hysperi pubescentis est insula Tragonisi Tournes. Cor.

Habitat in oriente herb. Tusc.

Caulis statim a basi ramosus, foliosus, vertus raris foliorum petiolorum, spithamosus.
Folia cuneata, unguiculata, basi in petiolum attenuata basi dilatata, angusta,
utrinque punctis elevatis notata, obtusa. Scopis dichotomis, scabriusculis et punctis
similibus foliorum frequentissimis. Spica terminali. Squama lanciolata.

Statice

Statice capillaris folio minor Tournef. F. R. herb

~~Statice~~ Habitat in Lusitania herb. Lus.

Folia medicinalia, setacea, pollicaria. Scapi Digitales vel parum ultra. Capiti-
fulum ut in Armenia

prolifera Statice.

Limonium gracium fruticosum proliferum bellidis folio Tournef. Cor.

Habitat in graecia herb. Tus.

Caulis erectus pedalis ^{articulatus,} primum trichotomus dein dichotomus, teres: rami elevato punctati, scabri. Folia ad articulos, aggregata, crassa, spatulata, obtusissima saepe emarginata, aserna, enervia, oculo armato confertim punctata. Spicae superne ad articulos, aggregatae, ex his squamis lanceolatis ferrugineis marginariis.

grandiflora Statice

Statice Lusitanica fruticosa maritima magno flore Tournef. Inst. R. 44.

Habitat in Lusitania herb. Tusf.

Caulis teres, superne in ramos nonnullos divisus, foliis vertic. Folia linearia, basi membranacea ferruginea dilatata cuneata, demum teretia, incurva, glabra. Scapus erectus semipedalis, lavis, glaber. Calyx squamis ferrugineis imbricatus; exteriores ovatae, interiores oblongae, margine membranaceae.

Statice

Habitat ad fretum magellanicum herb. Gus.

Facies omnino armena, an modo varietas?

Folia radicalia, lanceo-nis, glabra, striata, 4-5 pollicaria. Scopi semipedales. Involucrum tubulosum, laciniatum: laciniis dentatis. Capitula aureo ferruginea.

Squamis lato-lanceolatis