

Hrychnos

Rouhamon lupi - genus

Calyx pentaphyllus ^{inferius} foliolis linearibus corolla brevioribus
Corolla monopetalala campanulata & fida laciniis ovatis acutis, pater-
tibus fauce villosa
Stamina filamenta fere nulla. Antheris & que subsessilibus inter laci-
nias corollae.
Pistillum germen convexum. Stylus l. corolla parum longior. Stigma
capitatum

Strychnos

articulato baccata ovata

Capsula magnitudine nucis

Coryli, oblonga, laxis glabra, tripartita vis
apice stylo persistens. semibus 2 minutis

numen
triloculari
in

valvulis equali a basi ad apicem scidentibus Valvula equali

sepe iterum exteras. Capsula interior stylo brevi Ingonis, luti

triloculari
in 6

Semen oblongum sub apice receptaculi pendula usque furculorum

ovata

biliciale brevi intus brevi superne pro utraque parvis compressis

sinuatum apice inferiori obtusiore magnitudine fere
lasi minori membranae subcartilaginea tecta obrotum

subuni 6
in

in pulpa involutum. obrotum magnitudine seminis
nuci atq. superne atum. tunica exterior subcartilaginea

Idem dicitur baccata
triloculari tunica
interior choriata

interior tenuis. albumen magnitudine et figura seminis
embryo radice alba fusca versu lineari cotyledonibus

triplo brevior cotyledones subrotundo una basi parum emargina

Galla oblonga unguiculari
apice acuminata
corolla

receptaculum longitudine capsula ^{triangulari} ~~triloculari~~ ^{superne}
setis supra parum crassiora membranae ^{ad rotum apicem} ~~triloculari~~ ^{superne} ~~triloculari~~ ^{superne}

superior

brachyactis Strychnos

Strychnos brachyactis foliis oppositi ovati oblongisque acuminati quaque
nervis, caule brachyactis terati, floribus corymbosis &c. Panus # 2. pag

30 Feb. 157. fig. 6.

Habitat in Peninsula nemoribus imis &

Caulis fruticosus, ramosissimus, scandens, fusco-cinereus Rami longissimi
ramosi, brachyactis, dependentes, granulosi, obtuse tetragon. Vicia oppo-
sita, breviter petiolata, ovata, oblonga ovaleque, acuminata, integerrima,
nervis, sive nervis, nervis lateralibus obsolete, supra nido-
ulnigine glabra, dense venosa, venis horizontalibus. Petioli connati,
sub contorti. Circhi raris & simpliciter, spirales, breves. Pedunculi
axillares, terminalesque, solitarii, corymbosi, multiflori. Pedicelli
breves, basi bractea linear-subulata, decidua. Calyx minimus penta-
phyllus, hirsutus, deciduus. Corolla alba, extus hirsuta, calyce qua-
druplo longior. Bacca magna, globosa, corticosa, fragilis, crassa ~~fragi-~~
~~li, cinerea~~ ~~tecta~~ viridi-lutescent, dependens, uniloculari, pulpa
lutea referta. Semina pedunculata, suborbiculata, compressa, pleneque
vane angulata, glabra, fulva, cornea, pellucida, crassa ~~fragili~~ ^{cinerea} tecta
&c. Panus - C. C.

acuminata *Arychnos*

Habitat in Guiana Richard herb. Guss.

Rami teretes, glabri. Folia petiolata, bipollidaria, lanceolata-oblonga, acuminata, glabra, trinervia, subtus nitida venosa. Flores non vidi. an *Nonhamon Aubletii*

Fraxinea Strigosa

Habitat in insula Bourbonia Commes. herb. Mus.

Rami teretes, oppositi, glabri, cinerei vel purpurascens. Spinis oppositis, unguiculatis, purpureis, patentibus. Folia breviter petiolata, pollicaria, oblonga, ovata, vel obovata, glaberrima, obtusa, mucronata vel subretusa, supra nitida, arenaria, quinque nervia, basi parum attenuata. Cymbrum terminale. Racemum oppositi villosi. Perianthia nulli. Calycis laciniis linearibus. Corolla squae dentata, faucibus hispida. Stylus 1. Balda sphaerica.

— Mus.

Spina Styracis

Habitat herb- La Martini

Rami horizontales. Spino axillares et terminales patentissimi.

nux vomica L. foliis ovatis, caule inermi Lin fl. reuel. 91. S. V

nux vomica officinarum Bauh. pin. 511. Raj hist. 1014.

Colubini ligni Vestrum genus in Malabar Bauh. pin. 301.

Canuram Rheed. Mal. t. p 67. 7. 37. Raj hist. 1661.

Habitat in India

Rami teretes, glabri cortice cinereo, adiacenti ^{in ramulis} ~~sed~~ articulis parum
compressi dilatati. Folia ^{in ramulis} petiolata, opposita, glabra
senesca integerrima basi subrotunda obtusa, ternaria
tri vel quadrifolia, pedunculati in apice ramulorum
solitarii vel bifidi ramosi quos corymbosi.

obtusis ternariis
— ovato oblongo ovatis, pedunculatis terminalibus ramorum

Colubrina L. foliis ovatis acutis, ^{trinerviis} cinnis simplicibus Len. S. V. am. ac. 2. p. 119

Coluber Clematis indica ~~purpurea~~ foliis luteis Bauh. pin. 301.

Fructus orbiculati major fusci striatus Bauh. pin. 405.

Modia Canum Rheed. Malabar. p. 10. t. 5. Rajah 1661.

Habitat in India

a priori non diversa contendunt Botanici: India

— pedunculis terminalibus

non videtur

rotatorum s. inermis, foliis oppositis ovatis petiolatis acutis paniculis
verticillatis Lin. Suppl. p. 140. & S.V

Habitat in montosis ad Madras. &

Rami oppositi glabri cortice cinereo folia opposita brevissime petiolata
ovata utraque glabra oblique ~~multinervi~~⁵⁻⁹ nervi sive duo super
ne duo inferne venosa. Lami cymulati ad articulos superiores
ramorum 2-4-6 verticillatimposita pedunculata parva pedun-
culis glabris. Flores nutantes. Corolla fauce vellis clausa. Bacca
globosa magnitudine Ceras.

pedunculis
foliis ovatis acutis sive nervis paniculis
catenulis subverticillatis

Spinoza *Stycknera* ramis spinosis, foliis obovatis, corollae faucibus barbatis calycis
sex longioribus. La Mart. T. E. 2. pag 30

Pluch. t. 170 fig 4.

Habitat in Madagascar.

Bacca globosa maxima. Semina globosa compressa margine subrecto

muti quatuor Agnoscant ad *Stycknera* revertat La Mart