

Trigonia

Trigonia

Character Essentialis Calyx quinque partedius. Petala quinque, inqua
lia; & iu premium basi intus foveolatum. Stamina ~~non~~ Nectarini
~~squamo~~ due ad basin germinis. Filamenta non sialla sterilia.
Capsula leguminosa, Trigona, trilocularis, trivalvis

97.4.8

Character Essentials

Mabes

²
Corri Trigonia foliis oblongis utrinque glabris nitidis ruscansis ciliatis Ver
minatae sanguine.

Trigonia corri foliis ovatis, fructu brevi, a fratre Aubl. fl. Guiana pag.
390. Tab. 150.

Habitat in Guiana Dr von Röhr 5

Rami punctis numerosis spinis elevatis inferne ad fructu ^{glabri} teretes; superne
compressi - subtriangulari vellosi sculi.

Folia petiolata, opposita, bi. f. triplicaria, oblonga, acutifolia, integrer
nima, utrinque glabra excepto costa, venosa, nitida, pubescens palli
ciora

Petoli semiunguisculares; pedicelli. Ripulo nulla.

Racemi ex astillis supremis et apice ramorum, erecti; fructi: Vermi
nales. Triplicares, bafi ramis nonnullis oppositis, astillares sim
plificantes, pollicares vel planum ultra. Flores dimidiis minores
quam in T. velloso, pleniusque solitarii, intemperie gemini, breve
pedicellati, extus canescentes. Bractea minuta ad bafi pedicel
larem.

Palycis foliola lanceolata, subcanescens

nectarium ut in fermentata

gel 100

18^o 30 Dr v. Röhr

Trigonia

Trigonia villosa, foliis obvolutis subtus tomentosis incanae

Trigonia villosa: foliis ovatis integerrimi, fructu longo rufescente aublet.
fl. Quan. pag. 180. Tab. 149.

Lctr. O. p. 53
In Payenna legit vva Rohr

Rami teretes, inferne glabri, superne villosi: ramuli incano-tomentosi.

Solia in ramulis junioribus, petiolata, opposita, bi-vel tripli-lacinia,
ovata, basi versus attenuata, e Costa parum ultra folium
excurrende macronata, margine parum reflexa, supra glabra,
in Costa tantum flavecenti villosa, secundum nervos lineata;
Subtus tomentosa, incana, venoso-reticulata, nervis et venis
elevatis flavecenti villosis.

Racemi ex ultimi astillis simplicis sumis, terminaler compositi:

rami oppositi, patentissimi, basi versus decrescentes.

Pedielli oppositi vel alterni, patentissimi, uti pedunculi flavi

tomentosi. Bractea duo getaceo ad basin singuli pedicelli

Polyt. quinque-partitus ^{lacinia} pentaphyllo: folioli ovatis, acutis, extus tomentosis, incanis, intus
glabris, unicum interius, relexis parum latiss., extus nivis in dorso
linea elevata vindi, corollam fere aequalibus

Corolla pentapetala: pedalem superiori reliqui laticeo, ex his basi gibbum, intus
e regione gibbi foræola marginibus villosa: duo lateralia parum longiora
lineari-cuneata, subfoliata, duo inferiora oblonga in medio latens in venis
versus marginem gibba

Nectarium squamo duo, rigidiusculo, minuti, rufescentes, glabri, utraque
ad basin germinis, basi altero marginat^a base parum connata.

^{re redit}
Stamina filamenta ~~decomposita~~ alterum latius germinis, primum in unum
connata, demum in duo corpora divisum, filamenta quinque in
angulo: ~~læcimis~~ ^{tribus} squalibet antheriferis; duo ^{bilobis} sibi mino-
ribus & veribus. Anthera oblonga, parva, subrotunda.

Pistillum. Germen ovalum, villosum, incanum. Stylus unicus.

Stigma planum

{ Cum filimenta non nisi rufescente germine in duo corpora discedant, sed his
forte erit Trigonio genus. Clasi monade lphaerum infere se