

Ulex

• / europeus Ulex

parviflora Ulea
aphylla

Genista caule aphylla, fructis comppositis, floribus lateraliibus
solitariis, calycibus bipartiti.

Habitat in Panamia

Rami profunde striati, sparsi, purpureo-pentes: ramuli alterni
est foliis hirsutis ramosus constantes apice, eti rami in qua
mam flavescens deponentes, virides, striati; versus apicem
viles ramicibus vix conspicuus.

Spirae laterales ramorum alterni, inferiores versus basi spinis
lateralis, superiores simplicies.

Solia nulla nisi squama ovata, parva, acuminata rigida
sub fragulo frumenta partiali

Pedunculi laterales in fructis partikulis breves, cinerei.

Calyx sericeus, bipartitus: lacrima superior brevis, ovata;
inferior oblonga, concava.

Ventilum subrotundum, est tuis sericeum. Canna bifida est ut
sericea. ala lancolata, glabra, canna dimidi breviora.

21 reticulata ~~foliis vernatis oblongo-lanceolatis pubescentibus fustis~~
~~reticulato venosi, racemi axillaris subfusilebus; legumi~~
~~nibus oblongis compressi Iw. prod. p. 105.~~

Habitat in Jamaica

~~Pauli volubilis angulatus, molles vellosus pubescens. Polea
petiolata: Poleola petiolata inter medium oblongo subrombeum
seque pellucens.
Lateralia exstis gibbosa etsi lanceolata omnia acuta utra
que mollia, sub supra subrugosa fustis reticulata pallidio
ra. petoli tomentosi. Stipulo orato. Racemi axillares
solidari foliis longioris. Pedicelli solidari, d' Hanter. Oalyx
quinquedidi laciniis subulato. Legumea oblongum utraque
acutum pubescens propositim margine utraque fusco. Semina
d' us~~

~~reticulata~~ ~~obtuso-oblongis subrombeis sub molles fustis~~
~~reticulato venosi~~

nanus L. Denthibus calycinis lanceolatis distantibus, bracteis minutis adpresso, rameulis
decumbentibus Smith. fl. Engl. 2. pag. 757.

Genista spinosa minor Reji Syn. 475.

Ulex europeus B. Lin Spec: 1045

Habitat in ericetis & latronibus ^{humilior}

~~long~~ ^{minor} europeo, rameis decumbentibus, fructu brevioribus, floribus minoribus:

Bracteis ex quo sope vix manifesta, ~~est~~ Smith

ulix

Habitat in huperca herb. Desforest.

genus & spadum lusitanicum lanuginosum aculeis tridentatis longioribus
munitum Fournet. f. v. hctb.

Rami pelosi, fraxi, per intervalla conferti, s'niati. Aculei pennati, apice
florescentes. Folia pectinata, attenuata, s'niata. Flores in apice, pedicellati;
pedicelli tomentosi. Calyx et Corolla glabra, longitudine squamis, magnitudi-
ne fere Ulii europeo.

Venofum Hedyotinum

~~Habitat in Basaria~~

Caulis rist. ullus. Solia radicata, 4-6 juga. foliola lineana & pilosa, ut
petoli, alata, semi-pollucaria. Si pula pubulata, hastata. 8 cores.

minor Ulmus pumila parkii, colycis foliolato inferiore videntata foliis
lineariibus glabris Roth Cod. bot. 1. p 83.

Nepo Theophrasti Lobel. f. pag. 788.

Genista pumila major, brevioribus aculeis Baeh. p. 394

Ulet europeus B. Lin

Habitat in Gallica

~~pendulina R. germinibus glabris, pedunculis caulegue hispidis, petiolis
inermibus, fructibus pendulis l. m. s. v.~~

~~Rosa sanguinolenta majori folio, fructu longo pendulo L. C. L. 6th
328. f. 245. f. 317.~~

~~Habitat in Europa~~

europaeus Ulex. Lin. S. V.

fl. Dan. t. 608.

Genista spinosa Dad. pempt 659;

Genista spinosa major, breviori longioribus aculeis Brauck. p. 394

3 Genista spinosa major, brevioribus aculeis Brauck. p. 394

Habitat in Anglia, Gallia, Brabantia,

romis spines que recte ^{tulicato} floribus laterali

parriflora

U ratris punci que floratis ex gloribus vestillis can
nique fricci

floribus ~~et~~ ^{terram} cum parris tunc solum quibus
parris

~~Calyx pentamerum per 5 apphyllum foliolis oblongis~~
~~Corolla pentapebala. Petalis oblongis longioribus calycis~~
~~Stamina. filamenta nulla Anthers 10. vel 8.~~