

Famia

angustifolia Zamia.

Zamia angustifolia Walp. coll. 3. pag. 263.

Habitat in insulis Bahamae sive 4
Radix in capite conicum obtus. 2. 3 unciali, post 7 annos magnitudine pugni
imbricatus squamis latis acuminatis exserti villosi. foliatis more felinum
adulto folia glabra nitida, inermia exserti villosa abrupte pennata. Stipes
ter. calamo duplo gracilior, inferne apophyllus 4, pedalis. Foliola utrinque a 4 ad
20. linearia simplicialia, 3 lineas lata, obtusa, alterna vel opposita, planis apice
fida, patentissima, subtilesque striatula, intermedia, apice papillis 3 bus
indistinctis. e pinnitate 4 unci: squamis unus vel alter, basi squamis,
lineari lanceolatis acuminatis, dextr. reflexis, apertis, erectis, 4 pollicibus
terminatur in stipulis pedice masculo in foemineo foemineo.

Cycadis 7. frondibus pennatis: folioliis lanceolatis acuto spinosi Len.

Suppl. p. 443. & L. V. pag. 926

Cycas castra Thunb. in Act. Ups. 2. p. 204. T. 5

Habitat in Cap. b. Spei

rumula ē frondibus bipinnatis: folioliis linearibus obtusis Len Suppl.
p. 443. P.V. p. 926
Palma americana, crassè rigidisque foliis Herm. parad. 210.
t. 210.

Habitat.