

Tannonia

indico 7

Zannonia Campelia

Compositae Zannonia Lin

Habitat herb Richard.

Caulis erectus, 2-4 pedalis, carnosus, teres, lavis, glaber, suffruticosus. Ramuli
floriferi erecti, terminati bracteis binis foliaceis, amplexantibus, contiguis, pa-
rentisimis recurvis, subundulatis, ad oras discoloris, pubescente, oblongo cordata,
Pedunculi infra post bracteam, glomerati, breves, uniflori, interstiti paleis
membranaceis, subdentatis, recurvis, glabris carnosis, sursum parum incrassatis
Flores cernui, exallidi. Corolla 6 petala ^{partia} lacinis 3 exterioribus ovatis, con-
cava, undulato albidis, superioribus majore, 3 interioribus amplioribus, tenuioribus,
candidis, subrotundo-ovatis, concavis, marginatis. Stamina 6, conformia
calyci inserta rursus breviora. Filamenta primum quasi plicata flexuosa, demum
rectiuscula, fide formia. Anthera connectivo litteram V. in resham fere referens
terminato utroque brachio loculo subrotundo. Germen subrotundum trique-
trum, apice submarginatum. Stylus primum plicato contractus. Stigma
excavatum Richard

India Zannonia L. f. V

Penas-valli mascula Rheed. mal. p. p. 39. 7. 49

Penas-valli foemina Rheed. mal. p. p. 31. 8. 47. 40.

Habitat in Malabaria

Florae racemose. Solia ovato lanceolata